



Unresolved gamma-ray point sources in the inner Galaxy at $E > 10$ GeV

Silvia Manconi

Laboratoire d'Annecy-le-Vieux de Physique Theorique (LAPTh)
CNRS (France)

September 14, 2023

TeVPA 2023, Naples (IT)

The gamma ray Galactic Center Excess (GCE)

Statistically significant excess in Fermi-LAT data [talk by D.Hooper]
few % of inner Galaxy flux

Debated nature:

Millisecond pulsar-like (MSP) in Galactic Bulge vs. Dark Matter (DM)





image credits: CLUMPY (J factor map), diffuse model [Storm+17], spectra [Gautam+21, Calore+14], morphology maps, [Storm+17], sim statistics [Lee+15]. *[Blame me for this collage]*

The gamma ray Galactic Center Excess (GCE)

Statistically significant excess in Fermi-LAT data [talk by D.Hooper]
few % of inner Galaxy flux

Debated nature:

Millisecond pulsar-like (MSP) in Galactic Bulge vs. Dark Matter (DM)



Characteristics: spectrum, morphology, photon statistics


image credits: CLUMPY (J factor map), diffuse model [Storm+17], spectra [Gautam+21, Calore+14], morphology maps, [Storm+17], sim statistics [Lee+15]. [Blame me for this collage]

The gamma ray Galactic Center Excess (GCE)

Statistically significant excess in Fermi-LAT data [talk by D.Hooper]
few % of inner Galaxy flux

Debated nature:


Millisecond pulsar-like (MSP) in Galactic Bulge vs. Dark Matter (DM)



Interstellar diffuse emission mismodeling

image credits: CLUMPY (J factor map), diffuse model [Storm+17], spectra [Gautam+21, Calore+14], morphology maps, [Storm+17], sim statistics [Lee+15]. [Blame me for this collage]

Focus: the GCE high energy tail



Crucial discrimination power: naturally explained by inverse Compton of e^\pm in MSP (see [\[GRA Monday session\]](#)), DM annihilation needs more tuning

Last dedicated work: [\[Linden+PRD'16\]](#) 7y of data using standard template fitting + photon count statistics (NPTF), likely same issues as [\[Lee+PRL'15\]](#) as pointed in [\[Leane+PRL'19\]](#) (see backup)

in collaboration with:


Francesca Calore (LAPTh, CNRS) Fiorenza Donato (University & INFN, Torino)

Assess GCE significance and measure characteristics of inner Galaxy gamma rays at > 10 GeV



- 1 Combining *state-of-the-art methods*, as pioneered in [\[Calore,SM+PRL'21\]](#)
 - * *Photon count statistics*: Measure faint point sources
 - * *Adaptive template fitting* : Minimize/study diffuse emission model systematics, crucial for photon count statistics
- 2 Preliminary results
- 3 Summary & Outlook

Photon count statistics of gamma rays

Statistical analysis of photon counts¹ to decompose the gamma-ray sky and measure source count distribution dN/dS below catalog flux threshold



1p-PDF



Main application: isotropic, extragalactic sources


Bright diffuse backgrounds + mis-modeling could bias method at low latitudes

[Leane&Slatyer PRL'19,PRD'20,Buschmann+PRD'20,Calore,SM+PRL'21]

¹Two main implementations: NPTF Lee+PRL16,Mishra-Sharma+AJ'17; 1pPDF Zechlin+ApJS'16,+ApJL'16 based on formalism introduced in [Malyshev+ApJ2011]. Main applications include: extragalactic sources [Lisanti+ApJ2016,DiMauro,SM+ApJ'18], blazar models [SM+PRD'20], DM halo, subhalo constraints [Zechlin,SM+PRD'18,Somalwar+ApJ'21], ...


SkyFACT: overcoming diffuse emission mismodeling

Model to fit Fermi-LAT data: Σ_{pixels} energy spectrum x spatial morphology



- Standard fitting techniques: up to 30% residuals [Cholis+PRD'22,Pohl+PRD'22]
- Mismodeling at low angular scales: *spurious evidence* for new components such as point sources [Leane&Slatyer PRD'20, Karwin+22]
- **SkyFACT** [Storm+JCAP'17]: **account for intrinsic uncertainties in spectral/spatial predictions** by introducing very large number of parameters w/ regularisation conditions for the likelihood [\rightarrow C.Eckner's talk]



Combining 1pPDF and SkyFACT



- **SkyFACT**: reduce diffuse mis-modeling
- **Photon-count statistic**: model faint sources after reducing residuals



First application to inner Galaxy [Calore,SM+PRL21, arXiv:2102.12497] at energies 2-5 GeV

Inner Galaxy, energies 2–5 GeV: results



- Stellar-bulge morphology *preferred over dark matter*: SkyFACT only (10σ) and modeling faint sources ($\ln B > 20$), confirms [Bartels+NA'18, Macias+NA'18, JCAP'19]
- *Unresolved* point sources resolved down to $\sim 5 \cdot 10^{-11} \text{ ph cm}^{-2} \text{ s}^{-1}$
- Diffuse mismodeling *strongly affects faint source reconstruction*

Inner Galaxy, energies 2–5 GeV: results




- Stellar-bulge morphology *preferred over dark matter*: SkyFACT only (10σ) and modeling faint sources ($\ln B > 20$), confirms [Bartels+NA'18, Macias+NA'18, JCAP'19]
- *Unresolved* point sources resolved down to $\sim 5 \cdot 10^{-11} \text{ ph cm}^{-2} \text{ s}^{-1}$
- Diffuse mismodeling *strongly affects faint source reconstruction*
- Faint sources *not purely isotropic*, few % of total 2-5 GeV flux

**Corroborating a possible, (at least) partial stellar origin
of the Galactic center excess**

Extending to energies > 10 GeV

Fermi-LAT data selection: ULTRACLEANVETO, all PSF (2-5 GeV was PSF3)




Extending inner Galaxy region of interest to 20×20 deg, cut plane at $b < |l|$ deg



SkyFACT results: inner Galaxy spectrum

GCE input spectrum: power law




Model components compatible between
full energy (0.2-500 GeV) and high energy (> 10 GeV) fit


work in progress: MSP-like GCE input spectrum

Stellar bulge of dark matter?

SkyFACT fit, nested model comparison




GCE: NFW ($\gamma = 1.26$)



Comparing to no GCE: 5.5σ
Comparing to boxy + nuclear
bulge: no evidence

GCE: stellar + nuclear bulge



Comparing to no GCE: 9.7σ
Comparing to NFW: 7.9σ

Photon count statistics, Bayesian model comparison $B_{ij} = \exp(\ln \mathcal{Z}_i - \ln \mathcal{Z}_j)$

no GCE

Comparing to no GCE:
 $\ln(B) = 17$


Comparing to no GCE:
 $\ln B = 37$

Comparing to NFW: $\ln B = 21$

unresolved sources + norm of diffuse, GCE templates ~ 1

**Evidence for a GCE at > 10 GeV
better described by a stellar bulge morphology**

→ Compare source count distribution and density to inner Galaxy





- High latitudes: stable and robust measurement, no matter diffuse model
- Low latitudes: large residuals with modA^2 drive spurious source reconstruction around catalog threshold
- Sources resolved at fluxes 1 order of magnitude lower with respect to catalog threshold, see e.g. 3FHL [DiMauro, SM+ApJ'18]

²GALPROP model from Fermi-LAT IGRB analysis [Ackermann+ApJ'15]

dN/dS results: control regions (10–300 GeV)

→ Compare source count distribution and density to inner Galaxy





- High latitudes: stable and robust measurement, no matter diffuse model
- Low latitudes: large residuals with modA² drive spurious source reconstruction around catalog threshold
- Sources resolved at fluxes 1 order of magnitude lower with respect to catalog threshold, see e.g. 3FHL [DiMauro, SM+ApJ'18]

² GALPROP model from Fermi-LAT IGRB analysis [Ackermann+ApJ'15]

Source-count distribution measured for 10–300 GeV:

20x20deg, cut plane at 1deg



- Gamma-ray point sources resolved down to $\sim 3 \cdot 10^{-12}$ ph cm $^{-2}$ s $^{-1}$
- dNdS reconstruction robust against modification of diffuse emission model
- *work in progress*: spatial distribution

Evidence for a stellar bulge-like Galactic Center Excess from $E > 10$ GeV Fermi-LAT data

- ✓ Addressing diffuse emissions mis-modeling with adaptive template fitting
- ✓ Measuring bright and faint point sources with photon count statistics

Work in progress:

- Spatial distribution of faint point sources in the inner Galaxy
- Consistency of results between energy bins
- Predictions for forthcoming surveys of high-energy gamma rays

Thank you for listening!



Dr. Silvia Manconi

silvia.manconi@lapth.cnrs.fr

Marie Skłodowska-Curie fellow

Laboratoire d'Annecy-le-Vieux de Physique Théorique, CNRS (France)

<https://silviamanconi.wordpress.com/>


I acknowledge the European Union's Horizon Europe research and innovation programme for support under the Marie Skłodowska-Curie Action PF2021, grant agreement No.10106280, project VerSi.

Backup

The dN/dS of gamma ray sources

Fermi-LAT source catalogs ³ \rightarrow source count distribution dN/dS

Table Browser for 1: gll_psc_v27.fits							
Source Name	DataR	RAJ2000	DEJ2000	GLON	GLAT	Conf. 68 S.	Conf. 68 S.
1	4FGLJ0000.3-7955	1	0.0963	-79.922	307.709	-42.7295	0.032278
2	4FGLJ0003.3+0743	2	0.1379	7.7373	181.056	-53.0295	0.049484
3	4FGLJ0001.2+4741	1	0.1126	47.6859	114.25	-14.3381	0.03688
4	4FGLJ0001.2-0747	1	0.1151	-7.7971	89.0327	-67.305	0.01844
5	4FGLJ0001.3+2113	1	0.3655	21.2183	107.646	-43.1477	0.02026
6	4FGLJ0001.6-4156	1	0.4165	-41.9425	334.226	-72.0285	0.042678
7	4FGLJ0002.1-6728	1	0.5378	-67.4746	310.085	-48.9635	0.021709
8	4FGLJ0002.1+6723C	1	0.5419	67.3578	118.203	-4.9349	0.069393
9	4FGLJ0002.3-0815	2	0.5537	-8.2652	89.0434	-67.8483	0.041752
10	4FGLJ0002.4-5156	2	0.6131	-51.9355	319.999	-63.539	0.094456
11	4FGLJ0002.7+7220	2	0.6786	72.3479	119.292	0.62628	0.06879
12	4FGLJ0002.8+6217	1	0.7261	62.2905	117.32	-0.051038	0.021154
13	4FGLJ0003.1-5248	1	0.7617	-52.8071	318.99	-62.7889	0.020537
14	4FGLJ0003.2+2203	1	0.8058	22.1922	108.439	-39.3888	0.040669
15	4FGLJ0003.3+2511	1	0.8323	25.1912	109.382	-36.4115	0.062968
16	4FGLJ0003.3-1928	1	0.8465	-19.4676	65.1978	-76.581	0.043964
17	4FGLJ0003.3-5005	1	0.8483	-50.0922	314.239	-56.9659	0.079319
18	4FGLJ0003.6+3059	1	0.9045	30.9888	111.004	-30.772	0.049955
19	4FGLJ0003.9-1149	1	0.9996	-11.6251	84.5954	-71.0751	0.032317
20	4FGLJ0004.0+5715	1	1.0017	57.2578	115.526	-5.02827	0.052945
21	4FGLJ0004.0+0840	1	1.0073	8.6757	103.496	-52.3978	0.050283
22	4FGLJ0004.3+4614	2	1.0757	46.2427	114.489	-15.8549	0.051805
23	4FGLJ0004.4+4337	1	1.1061	43.6233	323.886	-67.5378	0.027814
24	4FGLJ0004.4-6001	1	1.1184	-40.0251	336.991	-73.8453	0.083752
25	4FGLJ0005.6+6746C	1	1.4155	67.7688	118.608	5.26213	0.119523




(# of sources N per $d\Omega$ with integral flux in $(S, S + dS)$)


Limited by detection threshold of given survey

Characterizes gamma-ray sky composition and source populations


³ 4FGL-DR3, 12 years [Fermi-LAT coll., ApJS'22]






SkyFACT: gas smoothing scale





skyFACT: influence of smoothing scale of templates on dN/dS and residuals (2–5 GeV)




dNdS results: control regions (2–5 GeV)



dNdS results: control regions (2–5 GeV)




[Calore, SM+PRL '21]



Source count distribution 2–5 GeV: systematics

Stability of results tested against many systematics, see supplementary [Calore, SM+PRL'21]

Diffuse emission mismodeling




1pDPF modeling: flux cuts, dNdS breaks


Test of 1pPDF method with Fermi-LAT data

In 1-10 GeV

[Zechlin+ApJS 2016]




Adding a galactic dark matter template




To constrain blazar models

[Manconi+PRD 2020]



In > 10 GeV, with efficiency corrections


[DiMauro, SM+AgJ 2018]



Luminosity functions of MSP explaining the excess


[Dinsmore&Slatyer'JCAP'22]: Comparing 7 models for MSP properties explaining the excess:

sources detectable? # overall to explain excess? viable luminosity functions?



Number of detectable MSP highly depends on luminosity function model, which is uncertain

Factor 5-20 sensitivity: 30% of the excess would give detectable MSP in Fermi-LAT



Models can produce 100-10⁶ MSP w/o overproducing 4FGL sources

Photon count statistics: timeline [until 2020, using NPTF]




Is the excess diffuse or point-like (sources spatially distributed as dark matter)?

2016 : Excess entirely due to unresolved **point-sources** [Lee+PRL'16]

2019 : Earlier results not robust: smooth dark matter not reconstructed even if injected **dark matter strikes back?** [Leane+PRL'19]

2019/2020 : Explain why injection test failed: background mismodeling! excess still consistent with being **partially point sources** with updated diffuse models [Chang+PRD'20, Buschmann+PRD'20]


2020 : Preference for point sources influenced by spurious sources/ excess north-south asymmetry, robustness further casted into doubt [Leane+PRL,PRD'20]



Robustness of results highly dependent on serious systematics from Galactic diffuse emission mismodeling

Galactic diffuse mismodeling: residuals

Model to fit Fermi-LAT data: Σ_{pixels} energy spectrum x spatial morphology



Template fitting: still up to 30% residuals

Mismodeling at low angular scales, north-south: *spurious evidence* for new components such as point sources [Leane&Slatyer PRD'20, Karwin+22]


How do we reduce residuals?

Data-driven:


- Spherical harmonic marginalization
[Buschmann+PRD20]
- Gaussian Processes
[Mishra-Sharma, Cranmer, '22]
- SkyFACT: sky factorisation with adaptive constraining templates
[Storm+JCAP'17]

Improve models:


Better estimates of target H_I, H_2, H_{II} gas column density, inferred by line spectra, dispersion measures



- new atomic HI reconstr, with radiation model of emission + absorption [Shmakov+22]
- convolutional neural nets to fill gaps in molecular H_2 tracers like CO [Shmakov+22, Karwin+22]
- bayesian inference of 3D CO maps [Mertsch&Vittino'20]





Extended up to $\sim 10\text{deg}$ ($\sim 1.5\text{kpc}$)



Early works: spherical symmetric around GC, contracted NFW profile $\gamma = 1.26$

Recent debate: [DiMauro PRD'20,21, Pohl+ApJ'22, McDermott+22], ...



- Evidence for stellar bulge-like [Bartels+NA'18, Macias+NA'18, JCAP'19, Calore, SM+PRL'21]
- Longitudinal asymmetry at ~ 10 deg

Discriminate interpretation? Yes!

Morphology: still debated



Independent groups find different results for the morphology of the excess

Spherical symmetric (dark matter-like)

- Early works often *not testing* other morphologies
- Recent works: [DiMauro PRD'20,21, Cholis+PRD'22,McDermott+22], ... using astrophysical models and varying many parameters

Stellar bulge (MSP-like)

- Two independent groups in 2018 [Bartels+NA'18,Macias+NA'18,JCAP'19]
- Subsequent works with even more significance: [Calore,SM PRL'21,Pohl+ApJ22] (ring based, skyfact based)



Difficult to close this debate using only template fitting:
current residuals too high