

The X17 search with the MEG-II apparatus at PSI

Hicham Benmansour, INFN Pisa

ITN Intense Interim Review Meeting, November 23rd 2022

- Hicham Benmansour: From Paris, France - French-Algerian
- **Bachelor's Degree** in Engineering at Ecole Centrale de Lyon, France
- **Master's Degree** in Engineering at Ecole Centrale de Lyon, France



- **Master's Degree** in Physics at Queen's University, Canada



—> Master Thesis on DEAP-3600, dark matter direct detection experiment
—> studies of WLS fluorescence

- Since September 2021: **PhD** in Particle Physics at University of Pisa, Italy

—> PhD Thesis on the MEG-II experiment: hands-on work and data analysis



- **Particle Physics** - exam: July 4th
- **Instrumentation for Fundamental Interaction Physics** - exam: July 4th
- **Italian, A2 level** - exam: June 14th

Conferences and trainings


- International Workshop on Cosmic-Ray Muography, Ghent - November 2021
- 15th Pisa meeting on Advanced Detectors, Elba - May 2022
- International Conference on High Energy Physics XLI, Bologna - July 2022
- PSI Particle Physics Summer School - Vision and Precision - Zuoz (CH) - September 2022

1) The X17 anomaly at ATOMKI

2) The X17 search with MEG-II

3) Data collection and analysis


- Analysis procedure
- Gamma rate from BGO analysis
- Trigger rate estimate
- Optimized trigger
- Significance estimate
- First observables from data



1) The X17 anomaly at ATOMKI

The Beryllium Anomaly

Atomki Collaboration



${}^7\text{Li}(p, e^+e^-){}^8\text{Be}$ studied at
 $E_p = 450, 650, 800, 1100$ keV



→ e^+/e^- energy sum and angular correlation Θ

- Internal Pair Conversion (IPC) distribution shows excess at $\Theta \sim 140^\circ$ at several beam energies

→ decay of a light particle emitted during proton capture

→ best fit $m_X = 16.95 \text{ MeV}/c^2$
 $BR(X) = 6 \times 10^{-6}$

→ protophobic vector boson X17? mediator of a fifth force?




arXiv:2205.07744

Phys. Rev. Lett. 116, 042501

Phys. Rev. D 95, 035017

2) The X17 search with MEG-II

The MEG-II experiment



- MEG-II experiment searches for charged lepton flavour violating decay: $\mu \longrightarrow e\gamma$


Eur. Phys. J. C, 76(8):434

- 1 order of magnitude sensitivity improvement wrt MEG: $BR(\mu \longrightarrow e\gamma) \longrightarrow 6 \times 10^{-14}$

Eur. Phys. J. C 78, 380

MEG-II results from an intense upgrade program

Proton beam direction for X17 search



- The new MEG-II highly performing spectrometer can be used for X17-boson search:

→ X17-dedicated target in place of the muon target

→ gamma auxiliary detectors

→ MEG-II CW accelerator as proton beam


→ optimized TDAQ

→ adjusted magnetic field

3) Data collection and analysis

- X17 runs from February 10th to February 22nd: sample of **90 M** events
 - 10-17: LiF target (**55 M**)
 - 17-22: LiPON target (**35 M**)

- Signal



- Two types of backgrounds

IPC = Internal Pair Conversion
 → direct e^+/e^- pair creation

EPC = External Pair Conversion
 → gamma conversion outside nucleus


Analysis procedure

- Electron and positron tracking
- Example of reconstructed pairs

MEG reconstruction focuses on e+. It was adapted for e-. Performance was evaluated.

Positrons: +B search


Vertex Position - Y vs X



MC: 9 MeV e+/e-

Electrons: +B search

Vertex Position - Y vs X




5% relative contamination

$\theta \sim 90^\circ$

(orthogonal to beam) are tricky to identify


Positrons: -B search

Vertex Position - Y vs X



Electrons: -B search

Vertex Position - Y vs X





Mitigated by checking track correlation (+B/-B)

Reconstructed pair events from data

- Event A

$p_+ = 6.7$ MeV
 $p_- = 8.3$ MeV
 angle = 141°





LiPON data
02/22

• e+ hit
 • e- hit
 target

- Event B

$p_+ = 6.7$ MeV
 $p_- = 6.9$ MeV
 angle = 101°



Gamma rate from BGO analysis


Why a gamma detector?

- Understanding of background
- Stability monitoring
- Signal normalisation




- Two gamma detectors

→ Understanding of background → Stability monitoring → Signal normalisation



Bismuth Germanate (BGO) crystal matrix (4x4)




Lanthanum Bromide (LaBr3) crystal





- LXe calorimeter on maintenance during run



- ➡ BGO run (50k events) taken 1-2 times per day
- ➡ Crystals calibrated with F line at 6 MeV
- ➡ Sum on crystals
- ➡ Events with maximal energy release in the 4 central crystals
- ➡ Energy scale still to be corrected
- ➡ BGO 70 cm away from COBRA center → 4 central crystals: **0.14%** of full solid angle





Rate of gammas from Be15+18 = **16 Hz** (0.14% of full solid angle)

➡ **12 kHz** on full solid angle

➡ consistent with Brilliance rate

Assuming perfect efficiency and isotropic emission

Trigger rate estimate


- % of triggered events from MC
- Combination with BGO gamma rates
- Comparison with trigger rates from data

MC with gammas generated isotropically and uniformly [1,19MeV]


Combined trigger (1 SPX and 10&10 DCH)



#events triggered as a function of the generated gamma momentum



direction of gamma leading to trigger



→ **0.35%** of all 18 MeV gammas lead to trigger through EPC and Compton

→ **High energy** gammas induce trigger

→ Gammas convert through the **copper ring**

Estimated rates

- **12 kHz** (gammas on full solid angle)
- **42 Hz in trigger**

Raw rate from data

LiPON/ 2uA (#418137)
66 Hz

→ DAQ dominated by **EPC and Compton**: trigger needs to be optimized

Optimized trigger



A better choice of trigger for a potential next data taking

EPC MC: 200k gammas generated

<u>Trigger</u> 1 SPX hit n&n DCH hits	Triggered events (%)	#pairs in signal region
n = 10	0.35 %	0
n = 30	0.10 %	0
n = 50	0.025 %	0

EPC/Compton rate **divided by 14**
going from 10&10 to 50&50

X17 MC: 50k pairs generated

<u>Trigger</u> 1 SPX hit n&n DCH hits	Triggered events (%)	#pairs in signal region
n = 10	18 %	136 (1.5%)
n = 30	12 %	135 (1.5%)
n = 50	8 %	105 (2.6%)

X17 rate **divided by 2** going
from 10&10 to 50&50

Signal region:

- $16 \text{ MeV} < \mathbf{Esum} < 20 \text{ MeV}$
- $15 \text{ MeV} < \mathbf{InvMass} < 18 \text{ MeV}$
- $120^\circ < \mathbf{Angle} < 160^\circ$

Advantages

- ➔ Background largely removed from trigger while only losing 25% of signal
- ➔ Can be compensated by increasing proton current

Significance estimate

- significance from last data taking
- challenge: get best significance while keeping trigger rate below limit
- significance with optimized trigger

Trigger 10&10 and 2uA (I_{CW}) - prescaling 2 (last data taking)

- EPC + Compton:
6 kHz \rightarrow 21 Hz in trigger \rightarrow **0 pairs**
0.35% induce trigger *0% signal pair reco*

Total trigger rate: 24 Hz

All

- IPC18+15:
6 kHz \rightarrow 18 IPC generated \rightarrow 2.7 Hz trigger
BR 3e-3 *15% induce trigger*

- IPC18:
3 kHz \rightarrow 9 IPC generated \rightarrow 1.4 Hz trigger \rightarrow **5.4e-4 pairs/s**
BR 3e-3 *15% induce trigger* *0.04% signal pair reco*

**18 MeV
line**

- X-Boson:
3 kHz \rightarrow 1.8e-2 X17 generated \rightarrow 3.2e-3 Hz trigger \rightarrow **4.9e-5 pairs/s**
BR 6e-6 *18% induce trigger* *1.5% signal pair reco*

Significance

In 10 days:

- significance = $S/\sqrt{S+B}$ = **1.9**

For 5 sigma:

T = 71 days

\rightarrow to compare with data from last February (<1 sigma)

Trigger 50&50 and 10 uA (l_{cw})

All

- EPC + Compton:
60 kHz → 15 Hz in trigger → **0 pairs**
0.025% induce trigger *0% signal pair reco*

Total trigger rate: 20 Hz

- IPC18+15:
60 kHz → 180 IPC generated → 5 Hz trigger
BR 3e-3 *3% induce trigger*

→ increases by **x10** (l_{cw} and no prescaling)
→ decreases by **x14** (more constraining trigger)

18 MeV line

- IPC18:
30 kHz → 90 IPC generated → 2.7 Hz trigger → **4.5e-3 pairs/s**
BR 3e-3 *3% induce trigger* *0.17% signal pair reco*

- X-Boson:
30 kHz → 1.8e-1 X17 generated → 1.4e-2 Hz trigger → **3.8e-4 pairs/s**
BR 6e-6 *8% induce trigger* *2.6% signal pair reco*

Significance

In 10 days:

- significance = $S/\sqrt{S+B}$ = **5.1**

Observables from data


→ What were we able to extract from the data we have?

LiPON data - 2.1M events - 02/22 - All statistics from 22 hours of non-ZS LiPON data

Selection on:

- > *ngoodhits*
- > *vertices positions and distance*
- > *z position*


From CDCH track reconstruction




→ In pair events, e^+ and e^- energy can be summed to reconstruct the transition energies (including some energy loss)

LiPON data - 2.1M events - 02/22 - All statistics from 22 hours of non-ZS LiPON data

From CDCH track reconstruction



From BGO analysis




→ In pair events, e^+ and e^- energy can be summed to reconstruct the transition energies (including some energy loss)

→ Consistent with Be15 and Be18 gamma lines

Angular opening: data vs MC

LiPON data - 2.1M events - 02/22 - All statistics from 22 hours of non-ZS LiPON data



Assuming same angular opening distributions for IPC18 and IPC15

→ Data consistent with a mix of EPC and IPC (~55%/45%)

Summary:

- e+/e- tracking procedure was developed
- **data-MC consistent**
- from BGO analysis, **gamma rate estimated**
- with MC, **trigger rate understood** and optimized trigger identified for next data taking: **5 σ in O(few weeks)**

Next:




- characterize fake pairs

Backup

→ $\theta \sim 90^\circ$ tracks lead to **fake pairs** (two pieces of tracks seen as opposite sign)

What if we request 1 particle US and 1 particle DS: $z_{e+} \times z_{e-} < 0$

$P(A)$: probability that a pair with angular opening A has two particles going in opposite sides of the CDCH



Simulated vs reconstructed opening angle from MC requesting $z_{e+} \times z_{e-} < 0$

Estimated advantages:

- we lose mostly low opening angles
- we get rid of fake pairs
- we lose only ~20% of signal (X17 expected **opening angle $\sim 140^\circ$**)


Still work in progress

Increasing CDCH multiplicity


IPC MC → [2-16] MeV energy range

Generated momentum of reconstructed particles


Trigger (1 SPX and 10&10 DCH)



Trigger (1 SPX and 50&50 DCH)



CDCH multiplicity to reconstruct X17 pairs




We're looking for X17 signal as symmetric energy particles (8-10 MeV)

Requesting 50&50:

- Gets rid of high energy particles (for which pairs cannot be reconstructed)
- Little loss on 8-10 MeV particles
- Hard to reconstruct pairs with low CDCH multiplicity anyway

Angular opening: data vs MC


LiPON data - 2.1M events - 02/22 - All statistics from 22 hours of non-ZS LiPON data




Assuming same angular opening distributions for IPC18 and IPC15

→ Data consistent with a mix of EPC and IPC (~55%/45%)

For LiPON data, trigger was **10 CDCH hits over 40 mV** (US and DS) and 1 SPX hit over 70mV
→ with zero-suppression all **waveforms below 40 mV** were suppressed



→ Most hits at 20mV (likely to be largely noise)




→ Hits on a fitted track
→ Good hits also below 40 mV

Likely to explain a **factor 20 less events** reconstructed in ZS data compared to non ZS-data

→ **Gammas only generated isotropically and uniformly [1,19MeV]**


#events triggered as a function of the generated gamma momentum

TC only (1+ hit)




→ **1.2% of all 18 MeV gammas lead to trigger**

CDCH only (10&10)



→ **1.5% of all 18 MeV gammas lead to trigger**

Combined



→ **0.35% of all 18 MeV gammas lead to trigger**

Gamma rate

- **12 kHz** (on full solid angle)
- **180 Hz in CDCH single**
- **140 Hz in TC single**
- **42 Hz in trigger**

RAW RATE

LIPON/ 2uA (#418137)


CDCH trigger: 1.6 kHz

TC single: 580 Hz

« Michel »: 66 Hz


Data

Angular opening between e+ and e- [deg]




IPC

Angular opening between e+ and e- [deg]




EPC

Angular opening between e+ and e- [deg]



LiPON data - 2.1M events - 02/22 - All statistics from 22 hours of non-ZS LiPON data



→ Data consistent with a mix of EPC and IPC (~50/50)

→ A fraction of pairs at large angles are **artifacts** from reconstruction (two pieces of tracks seen as opposite sign)

Potential solution

- Request 1 particle US and 1 particle DS
- Work still ongoing