


Measurements of Transverse Spin Dependent $\pi^+\pi^-$ Azimuthal Correlation Asymmetry and Unpolarized $\pi^+\pi^-$ Cross Section in p+p Collisions at STAR at RHIC

Bernd Surrow



(On behalf of the STAR Collaboration)



15th European Research Conference on Electromagnetic Interactions with Nucleons and Nuclei
 29 October - 04 November 2023 | Annabelle Hotel, Paphos, Cyprus

Pre-conferences: 29 - 30 October 2023
 Frontiers and Careers in Photonuclear Physics - PhD Development and talks for students

Main conference: 31 October - 04 November 2023

EINN2023

Conference Topics

- Nuclear form factors and low-energy hadron structure
- Photon structure of nucleons and nuclei
- Photon excitation physics and new physics searches
- Meson structure
- Heavy and light-meson spectroscopy
- Nuclear effects with low hadron pT

Parallel Workshops

- QCD in the 21st Century: Opportunities and Director Challenges
- QCD analysis of nuclear structure

Poster Session

We invite you to submit abstracts for talks at the conference and for the poster session. Contributions not accepted for talks will be given the option of a poster presentation.

There will be a poster prize for the three best posters, which will also be presented for a general talk at the conference.

ORGANIZERS

Conference

- Esther Paschos-Chan (Cyprus)
- Cyrill Koppel (Austria)

Workshops

- André Meyer (Germany)
- Werner Vogelsang (Germany)

Pre-conference

- Elin Sjöstrand (George Washington U.)
- Francis Handberg (Austria)

SPONSORS




DOE NP contract: DE-SC0013405

Bernd Surrow

Outline

- Theoretical Foundation
- RHIC Collider and STAR experiment
- Analysis Details - $\pi^+\pi^-$ Asymmetry
- $\pi^+\pi^-$ Asymmetry Results
- Analysis Details - $\pi^+\pi^-$ Cross-Section
- $\pi^+\pi^-$ Cross-Section Results
- Summary



Theoretical foundation

□ Probe transverse proton spin structure using high-energy polarized p+p collisions

- Important new insight into the **transverse proton spin structure** at STAR in **polarized p+p collisions at high energies** using **well established processes** both theoretically and experimentally **involving jets / hadrons**
- **Transversity-related measurements**: Important insight into transverse spin structure - **Need coupling of transversity (h_1) to chiral-odd transverse spin dependent fragmentation function (FF)**:


- **Collins TMD FFs**: Azimuthal single-spin asymmetries of charged pions in jets

$$\sum_{i,j,k} h_1^{i/p_a}(x_a) f_1^{j/p_b}(x_b) H_1^\perp{}^{h/k}(z, k_T)$$

- **Di-hadron FFs**: Azimuthal correlations of charged pion pairs

$$\sum_{i,j,k} h_1^{i/p_a}(x_a) \otimes f_1^{j/p_b}(x_b) \otimes H_1^\triangleleft{}^{h_1 h_2/k}(z, M_h)$$


- Deepen our understanding concerning universality, factorization and evolution!



FF Review: A. Metz and A. Vossen, Prog. Part. Nucl. Phys. 91 (2016) 136.

Theoretical Foundation

□ Proton spin structure



		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$		$h_1^\perp = \uparrow - \downarrow$ Boer-Mulders
	L		$g_{1L} = \rightarrow - \leftarrow$ Helicity	$h_{1L}^\perp = \rightarrow - \leftarrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$ Sivers	$g_{1T}^\perp = \uparrow - \downarrow$	$h_1 = \uparrow - \downarrow$ Transversity

○ Proton spin structure in terms of parton distribution functions (PDFs)

○ Three leading twist collinear PDFs, integrated over parton transverse momentum k_T :

□ $f_1(x) =$ Unpolarized PDF

□ $g_1(x) =$ Helicity PDF


□ $h_1^q(x) =$ Transversity PDF

□ Motivation: Measurement of **observable to constrain $h_1^q(x)$ in collinear framework** in polarized p+p collisions **employing chiral-odd di-hadron fragmentation function (DiFF)**!



Theoretical Foundation

□ Transversity

Correlation between nucleon transverse polarization and transverse polarization of quarks - no gluon transversity!



- First transversity global analysis by M. Radici and A. Bacchetta (Phys. Rev. Lett. 120, 192001 (2018))
- New global analysis by JAM global analysis (arXiv 2308.14857)!
- Important connection to Lattice QCD!



Theoretical Foundation

Observables for transversity - Theoretical formulation

Di-hadron channel: $p \uparrow + p \rightarrow h^+ h^- + X$



Asymmetry: $A_{UT}^{pp} = \frac{\mathcal{H}(M_h, P_{hT}, \eta)}{\mathcal{D}(M_h, P_{hT}, \eta)}$

$$\mathcal{H}(M_h, P_{hT}, \eta) = 2P_{hT} \sum_i \sum_{a,b,c,d} \int_{x_a^{\min}}^1 dx_a \int_{x_b^{\min}}^1 dx_b \frac{dx_b}{z} h_1^a(x_a) f_1^b(x_b) \frac{d\Delta \hat{\sigma}_{a \uparrow b \rightarrow c \uparrow d}}{d\hat{t}} H_1^{\langle a,c \rangle}(z, M_h)$$

$$h_1 \leftrightarrow f_1, H_1^{\langle a \rangle} \leftrightarrow D_1$$

Unpolarized cross-section:

$$\mathcal{D}(M_h, P_{hT}, \eta) = 2P_{hT} \sum_i \sum_{a,b,c,d} \int_{x_a^{\min}}^1 dx_a \int_{x_b^{\min}}^1 dx_b \frac{dx_b}{z} f_1^a(x_a) f_1^b(x_b) \frac{d\hat{\sigma}_{ab \rightarrow cd}}{d\hat{t}} D_1^c(z, M_h)$$



Theoretical Foundation

□ Observables for transversity - Experimental measurement

- Di-hadron azimuthal correlation asymmetry, A_{UT} , for $p \uparrow + p \rightarrow h^+h^- + X$:

$$A_{UT} = \frac{d\sigma_{UT}}{d\sigma_{UU}} = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow} \propto \frac{\sum_{i,j,k} h_1^{i/p_a}(x_a) f_1^{j/p_b}(x_b) H_1^{\leftarrow h_1 h_2 / k}(z, M_h)}{\sum_{i,j,k} f_1^{i/p_a}(x_a) f_1^{j/p_b}(x_b) D_1^{h_1 h_2 / k}(z, M_h)}$$


- Independent measurement of H_1^{\leftarrow} is required from e^+e^- experiments (e.g. BELLE!)
- $D_1^{h_1 h_2}$ is least known, specifically for gluon fragmentation (New constrain from STAR!)

- Unpolarized di-hadron cross-section, $d\sigma_{UU}$, for $p \uparrow + p \rightarrow h^+h^- + X$:

- $d\sigma_{UU}$ is crucial for $D_1^{h_1 h_2}$ providing access to quarks and

gluons

- $d\sigma_{UU}$ and A_{UT} allow model-independent extraction of transversity, $h_1^q(x)$!



Theoretical Foundation



□ First proof-of-principle measurements at 200 GeV and 510 GeV

○ STAR observed significant $\pi^+\pi^-$ correlation asymmetry, A_{UT} , using 200 GeV and 500 GeV


○ $A_{UT} \propto h_1^q(x) H_1^{\langle\pi^+\pi^-\rangle}(z, M_h^2)$

○ A_{UT} enhanced around ρ -mass region.

Radici et. al. Phys. Rev. Lett. 120 (2018), 19 192001



Significant impact on $h_1^q(x)$ from STAR data at $\sqrt{s} = 200$ GeV




STAR, Phys. Lett. B 780 (2018) 332
 STAR Phys. Rev. Lett. 115 (2015) 242501



RHIC Collider and STAR Experiment


- Polarized p+p collider facility at BNL



RHIC Collider and STAR Experiment


□ Transverse spin-polarized p+p production runs

- Di-hadron FFs: 2006 at 200GeV and 2011 at 500GeV measurements and updates presented here!
- TMD Collins FFs: 2012 / 2015 at 200GeV and 2011 at 500GeV measurements
- Large data samples in 2015 at 200GeV and 2017 / 2022 at 510GeV!



RHIC Collider and STAR Experiment

□ Overview of STAR experiment



RHIC Collider and STAR Experiment

- Polarized p+p data samples and kinematic coverage

Collision mode	proton-proton						
Polarization type	transverse						
Year	2006	2011	2012	2015	2017	2022	2024
\sqrt{s} (GeV)	200	500	200	200	510	508	200
L_{int} (pb ⁻¹)	~1.8	~25	~22	~52	~320	~400	~190
$\langle P_{\text{beam}} \rangle$ (%)	~60	~53	~57	~57	~55	~52	

Published IFF A_{UT}
 STAR, Phys. Lett.
 B 780 (2018) 332
 STAR, Phys. Rev. Lett.
 115 (2015) 242501

STAR Preliminaries
 @ $\sqrt{s} = 200$ GeV
 Unpolarized $\pi^+\pi^-$
 Cross Section
 (2012)
 IFF Asymmetry
 (2015)


STAR IFF
 Preliminary @
 $\sqrt{s} = 510$ GeV

Planned IFF and
 Cross Section
 Measurements

RHIC Collider and STAR Experiment

□ Kinematic coverage

Collision mode	proton-proton						
Polarization type	transverse						
Year	2006	2011	2012	2015	2017	2022	2024
\sqrt{s} (GeV)	200	500	200	200	510	508	200
L_{int} (pb^{-1})	~ 1.8	~ 25	~ 22	~ 52	~ 320	~ 400	~ 190
$\langle P_{\text{beam}} \rangle$ (%)	~ 60	~ 53	~ 57	~ 57	~ 55	~ 52	




STAR Kinematic Coverage:

- Covers *larger Q^2 values* compared to HERMES and COMPASS.
- *Intermediate x coverage*, probing predominantly *valence quark region*.

Analysis details - $\pi^+\pi^-$ Asymmetry

□ Kinematic variables and selection cuts

Polarized parton fragments to $\pi^+\pi^-$:



Two crucial vectors: $\vec{p}_h = \vec{p}_{h_1} + \vec{p}_{h_2}$ and $\vec{R} = \frac{1}{2}(\vec{p}_{h_1} - \vec{p}_{h_2})$


○ Access to the quark polarization via correlation: $\vec{S} \cdot \vec{R} \times \vec{p}_h$

○ Pion identification by measuring the ionization energy loss (dE/dx) with $p_T^\pi > 1.5 \text{ GeV}/c$ and $|\eta| < 1$

○ Oppositely charged pion pairs, $\pi^+\pi^-$

○ Direction of \vec{R} always points from π^- to π^+ A_{UT} gets otherwise diluted

$\pi^+\pi^-$ reaction plane



Analysis details - $\pi^+\pi^-$ Asymmetry



□ Asymmetry determination

- Cross-ratio formula: ϕ_{RS} binning in A_{UT} extraction

$$A_{UT} \sin(\phi_{RS}) = \frac{1}{P} \frac{\sqrt{N^\uparrow(\phi_{RS})N^\downarrow(\phi_{RS} + \pi)} - \sqrt{N^\downarrow(\phi_{RS})N^\uparrow(\phi_{RS} + \pi)}}{\sqrt{N^\uparrow(\phi_{RS})N^\downarrow(\phi_{RS} + \pi)} + \sqrt{N^\downarrow(\phi_{RS})N^\uparrow(\phi_{RS} + \pi)}}$$

- Free from relative luminosity terms (cancels out in symmetric detector system!)
- Two transverse polarization states: \uparrow, \downarrow
- 16 ϕ_{RS} bins of uniform widths over $[-\pi, \pi]$.
- Symmetry between $[-\pi, 0]$ and $[0, \pi]$ hemispheres.
- Count $\pi^+\pi^-$ yields in each 16 ϕ_{RS} bins for each polarization states: $N^\uparrow(\phi_{RS}), N^\downarrow(\phi_{RS})$.

$P \equiv$ Average beam polarization





ϕ_{RS} binning scheme

Analysis details - $\pi^+\pi^-$ Asymmetry



□ Systematic uncertainties

- STAR PID relies on the measured ionization energy loss (dE/dx) by the TPC at low p_T .
- Time of Flight (TOF) helps to improve the STAR PID, in conjunction with the TPC via dE/dx
- The fraction of proton, kaon, and electron (backgrounds) in the pion signal region estimates the PID systematic uncertainty



$\pi^+\pi^-$ Asymmetry Results



- Asymmetry vs. pseudo-rapidity $\eta^{\pi^+\pi^-}$ at 200GeV and 510GeV



- A_{UT} increases with η at 200GeV (Run 15) and 510GeV (Run 17) - Sizable $h_1^q(x)$ expected for $\eta > 0$, i.e., large x !
- Improved PID treatment for 510GeV (Run 17) using TPC/TOF, whereas 200GeV (Run 15) based on TPC PID only so far, TOF PID incl. for final result for 200GeV (Run 15)
- Systematic uncertainties: PID and Trigger bias

$\pi^+\pi^-$ Asymmetry Results


- Asymmetry vs. invariant mass $M_{inv}^{\pi^+\pi^-}$ integrated in p_T at 200GeV and highest p_T bin at 510GeV



- A_{UT} asymmetry is enhanced around $M_{inv}^{\pi^+\pi^-} \sim 0.8$, consistent with the previous measurement and theory prediction
- Theory calculations overshoots the new measurement beyond the ρ resonance peak
- Statistical precision is significantly improved by the new result

$\pi^+\pi^-$ Asymmetry Results



- Asymmetry vs. invariant mass $M_{\text{inv}}^{\pi^+\pi^-}$ in p_T bins at 200 GeV and 510 GeV



- $A_{\text{UT}}^{\text{sin}(\phi_{\text{RS}})}$ vs $M_{\text{inv}}^{\pi^+\pi^-}$ in different p_T and $\eta^{\pi^+\pi^-}$ bins
- Signal grows stronger at higher p_T in forward $\eta^{\pi^+\pi^-}$ region / Resonance peak around $M_{\text{inv}}^{\pi^+\pi^-} \sim 0.8 \text{ GeV}/c^2 \sim M_\rho$.
- Backward $\eta^{\pi^+\pi^-}$ signal is small, mainly from low x quarks from polarized beam

$\pi^+\pi^-$ Asymmetry Results

- Asymmetry vs. transverse momentum p_T in $M_{\text{inv}}^{\pi^+\pi^-}$ bins at 200GeV and 510GeV





- Large asymmetry signal at higher p_T in forward $\eta^{\pi^+\pi^-}$ region. Stronger signal when $\langle M_{\text{inv}} \rangle \sim M_\rho$.
- Backward $\eta^{\pi^+\pi^-}$ signal ($\eta^{\pi^+\pi^-} < 0$) is small, mainly from low x quarks from polarized beam.

Analysis details - $\pi^+\pi^-$ Cross-Section




□ Selection criteria

- Di-hadron channel, $p + p \rightarrow \pi^+\pi^- + X$:
- Inclusive $\pi^+\pi^-$ differential cross section:
 - As a function of invariant mass, $M_{inv}^{\pi^+\pi^-}$, in $|\eta| < 1$.
 - Much needed for the $D_1^{h_1h_2}$ extraction.
 - Access to $D_1^{h_1h_2/g}$.
- STAR Run 2012 dataset @ $\sqrt{s} = 200$ GeV
- Triggers: JP0, JP1, JP2
- Lower trigger threshold provides better gluon sensitivity than Run 2015.
- $\pi^+\pi^-$ construction is same as in the IFF analysis, except for the track $p_T > 0.5$ GeV/c.



Analysis details - $\pi^+\pi^-$ Cross-Section

- Cross-section determination and systematic uncertainties



- PYTHIA simulated events, reconstructed through GEANT package embedded with real collision events to effectively reconstruct STAR detector responses (Embedding)
- Unfolding accounts for the bin migration effect and backgrounds
- Unfolding is performed for each trigger, allowing independent measurement of triggered cross-section

Analysis details - $\pi^+\pi^-$ Cross-Section

□ Preliminary di-hadron cross-section result

○ Top Panel:


- First unpolarized $\pi^+\pi^-$ cross-section measurement
- Good agreement in comparison to PYTHIA simulation and JAMDiFF prediction

○ Bottom Panel:

- Systematic uncertainties (Green band!)
- Statistical uncertainties (Red band!)
- Relative difference to PYTHIA / JAMDiFF shown in black/blue

○ Access to $D_1^{h_1 h_2}$ for gluons




○ Path to model-independent extraction of $h_1(x)$



Summary and Outlook

□ Summary

- New measurements of IFF di-pion asymmetries at 200GeV (2015) and 510GeV (2017)
- First di-pion cross-section measurement at 200GeV (2012)
- Improved PID systematics (Combination of TPC+TOF) for Run 17 data at 510GeV, to be applied to 200GeV measurement
- Publication of 200GeV and 510 di-pion measurements: Input to global analysis for transversity extraction!



Summary and Outlook

□ Outlook

- Precision measurement of IFF asymmetries for pions / kaons from 2015+2024 at 200GeV and 2017+2022 at 510GeV
- Planned cross-section measurements for pions at 510GeV and Kaons at 200/510GeV

