

Measurement of Transverse Spin Dependent $\pi^+\pi^-$ Asymmetry and Unpolarized $\pi^+\pi^-$ Cross Section in polarized pp Collisions at RHIC

Wednesday, 1 November 2023 17:30 (30 minutes)

The transversity distribution function of quarks, $h_1^q(x)$, encapsulates the transverse spin structure of the proton at leading twist, where x represents the longitudinal momentum fraction carried by the quark q . Extracting $h_1^q(x)$ poses a formidable challenge due to its chiral-odd nature. Measurements of final-state hadron pairs in transversely polarized proton-proton ($p^\uparrow p$) collisions directly probe collinear quark transversity through its coupling with a chiral-odd interference fragmentation function (IFF), $H_1^{\text{spherical angle}, q}$. This coupling leads to an experimentally measurable azimuthal correlation asymmetry, A_{UT} .

To extract $h_1^q(x)$ from A_{UT} asymmetry measurements, precise knowledge of IFF and unpolarized di-hadron fragmentation functions is needed.

The former is provided from e^+e^- experiments, owing to the factorization and universality of the physics mechanism in the collinear framework.

On the other hand, the latter is largely unknown but can be extracted from unpolarized di-hadron cross-section measurements in pp collisions.

In this presentation, we will present preliminary results of A_{UT} using $p^\uparrow p$ data collected in 2015 at $\sqrt{s} = 200$ GeV and in 2017 at $\sqrt{s} = 510$ GeV, and the unpolarized cross section using pp data collected in 2012 for $\pi^+\pi^-$ pairs at $\sqrt{s} = 200$ GeV by the STAR experiment. The presentation will also discuss prospects for additional data at both $\sqrt{s} = 200$ GeV and $\sqrt{s} = 510$ GeV.

Primary author: SURROW, Bernd (Temple University)

Presenter: SURROW, Bernd (Temple University)

Session Classification: Parallel workshop 1