

Neutrinoless double-beta decay with the LEGEND Experiment

Riccardo Brugnera

Università degli Studi di Padova e INFN Padova
on behalf of the LEGEND Collaboration

Outline:

- The LEGEND Experiment: general aspects
- The first stage: LEGEND-200
- LEGEND-1000



The LEGEND Experiment: general aspects

- For the **physics motivation** and **experimental situation** in the $0\nu\beta\beta$ research field: [talk of Christoph Wiesinger](#)

Large Enriched Germanium Experiment for Neutrinoless $\beta\beta$ Decay - LEGEND

~270 members, 55 institutions, 12 countries
from GERDA and MJD experiments + other groups
Collaboration formed in October 2016



LEGEND mission:


“The collaboration aims to develop a phased Ge-76 based double-beta decay experimental program with discovery potential at a half-life significantly longer than 10^{27} years, using existing resources as appropriate to expedite physics results”



LEGEND: a staged approach

First Stage (LEGEND-200):



- upgrade of the existing infrastructure of GERDA up to 200 kg
- reduction of the BI of a factor 5 w.r.t. GERDA Phase II goal
- to reach 200 kg: 35 kg from GERDA + 30 kg from MJD. The remaining 140 kg are new



Further Stages (LEGEND-1000):

- 1000 kg (staged)
- timeline and budget: highest priority from DOE after the recent Portfolio review (July 2021)
- Background reduction of a factor 20 w.r.t. LEGEND-200
- LNGS is the preferred site, SNOLAB is the alternative

sensitivity and discovery



Plots details:

- ~69% efficiency (including: isotopic fraction, active volume fraction, analysis cuts)
- GERDA Phase II: 1.5 counts/(FWHM·ton·yr)
- LEGEND-200: 0.5 counts/(FWHM·ton·yr)
- LEGEND-1000: 0.025 counts/(FWHM·ton·yr)

N.B.: background-free^(*) condition is a prerequisite for a discovery


(*) average expected bkg events < 1.0 in the ROI for the entire exposure

The first stage: LEGEND-200


- L-200 uses the GERDA infrastructure (cryostat, clean room, water plan, ...) at LNGS
- new elements: part of the enriched Ge detectors, cables, LAr veto, FE electronics, DAQ
- **February 2020**: L-200 took over the GERDA infrastructure
- **November 2021**: start commissioning
- **March 2023**: start of the physics run with ~140 kg of enriched detectors
- **L-200 Background Index goal at $Q_{\beta\beta}$:**
 $2 \cdot 10^{-4}$ cts/(keV·kg·yr)
- **L-200 Sensitivity goal:**
 $T_{1/2} > 1.5 \cdot 10^{27}$ years (90% CL exclusion)
after **1 ton·yr** of exposure
 $m_{\beta\beta} < 27 - 64$ meV (90% CL exclusion)



LEGEND-200: the experiment




active background reduction tools




Single-site event topology (SSE)

- $2\nu\beta\beta$
- $0\nu\beta\beta$




Detector multiplicity

- scattered events



Pulse Shape Discrimination (PSD)

- scattered multi-site events (MSE)
- surface events




LAr-anti coincidence

- intrinsic backgrounds
- Ge cosmogenics

Water Cherenkov anti-coincidence


- muons

Energy Resolution and Stability







Weekly energy calibration between physics runs using ^{228}Th sources

- Excellent energy resolution @ $Q_{\beta\beta}$
- Energy scale very stable between calibrations




Pulse Shape Discrimination

- based on A/E parameter, evaluated for each event
- PSD tuned to 90% survival at ^{208}Tl DEP
 - very good rejection of MSE
 -






LAr Instrumentation



- Improved Si photo-multiplier (SiPM) readout
- Improved geometry + optically active PEN → less shadowing
- Improved wavelength-shifting (TPB) fiber coating

→ ~ 3 more light wrt. GERDA



First LEGEND-200 background data



Look at

- Background before and after LAr and PSD cuts
- Compare with GERDA

Dataset based on BEGe & ICPC detectors

- Directly comparable with GERDA
- Mono-parametric PSD (A/E)
- No blinding applied


Exposure (kg·yr)	BEGe	ICPC
10.1	2.1	8.0



10 strings – 142 kg – 101 detectors


Energy spectrum after quality cuts

- Data cleaning (DC)
- Muon veto (μ)
- Ge-detector anticoincidence (AC)




Background after quality cuts

Before analysis cuts (Golden dataset)




Before analysis cuts (Golden dataset)



- No unexpected background components
 - ²³⁸U & ²³²Th decay chains, ⁴⁰K, ⁴²K
- Improved peak to Compton ratio
 - Reduced Compton continuum
 - Higher detection efficiency due to larger detectors
- Higher rate from ²⁰⁸Tl compared to GERDA
 - Expected → more construction material
- Similar spectra

Before analysis cuts (Golden dataset)




LEGEND vs. GERDA BEGe + ICPC

Background decomposition after quality cuts


Decomposition before analysis cuts

- Well described by expected contributions with current statistics




Background after QC + LAr AC


After LAr veto cut (Golden dataset)



After LAr veto cut (Golden dataset)



After LAr veto cut (Golden dataset)





- Some gamma lines “vanishes” & Compton continuum suppressed
- LAr instrumentation
 - Improved background suppression
 - higher light yield & less shadowing
 - More self-vetoing material: fibers of the LAr veto & PEN plates

LEGEND vs. GERDA BEGe + ICPC




2νββ shape and uniformity after QC + LAr AC

- Spectral shape compatible with $2\nu\beta\beta$ after LAr instrumentation anti-coincidence (LAr AC)
- ^{40}K & ^{42}K Compton edges vanish
- Uniform rate/detector in (1000–1300) keV
 - Normalized to detector specific exposure
 - BEGe/ICPC different containment eff.
- After LAr AC: Medium energy region dominated by $2\nu\beta\beta$ events



Background after QC + LAr AC + PSD

- PSD cuts multi-site and alpha events effectively
- More powerful due to higher MSE probability in larger ICOC detectors
- PSD suppression in physics data depends on actual background composition and location




LEGEND vs. GERDA BEGe + ICPC

Background Index

- Analyzed first 10.1 kg·yr of LEGEND-200 data
- ICPC&BEGe detectors
- Events in the BI-window (1930–2190) keV after QC + LAr and PSD cuts
- BI is compatible with LEGEND-200 goal:

$2 \cdot 10^{-4}$ cts/(keV·kg·yr)

- Expect 0.4 cts
- Probability to observe #cts > 0 ~38%




	LEGEND-200 BI 68%CL(cts/keV/kg/yr)	GERDA Phase II unblinded BI 68% CL(cts/keV/kg/yr)
After LAr & PS	$4.1 [1.5-11.4] \cdot 10^{-4}$	$5.2 [3.9-6.8] \cdot 10^{-4}$


LEGEND-1000

performance parameters & timeline

$0\nu\beta\beta$ decay isotope	^{76}Ge
$Q_{\beta\beta}$	2039 keV
Total mass	1000 kg
Energy resolution at $Q_{\beta\beta}$	2.5 keV FWHM
Overall signal acceptance	0.69
Total exposure	10 t·yr
Background goal	$< 10^{-5}$ cts/(keV·kg·yr) < 0.025 cts/(FWHM·t·yr)
$T_{1/2}^{0\nu}$	$1.3 \cdot 10^{28}$ yr (90% C.L. discovery) $1.8 \cdot 10^{28}$ yr (90% C.L. sensitivity)
$m_{\beta\beta}$	9.4 – 21.4 meV (99.7% C.L. discovery) 8.5 – 19.4 meV (90% C.L. sensitivity)




General layout @ LNGS




LEGEND-1000 background projections

@ SNOLAB



Expected total spectrum from $2\nu\beta\beta$ decay and from all background components after all cuts




Projected background index after all cuts:

$< 10^{-5}$ cts/(keV · kg · yr)

LEGEND-1000 target sensitivities

- ◆ $m_{\beta\beta} = m_e / \sqrt{G g_A^4 M^2 T_{1/2}}$
- ◆ Inverted Ordering: $m_{\beta\beta} > 18.4 \pm 1.3$ meV
- ◆ the discovery sensitivity required depends on the matrix element used
- ◆ the range of values given depends on the matrix elements that has been calculated for each isotope
- ◆ LEGEND-1000 will fully test inverted order and a large part of the normal ordering

Agostini, Detwiler, Benato, Menendez, Vissani
PRC, 104 (4) L042501 (2021)



Summary



^{76}Ge (92% enr.)

- ◆ The LEGEND experiment combines the best technologies from the two Ge experiments: GERDA and MAJORANA-DEMONSTRATOR
- ◆ Key feature is the staged approach: leading results at each phase
- ◆ The first phase is LEGEND-200 at LNGS using the GERDA infrastructure: the aim is to reach the limit of 10^{27} yr in the half-life of the $0\nu\beta\beta$ decay of ^{76}Ge
- ◆ LEGEND-200 is now taking data: the analysis of the first data shows that the BI is compatible with the LEGEND-200 goal
- ◆ The ultimate phase will be LEGEND-1000 able to reach an half-life greater than 10^{28} yr covering the entire inverted ordering region
- ◆ The LEGEND-1000 approval process is already begun: DOE Portfolio review (July 2021) for the choice of the best Ton-scale experiment put highest priority on LEGEND-1000.

