



Contribution ID: 144

Type: **Invited talk**

Gas of Bethe ansatz wavepackets and an ab initio derivation of generalised hydrodynamics

Monday, 4 September 2023 09:30 (50 minutes)

The hydrodynamic approximation is an extremely powerful tool to describe the behaviour of many-body systems such as gases. At the Euler scale, the approximation is based on the idea of local entropy maximisation: locally, within fluid cells, the system relaxes to a state that takes the Gibbs form. In conventional gases, these are thermal states, which include the few conserved quantities admitted by the model. In integrable systems, these are the so-called generalised Gibbs ensembles, which include the infinite set of conserved quantities, and the corresponding hydrodynamic theory is called generalised hydrodynamics (GHD). GHD applies for instance to experimentally realized one-dimensional interacting Bose gases described by the Lieb-Liniger model, and many more one-dimensional integrable systems, such as classical soliton gases and the hard-rod model. However, the local entropy maximisation is an assumption, and in general it is hard to establish the hydrodynamic equations from first principles (from the microscopic dynamics of the model). In this talk I will explain how to construct a gas of wavepackets in the Lieb-Liniger model, based on the Bethe ansatz form of the wave function. Their effective dynamics gives rise, without the assumption of local entropy maximisation, to the GHD equations. This provides a blueprint for proving the GHD equations from the Schroedinger equations. The main idea is a map to the scattering coordinates of the wavepackets' dynamics, and is similar in spirit to the techniques used in the known rigorous proof of the hydrodynamic equations for the hard-rod gas.

Primary authors: DOYON, Benjamin (King's College London); Mr HUEBNER, Friedrich (King's College London)

Presenter: DOYON, Benjamin (King's College London)

Session Classification: Bologna Workshop CFT-IM