

# Quantum Sensors for New Particles

Yoni Kahn

University of Illinois Urbana-Champaign  
Quantum Technologies for Fundamental Physics  
Erice, Italy, 9/3/23




# What is a quantum sensor?


No universally agreed-upon definition!

An attempt at a classification (boundaries are fluid):


- Detecting a single quantum of something (classically)
- Using quantum mechanics to sense small (classical) things
- Both at once (Quantum 2.0)




light




magnetic fields



charge




impulse



single-photon counting beyond the Standard Quantum Limit

# Why should theorists care?



Broadband and Resonant Approaches to Axion Dark Matter Detection

Yonatan Kahn,<sup>1,\*</sup> Benjamin R. Safdi,<sup>2,†</sup> and Jesse Thaler<sup>2,‡</sup>

[2016]

Direct Detection of sub-GeV Dark Matter with  
Semiconductor Targets

Rouven Essig,<sup>a</sup> Marivi Fernández-Serra,<sup>b,c</sup> Jeremy Mardon,<sup>d</sup> Adrián Soto,<sup>b,c</sup> Tomer  
Volansky,<sup>e</sup> Tien-Tien Yu<sup>a</sup>

[2015]

Gravitational Direct Detection of Dark Matter

Daniel Carney,<sup>1,2,\*</sup> Sohitri Ghosh,<sup>1</sup> Gordan Krnjaic,<sup>2</sup> and Jacob M. Taylor<sup>1,†</sup>


[2019]

**New theory ideas exploiting these sensors let us  
cover 50 orders of magnitude in DM mass!**


# Single-quantum detectors

Have been around for a century (bubble chambers, LHC, ...),  
but recent advances are **eV energy thresholds** and **ultra-low dark rates**

Single-charge  
semiconductor detectors:





Skipper CCD: non-destructive  
charge measurements reduce noise



NTL effect: single charges give  
quantized phonon response

Superconducting nanowire  
photon detectors:



Dark rate of 1/day!!  
[see also A. Casey, A. Sonnenschein]





# Single-quantum detectors

Towards the future:

## Single-phonon detectors


[Hochberg et al 2015]



Transition-edge sensor:  
with low enough threshold,  
can see single optical phonons,  
 $E \sim 50$  meV

## Charge and light at sub-eV scale

[SPLENDOR collab.]




Exotic narrow-gap semiconductors  
coupled to universal charge amplifier:  
strong synergy w/condensed matter,  
materials science


# Measuring classical things quantum-ly

Two examples:

Superconductors  
for EM sensing



Standard workhorse: flux  
quantization (SQUID)




New tools: high-Q SRF cavities  
[A. Romanenko, R. Cervantes, T. Roy]

Optomechanical systems  
for force sensing


[more examples: G. Marocco]

[Monteiro et al 2020]



optically-levitated microspheres

[Matsumoto et al 2019]




mirrors as pendulums

# Quantum 2.0


To get beyond Standard Quantum Limit, need to  
measure or prepare an actual quantum state

Two examples from axion DM detection:



[ADMX 2020]


destructive (power) readout + thermal vacuum



non-destructive  
photon counting  
by coupling to  
qubit

[Dixit et al 2021]

[A. Chou, C. Braggio]




squeezed  
vacuum


[Backes et al 2020]

# Why are theorists crucial?


- Define theory targets
- Invent new uses for existing sensors
- Spur development of new sensors
- Help interpret new data (CM connections!)




[Cosmic Visions 2018]




[Dimopoulos et al 2007]




[Essig et al 2017]




[Du et al 2021]




[Blinov et al 2019]



[Carney et al 2019]





[Baym et al 2020]



[Mandava et al 2022]



# From theory to the lab



From theory paper to first data in **< 5 yrs:**  
rapidly-advancing field and much more progress remains to be made!