

Contribution ID: 61

Type: not specified

## Dark matter and axion dark radiation properties from energy cascade in dark matter flow

Monday, 3 July 2023 18:11 (3 minutes)

We present a new theory to predict dark matter (DM) particle mass, size, lifetime, and properties of possible dark radiation from DM particle decay. In self-gravitating collisionless dark matter, the existence of inverse mass and energy cascade from small to large scales facilitates the hierarchical structure formation. A scaleindependent constant rate of energy cascade  $\varepsilon_u \approx -4.6 \times 10^{-7} m^2/s^3$  can be identified. The energy cascade leads to a two-thirds law for pairwise velocity and a four-thirds law for halo density profile. Both scaling laws can be directly confirmed by N-body simulations and galaxy rotation curves. For collisionless dark matter with only gravity involved, scaling laws can be extended down to the smallest scale, where quantum effects become important. Combining  $\varepsilon_u$ , Planck constant  $\hbar$ , and gravitational constant G on that scale, we predict DM particles have a mass  $m_X = (\varepsilon_u \hbar^5 G^{-4})^{1/9} = 0.9 \times 10^{12}$  GeV, a size  $l_X = (\varepsilon_u^{-1} \hbar G)^{1/3} = 3 \times 10^{-13}$  m, and a lifetime  $\tau_X = c^2/\varepsilon_u = 10^{16}$  yrs, where c is the speed of light. The energy scale  $E_X = (\varepsilon_u^5 \hbar^7 G^{-2})^{1/9} = 10^{-9}$  eV strongly suggests a dark "radiation" field to provide a viable energy dissipation mechanism. If existing, the dark "radiation" should be produced by DM decay at early time  $t_X = (\varepsilon_u^{-5} \hbar^2 G^2)^{1/9} = 10^{-6}$ s (quark epoch) with a mass of  $10^{-9}$  eV such that axion can be a very promising candidate. If axion is the dark "radiation" responsible for the energy dissipation, it should have a mass around  $10^{-9}$  eV with a GUT scale decay constant  $10^{16}$  GeV and an effective axion-photon coupling constant  $10^{-18}$  GeV<sup>-1</sup>. The dark radiation energy density is around  $\Omega_a h^2 \approx 2.6 \times 10^{-7}$ , which is about 1 percent of the photon energy in CMB. Parameterized by the increase in the effective number of neutrino,  $\Delta N_{eff}=0.02$  can be obtained. Since the DM particle mass  $m_X$  is only weakly dependent on  $\varepsilon_u$  as  $m_X \propto \varepsilon_u^{1/9}$ , the estimation of  $m_X$  should be pretty robust for a wide range of possible values of  $\varepsilon_u$ . If gravity is the only interaction and dark matter is fully collisionless, mass of  $10^{12}$  GeV seems required to produce the given rate of energy cascade  $\varepsilon_u$ . In other words, if DM particle mass has a different value, there must be some new interactions beyond gravity. This work suggests a heavy dark matter scenario with a mass much greater than WIMPs. Potential extension to self-interacting dark matter is also presented. More details can be found at arXiv:2202.07240.

Primary author: Dr XU, Zhijie (Jay) (U.S. Pacific Northwest National Lab)
Presenter: Dr XU, Zhijie (Jay) (U.S. Pacific Northwest National Lab)
Session Classification: Poster Session