




18th Patras Workshop on Axions, WIMPS and WISPS
Rijeka, 3-7 July 2023



Axions from SNe in underground neutrino detectors

Alessandro Lella




Physics Department of «Aldo Moro» University in Bari
Istituto Nazionale di Fisica Nucleare

Based on...

- AL, P. Carenza, G. Co', G. Lucente, M. Giannotti, A. Mirizzi, T. Rauscher, “*Getting the most on Supernova axions*”, e-Print: [2306.01048](https://arxiv.org/abs/2306.01048) (2023)
- P. Carenza, G. Co', AL, G. Lucente, M. Giannotti, A. Mirizzi, T. Rauscher, “*Detectability of supernova axions in underground water Cherenkov detectors*”, e-Print: [2306.17055](https://arxiv.org/abs/2306.17055) (2023)
- AL, P. Carenza, G. Lucente, M. Giannotti, A. Mirizzi, “*Protoneutron stars as cosmic factories for massive axion-like particles*”, Phys. Rev. D 107 (2023) 10

SN explosion and neutrino emission

Core-collapse SN is the terminal phase of a massive star [$M \geq 8 M_{\odot}$]. After the gravitational collapse, a shock-wave driven explosion occurs.




- The 99% of emitted energy ($\sim 10^{53}$ erg) is released via (anti)neutrinos of all species.
- From SN 1987A neutrino burst observations:
 - Duration of the burst ~ 10 s.
 - $\langle E_{\nu} \rangle \approx 15$ MeV.
- Standard picture confirmed by SN 1987A observation.

QCD Axion production in SNe

➤ Nucleon-Nucleon bremsstrahlung

[Brinkmann & Turner, *Phys. Rev. D* 38 (1988)]

[Carena & Peccei, *Phys. Rev. D* 40 (1989)]




State-of-the-art calculation include [Carenza & al., *JCAP* 10 (2019) 10]:

- Beyond OPE corrections
- Multiple scattering effects
- Effective nucleon masses

➤ Pion Conversions

[Carenza & al., *Phys.Rev.Lett.* 126 (2021)]



Contributions from:

- Contact interaction term
[Choi & al., *JHEP* 02 (2022) 143]
- Δ -mediated diagrams
[Ho & al., *Phys. Rev. D* 107 (2023)]

ALP emission spectra

- If ALPs interact weakly with nuclear matter they can *free-stream* through the SN volume

$$\frac{d^2 N_a}{dE_a dt} = \int_0^\infty 4\pi r^2 dr \frac{d^2 n_a}{dE_a dt}$$

ALP emission spectra

- If ALPs interact weakly with nuclear matter they can *free-stream* through the SN volume


$$\frac{d^2 N_a}{d E_a dt} = \int_0^\infty 4\pi r^2 dr \frac{d^2 n_a}{d E_a dt}$$

- In case of strongly coupled ALPs, they could enter the *Trapping regime*
[Caputo & al., Phys. Rev. D 105 (2022)]


$$\frac{d^2 N_a}{d E_a dt} = \int_0^\infty 4\pi r^2 dr \left\langle e^{-\tau(E_a, r)} \right\rangle \frac{d^2 n_a}{d E_a dt}$$

$$\tau \sim \int_0^\infty dr \lambda_a^{-1} \text{ optical depth for nuclear processes}$$

ALP emission spectra




ALP emission spectra




The energy-loss argument

Emission of exotic particles could cause an excessive energy-loss from SN, affecting the neutrino burst.



[Raffelt & Seckel, Phys. Rev. Lett. 60 (1998)]




The energy-loss argument

Assuming that ALP emission did not shorten the duration of the neutrino burst more than $\sim 1/2$, we require that [Raffelt, Phys. Rept. 198 (1990)]:

$$L_a \lesssim L_\nu \approx 3 \times 10^{52} \text{ erg s}^{-1}$$

Searches for solar axions in SNO.

[Bhusal et al., Phys. Rev. Lett. 126 (2021)]



[AL & al.,
e-Print: [2306.01048](https://arxiv.org/abs/2306.01048) (2023)]


The energy-loss argument

Assuming that ALP emission did not shorten the duration of the neutrino burst more than $\sim 1/2$, we require that [Raffelt, Phys. Rept. 198 (1990)]:

$$L_a \lesssim L_\nu \approx 3 \times 10^{52} \text{ erg s}^{-1}$$

Searches for solar axions in SNO.

[Bhusal et al., Phys. Rev. Lett. 126 (2021)]




[AL & al.,
e-Print: [2306.01048](https://arxiv.org/abs/2306.01048) (2023)]

Axion signal in Kamiokande II

- In case of strong couplings the ALP flux would have produced a signal in Kamiokande II.
- Seminal idea by Engel, Seckel and Hayes: look for axion-induced excitation of oxygen nuclei [*Engel et al., Phys. Rev. Lett. 65 (1990)*].



- The computation of the event rate requires:
 - SN explosion models
 - An adequate treatment of trapping regime
 - State-of-the-art nuclear models



Axion-Oxygen cross section


Introducing $C_0 = (C_p + C_n)/2$ and $C_1 = (C_p - C_n)/2$, Axion-nucleons interactions reads

$$\mathcal{H}_{aN} = -\frac{g_{aN}}{2m_N} \partial_k a \underbrace{\bar{N} \gamma^k \gamma^5 (C_0 + C_1 \tau_3) N}_{\text{Hadronic current}}$$


By computing the transition matrix element, the total cross section is [P. Carenza, G. Co', M. Giannotti, AL, G. Luente, A. Mirizzi, T. Rauscher, to appear soon]

$$\sigma(E_a) \sim \underbrace{\frac{g_{aN}^2}{m_N^2} E_a}_{\text{Strength of nuclear interactions}} \sum_J \underbrace{\left| \langle J^\Pi | |T_J| |0^+ \rangle \right|^2}_{\text{Nuclear transition matrix element}} \delta(E_a - E_J) \rightarrow \text{Computed in RPA approach}$$

Axion-Oxygen cross section



Oxygen de-excitation



- Excited oxygen states can also decay through non radiative channels (α -particles, protons, neutrons together with secondary nuclei).
- Branching ratios computed through the *SMARAGD Hauser-Feshbach reaction code* [*T. Rauscher, computer code SMARAGD, version 0.9.3s, Vol. 103, 2015*].
- γ -emission accounts for $\sim 50\%$ of the total de-excitation processes.

Detector resolution

- Detector energy resolution spreads detected energies around true photon energies.

$$\mathcal{R}(E, \epsilon) = \sum_{\omega(\epsilon)} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(E-\omega(\epsilon))^2/2\sigma^2} BR[\omega(\epsilon)] \quad \text{where } \sigma = 0.6 \sqrt{\omega(\epsilon)/\text{MeV}}$$

Detector resolution


- Detector energy resolution spreads detected energies around true photon energies.

$$\mathcal{R}(E, \epsilon) = \sum_{\omega(\epsilon)} \frac{1}{\sqrt{2\pi}\sigma^2} e^{-(E-\omega(\epsilon))^2/2\sigma^2} BR[\omega(\epsilon)]$$

where $\sigma = 0.6 \sqrt{\omega(\epsilon)/\text{MeV}}$

- Detector efficiency can be modelled as
[Hirata et al., Phys. Rev. D 38 (1988)]


$$\mathcal{E} = \max \left[0, 0.93 - e^{-(E/9 \text{ MeV})^{2.5}} \right]$$



Events number in Kamiokande-II

$$N_{\text{ev}} = F_a \otimes \sigma \otimes \mathcal{R} \otimes \mathcal{E}$$


$M_{KII} \sim 2.4 \text{ kton}$



Events number in Kamiokande-II

$$N_{\text{ev}} = F_a \otimes \sigma \otimes \mathcal{R} \otimes \mathcal{E}$$


$M_{KII} \sim 2.4 \text{ kton}$



Axion events from SN 1987A

No excess in the background of K-II around SN 1987A event ($\bar{n}_{bkg} \simeq 0.02$ events/s)

[Kamiokande Coll., Phys. Rev. Lett. 58 (1987) 1490].




[P. Carenza, G. Co', M. Giannotti, AL,
G. Lucente, A. Mirizzi, T. Ruascher,
to appear soon]

[AL & al.,
e-Print: [2306.01048](https://arxiv.org/abs/2306.01048) (2023)]

Concluding remarks

- Hadronic axions from SN in trapping regime require an adequate treatment.
- Supernova arguments alone exclude QCD axion masses $m_a \gtrsim 10^{-2}$ eV.
- No “*hadronic axion window*” [Chang & Choi, Phys. Rev. Lett. 316 (1993)] .
- No signatures due to mass of HDM axions in future cosmological surveys.




A wide-angle photograph of a dark night sky. The Milky Way galaxy is visible in the upper center, appearing as a dense, glowing band of stars. Numerous smaller stars are scattered across the dark blue and black sky. In the lower right foreground, the silhouette of a large, leafless tree is visible against the lighter sky. The horizon shows a faint glow from distant lights or the setting sun.

Thank you for your
attention

Supernova Neutrinos


Neutronization burst

- Electron capture in the inner core




Accretion

- When shock stalls, ν powered by infalling matter



Cooling




- Cooling on ν diffusion time scale



ALPs nuclear interactions


- Axions and ALPs could interact with all the Standard model particles.
- In ChPT interaction vertexes with baryons and mesons [*Ho & al., Phys.Rev.D 107 (2023)*]

$$\mathcal{L}_{nuc} = \sum_N g_{aN} \frac{\partial^\mu a}{2m_N} \bar{N} \gamma_\mu \gamma_5 N + \frac{g_{a\pi N}}{f_\pi} \partial^\mu a (i\pi^+ \bar{p} \gamma_\mu n + h.c.) + g_{aN\Delta} \frac{\partial^\mu a}{2m_N} (\bar{p} \Delta_\mu^+ + h.c.)$$



ALP mean free path


$$\lambda_a^{-1}(E_a) = \frac{1}{2|\mathbf{p}_a|} \frac{d^2 n_a(\chi E_a)}{d\Pi_a dt}$$



Axion events from SN 1987A

$$N_{\text{ev}} \lesssim \begin{cases} 2 \sqrt{n_{\text{bkg}} \Delta t} & \text{if } m_a \lesssim 17 \text{ eV} \\ 2 \sqrt{n_{\text{bkg}} \Delta t_a} & \text{if } m_a > 17 \text{ eV} \end{cases}$$

$$\Delta t \approx 12 \text{ s}$$



$$\begin{aligned} \Delta t_a(m_a) &\approx t(E_{\min}, m_a) - t(E_{\max}, m_a) \\ &\approx 1.82 \text{ s} \left(\frac{m_a}{10 \text{ eV}} \right)^2 \end{aligned}$$

Summary plot, no pions

