

# EIC next 10+ years to start of operations

Abhay Deshpande

June 22, 2023

Discussion in the INFN General Meeting

These are not “the opinion” of the EIC Project (though it should be ;-))

These are my personal thoughts on this topic for discussions within the EICUG community based on experience over many years.

# From now until first physics collisions


1. EIC Project (1 machine + 1 detector) timeline & criticality of its success
2. Some thoughts on EIC early operations & planning (motivated by Pietro's questions and comments)

How would the EIC Operations evolve? – Historical precedent and experience


3. Is “theory” ready to meet EIC data?
4. Ultimate Success of EIC → A second detector  
*Expansion of EIC Science Scope for ePIC & science scope for Detector 2*

# Item 1: EIC project and its timeline


# EIC Accelerator Design



Center of Mass Energies:	20GeV - 140GeV
Luminosity:	$10^{33} - 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ / $10\text{-}100\text{fb}^{-1}$ / year
Highly Polarized Beams:	70%
Large Ion Species Range:	p to U
Number of Interaction Regions:	Up to 2!







**BROOKHAVEN NATIONAL LABORATORY**  
 D. Gibbs  
 Laboratory Director  
 R. Tribble Deputy Director for Science & Technology J. Anderson Deputy Director for Operations

**EIC BOARDS**  
**EIC Advisory Board**  
 S. Henderson, TJ Director, Chair  
**EIC Resource Review Board**  
 H. Gao, BNL Associate Lab Director for Nuclear & Particle Physics  
 D. Dean, TJ Deputy Director for Science  
 TBD Co-Chair, International Funding Agency


**ELECTRON-ION COLLIDER PROJECT**  
 J. Yeck (BNL), Project Director  
 F. Willeke (BNL), Deputy Project Director and Technical Director  
 K. Smith (BNL), Deputy Technical Director  
 R. Ent (TJ), Co-Associate Director for the Experimental Program A. Lung (TJ), Deputy Project Director for TJNAF Partnership  
 E. Aschenauer (BNL), Co Associate A. Seryi (TJ), Associate Director for Accelerator Systems & Technical Integration International Partnership  
 L. Lari (BNL), Project Manager


**EIC COMMITTEES**  
**Project Advisory Committee**  
 T. Glasmacher, Chair  
**Machine Advisory Committee**  
 T. Raubenheimer, Chair  
**Detector Advisory Committee**  
 E. Kinney, Chair  
**Infrastructure Construction Advisory Committee**  
 M. Fallier, Chair

**EIC USERS**  
**EIC User Group Steering Committee**  
 R. Fatemi, Chair  
 M. Radici, Co-Chair  
 ePIC Collaboration  
 TBD – Spokesperson


A. Deshpande (BNL)  
 EIC Science Director

M. Chamizo Llatas (BNL)  
 EIC In-Kind Manager  
 K. Amm (BNL)  
 EIC SC Magnet Production Manager





# ePIC Detector Design



## Tracking:

- New 1.7T solenoid
- Si MAPS Tracker
- MPGDs ( $\mu$ RWELL/ $\mu$ Megas)

## PID:


- hpDIRC
- mRICH/pfRICH
- dRICH
- AC-LGAD ( $\sim 30$ ps TOF)

## Calorimetry:


- SciGlass/Imaging Barrel EMCal
- PbWO<sub>4</sub> EMCal in backward direction
- Finely segmented EMCal +HCal in forward direction
- Outer HCal (sPHENIX re-use)
- Backwards HCal (tail-catcher)



# Project Schedule



# High Level Installation Schedule



## Tight Timeline:

- Solenoid and Barrel HCal need to be ready by Jan 2029
- all other subdetectors need to be ready between 06/29 to 06/30 depending on their location in the detector

EIC From now to the start of Operations: Discussion at the INFN

General Annual Meeting

Aschenauer & Ent





# Item 2:

## Transition to operation:

### Operations planning

Start with lowest possible luminosity to build confidence in the machine in order to avoid accidental damage to the detector components and the machine itself. As we go forward and we get some explicit guidance on luminosity rise from EIC/CAD – we should be able to make more concrete proposals for early running – energy, species and physics goals/outcomes.

# @RHIC Polarized protons harder than nuclei




# Historical Precedence:

## HERA Experience:

- HERA highest luminosity  $\sim 5 \times 10^{31} \text{ cm}^{-2}\text{sec}^{-1}$

## HERA & RHIC lessons:

- Start of the machine slow and deliberate
- Development of polarization (both beams along with luminosity) will take time but early investments pay-off



# Preliminary thoughts & estimates

Time on this slide starts at CD4

We should plan to see about  $1 \times 10^{32} \text{cm}^{-2} \text{sec}^{-1}$  in the 1<sup>st</sup> year (already twice the maximum reached at HERA), and hope on reaching  $10^{33}$  in ~2-3 years, and  $10^{34}$  in ~5+ years at 10 x 250 GeV polarized proton operation.

18 GeV e beam will be experimented with before CD4 (after CD4A) if RF is ready but not “operated for physics” until about 3-5 years into the program.

I expect polarization of electron and proton would be harder to achieve compared to e-A luminosity.

These coupled with the fact that we would like fast physics results: I think we will run e-A physics 10 x 100 GeV beams at the beginning, with highest luminosity possible and run at least two nuclei in the first two years. In those years, ample time would be given to develop polarization and luminosity for e-p.

# Start EIC physics program hence be:

**Begin with** electron-Nucleus/Ion Physics: Intermediate energy and high energy operation with different nuclei

- Search for saturation from inclusive to sem-inclusive  $\rightarrow F_2^A$ , disappearance of jet
- Interactions in color with nuclear matter: with multiple nuclear sized targets; study of jet production and its interaction with nuclear matter, jet internal structure, hadronization
- With luminosity increase: exclusive diffraction in e-A to establish saturation
- Comparison runs of e-p should be expected at moderate luminosity (not for polarization) but we should be ready to utilize what we get **for inclusive & start semi-inclusive spin physics**

Allow ample **time for luminosity and polarization development. In RHIC era, significant time of p-p was given to R&D.** This philosophy gave high returns in the end.



# Operational thoughts further....

$10^{33}$  luminosity  $\rightarrow$  20 fb<sup>-1</sup>/year including 70% accelerator & detector efficiency

White paper and Yellow Report hence assumed those luminosities and hardly any measurement (other than a few in TMDs and GPDs) showed more required luminosity in e-p.

- It is hence my guess that everyone's favorite measurement will be start happening after the 2<sup>nd</sup> year with polarized beams and unpolarized nuclear beams.
- Transverse spin measurements needing only proton beam polarization should be easier to achieve than double longitudinal spin measurements with high e-polarization

# Item 3:

## Is theory ready for EIC to start?

Greatly benefitted from [Daniel deFlorian's](#) recent talk at [DIS2023](#) and also input from [Werner Vogelsang](#) and a recent White Paper submitted to the NSAC Long Range Planning process:


“A Case for an EIC Theory Alliance: Theoretical Challenges of the EIC”, R. Abir et al. arXiv: 2305.14572v1

Based on: CFNS/Stony Brook & CFNS/MIT Workshops in 2022:

- 1) [Precision QCD for e-p at the EIC](#) and 2) [Theory for EIC in the Next Decade](#)


# Anticipated precision at the LHC drove higher precision calculations in theory

- ▶ Still plenty of room for new discoveries : two main scenarios



- ▶ Search for (and find) new states
- ▶ Resonance needs “descriptive” TH

- ▶ Most likely look for “new interactions”
- ▶ Small deviations from SM : PRECISION
- ▶ EFT description / BSM model




Theory  
uncertainties HAD  
to be reduced


# Why higher order corrections?

- ▶ Large Corrections  $\alpha_s \sim 0.1$

- ▶ Better TH/EXP matching



- ▶ Accurate Theoretical Predictions **Scale dependence reduced**



- DIS Single jet production at N3LO

Currie, Gehrmann, Glover, Huss, Niehues, Vogt (2018)


- ▶ uncertainties still larger than experimental errors (HERA)



Recall, EIC luminosity will be at least 100 (if not 1000) times larger than HERA

Clearly theory precisions needs to be visited immediately.

# The NLO revolution standard automation



- Final goal: Really automatic NLO calculations **zero cost for humans**
- Apart from very large multiplicities and loop induced processes automatic NLO “conceptually” solved

- in a few years a number of codes

HELAC-NLO, Rocket, BlackHat+SHERPA, GoSam+SHERPA/MADGRAPH, NJet+SHERPA, Madgraph5-aMC@NLO, RECOLA, OpenLoops+SHERPA..

- Uncertainties, EW corrections, masses, matching to PS, etc



pQCD @ HEC

Daniel de Florian

Automation at NLO already used and demonstrated.

Modern version: with ChatGPT LO tested(?) (Joke!)

## LO can (almost) be done by chatGPT



Can you write a Fortran 77 code that computes the DIS cross section at LO using LHAPDFs? Please work in a 5-flavor scheme and using the latest set of CTEQ to make Joey Huston happy



```
sql
Copy code

Sample Fortran 77 code to compute the DIS cross section at LO using LHAPDF
in a 5-flavor scheme and neglecting the contribution from the photon PDF

implicit none

Declare the input variables
real*8 x, q2, y, s, tau, pdf, sigma
integer flavor, nset, ievtnum
character*255 pdffile

Declare the LHAPDF functions
external initpdfset, evolvepdf, alphapdf, endpdfset

Set the input kinematics
x = 0.1d0
q2 = 10.d0**6
y = 0.5d0
s = q2 / x / y
tau = q2 / s


Set the LHAPDF set and initialize it
pdffile = 'cteq66'
call initpdfset(nset,pdf,0)

Compute the LO partonic cross section for each flavor
sigma = 0.d0
do flavor = 1, 5
  pdf = evolvepdf(x, q2, flavor) / x ! divide by x to get the PDF for t
  sigma = sigma + alpha_em**2 / (2.d0 * s * q2) * pdf * &
    ((4.d0 / 9.d0 + 16.d0 / 27.d0 * delta(flavor,2)) * (1.d0 + (1.
    (4.d0 / 27.d0 * delta(flavor,2)) * (1.d0 + y**2))
end do

Compute the DIS cross section by convolving with the PDF of the proton
sigma = sigma * 3.d0 / 2.d0 / x / q2 / (1.d0 - y)
```



# The NNLO revolution standard



New Standard at LHC is NNLO

But it took 20 years from 1990's to 2020's for all processes to be calculate & tested

- ▶ Inclusive DIS known at N3LO Vermaseren, Vogt, Moch (2005)
- ▶ Quark kernels almost known at N3LO Falcioni, Herzog, Moch
- ▶ Steps toward N4LO (first moments): N4LO DIS+evolutic  
Moch, Ruijl, Ueda, Vermaseren, Vogt (2022) A. Pelloni DIS23
- DIS Single jet production at N3LO Currie, Gehrmann, Glover, H
- ▶ What about other less inclusive observables?
- ▶ In many cases legacy calculations from HERA (missing pol
- ▶ Some hard to find (needed for validation), some wrong (I
- ▶ Need to reach level of LHC calculations and account for
- ▶ Set **new standards** for EIC

## Need to gather TH community for precision at EIC

### EIC Wishlist

Precision QCD predictions for  $ep$  physics at the EIC  
CFNS August 2022

#### Regarding DATA

- Measure cross-sections instead of ratios for a more dedicate analysi
- Release both QED corrected and uncorrected data
- Develop method for unbinned cross-sections

#### Regarding PDFs, FFs and more distributions

- Replication of PDF4LHC and HERA efforts for EIC : PDF4EIC
- Perform Global NNLO analysis of polarized PDFs
- Impact of QED corrections to polarized PDFs
- Perform Global analysis of DVCS
- Generate threshold resummed PDFs and FFs
- New set of photon PDFs (existing are outdated)

#### Regarding Perturbative corrections (QCD/QED)

- Jets in DIS: matched NNLO +  $q_T$  resummation
- DIS with QED/EW corrections
- Calculations for dihadron production

#### Regarding Theoretical Issues

- Discuss (non)Universality of TMDs
- Search for ideal observables to measure Wigner distribution
- Role of lattice in PDFs (in two slides!)
- Studies for Lambda polarization at EIC
- $N^*$ ,  $\Delta$  electro-couplings at  $Q^2 > \text{GeV}^2$
- Proton structure functions in transition regime  $\rightarrow$  DIS
- Dipole and quadrupole amplitudes

Precision QCD predictions for  $ep$  physics at the EIC (II)

<https://indico.bnl.gov/e/qcd4eic>

CFNS (StonyBrook)  
18-22 September 2023


# Conclusions

- ▶ Amazing progress in fixed order calculations during the last two decades
  - Automation of NLO **Driven by LHC**
  - Several NNLO processes  $2 \rightarrow 2$  and already a few  $2 \rightarrow 3$  ✓
  - Even N<sup>3</sup>LO for simpler kinematics and first set of splitting functions
  - Account for QED/EW effects
- ▶ But... **Reaching new bottlenecks**
  - ▶  $2 \rightarrow 3$  (Massive) 2-loop amplitudes complicated beyond leading color
  - ▶ Real radiation far from trivial (**numerical infrared treatment**)
  - ▶ N<sup>3</sup>LO beyond Drell-Yan like processes will require significant developments
  - ▶ Need a more rigorous treatment of TH uncertainties
- ▶ **From LHC to EIC**
  - ▶ By the time EIC starts taking data ~N<sup>3</sup>LO might be the new standard!


An enormous thrust in precision calculations in Theory including already emerging precision in Lattice QCD is needed in the next ~10 years before or around the same time the EIC data will become available

Item 4:  
EIC would be incomplete without  
the 2<sup>nd</sup> IR and a detector

# EIC Accelerator Design



Center of Mass Energies:	20GeV - 140GeV
Luminosity:	$10^{33} - 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ / $10\text{-}100\text{fb}^{-1} / \text{year}$
Highly Polarized Beams:	70%
Large Ion Species Range:	p to U
Number of Interaction Regions:	Up to 2!





# Two documents: with overlapping arguments



Ent and Milner et al for the EICUG SC

JLAB-PHY-23-3761

## Motivation for Two Detectors at a Particle Physics Collider

Paul D. Grannis<sup>\*</sup> and Hugh E. Montgomery<sup>†</sup>  
(Dated: March 27, 2023)

It is generally accepted that it is preferable to build two general purpose detectors at any given collider facility. We reinforce this point by discussing a number of aspects and particular instances in which this has been important. The examples are taken mainly, but not exclusively, from experience at the Tevatron collider.

arXiv: 2303.08228v2 March 24, 2023<sup>4</sup>

Case for two detectors being made from **Nuclear** and Particle Physics

While EIC project (machine and 1<sup>st</sup> detector)  
have to succeed....

I think we have everything we need to sow  
the seeds for a 2<sup>nd</sup> detector

## Opportunity for complementary detector designs for different IRs exists!

# Complementarity for 1<sup>st</sup>-IR & 2<sup>nd</sup>-IR

	1 <sup>st</sup> IR (IP-6) ePIC	2 <sup>nd</sup> IR (IP-8)
Geometry:	<p>ring <b>inside to outside</b></p> <p>tunnel and assembly hall are larger</p> <p><b>Tunnel:</b> <math>\varnothing</math> 7m +/- 140m</p>	<p>ring <b>outside to inside</b></p> <p>tunnel and assembly hall are smaller</p> <p><b>Tunnel:</b> <math>\varnothing</math> 6.3m to 60m then 5.3m</p>
Crossing Angle:	<p><b>25 mrad</b></p>	<p><b>35 mrad</b></p> <p>secondary focus</p>
Luminosity:	<p>different blind spots</p> <p>different forward detectors and acceptances</p> <p>different acceptance of central detector</p> <p>More luminosity at lower <math>E_{CM}</math> ?</p> <p>Optimize Doublet focusing FDD vs. FDF</p> <p>→ impact of far forward <math>p_T</math> acceptance</p>	
Experiment:	<p><b>1.7 Tesla</b> or <b>3 (?) Tesla</b></p> <p>different subdetector technologies</p>	

## Potential Physics topics beyond Core EPIC detector's mandate exist

### Focus first on Physics beyond the EIC's core (CD0) science

(there will be others: some overlapping, some exclusive due to different IR design)

#### Physics with nucleons and nuclear targets:

- Quark **Exotica**: 4,5,6 quark systems...? Much interest after recent **LHCb** led results.
- **Nuclear Fragments** from light and heavy nuclei : e-A – Connecting to low energy nuclear physics (exotic nuclei), studying the shapes of nuclei and their internal substructure; entanglement, entropy, fragmentation, hadronization and such phenomena

#### New Studies with proton or neutron target: (mostly overlapping?)

- Impact of precision measurements of unpolarized PDFs at high  $x/Q^2$ , on LHC-Upgrade results(?)
- Precision calculation of  $\alpha_S$  : higher order pQCD calculations, twist 3
- Heavy quark and quarkonia (c, b quarks) studies with 1000 times lumi of HERA (and polarization)

#### Precision electroweak and BSM physics:

- Electroweak physics & searches beyond the SM: Parity, charge symmetry, lepton flavor violation
- LHC-EIC Synergies & complementarity: (**muon detectors were of particular interest**)

# EIC Science from the perspective of High Energy Physicists

arXiv:2203.13199v1 [hep-ph] 24 March 2022

## Snowmass 2021 White Paper: Electron Ion Collider for High Energy Physics


R. Abdul Khalek,<sup>1</sup> U. D'Alesio,<sup>2,3</sup> Miguel Arratia,<sup>4,5,\*</sup> A. Bacchetta,<sup>6</sup> M. Battaglieri,<sup>7,1</sup> M. Begel,<sup>8</sup> M. Boglione,<sup>9</sup>  
R. Boughezal,<sup>10</sup> Renaud Boussarie,<sup>11,\*</sup> G. Bozzi,<sup>12,3</sup> S. V. Chekanov,<sup>10</sup> F. G. Celiberto,<sup>13,14,15</sup> G. Chirilli,<sup>16</sup>  
T. Cridge,<sup>17</sup> R. Cruz-Torres,<sup>18</sup> R. Corliss,<sup>19,20</sup> C. Cotton,<sup>21</sup> H. Davoudiasl,<sup>8</sup> A. Deshpande,<sup>8,19</sup> Xin  
Dong,<sup>18,\*</sup> A. Emmert,<sup>21</sup> S. Fazio,<sup>8</sup> S. Forte,<sup>22</sup> Yulia Furletova,<sup>1,\*</sup> Ciprian Gal,<sup>23,20,\*</sup> Claire Gwenlan,<sup>24,\*</sup>  
V. Guzey,<sup>25</sup> L. A. Harland-Lang,<sup>26</sup> I. Helenius,<sup>27,28</sup> M. Hentschinski,<sup>29</sup> Timothy J. Hobbs,<sup>30,31,\*</sup> S. Höche,<sup>32</sup>  
T.-J. Hou,<sup>33</sup> Y. Ji,<sup>18</sup> X. Jing,<sup>34</sup> M. Kelsey,<sup>35,18</sup> M. Klasen,<sup>36</sup> Zhong-Bo Kang,<sup>37,38,20,\*</sup> Y. V. Kovchegov,<sup>39</sup>  
K.S. Kumar,<sup>40</sup> Tuomas Lappi,<sup>27,28,\*</sup> K. Lee,<sup>41,42</sup> Yen-Jie Lee,<sup>43,44,\*</sup> H.-T. Li,<sup>45,46,47</sup> X. Li,<sup>48</sup> H.-W. Lin,<sup>49</sup>  
H. Liu,<sup>40</sup> Z. L. Liu,<sup>50</sup> S. Liuti,<sup>21</sup> C. Lorcé,<sup>51</sup> E. Lunghi,<sup>52</sup> R. Marcarelli,<sup>53</sup> S. Magill,<sup>54</sup> Y. Makris,<sup>55</sup>  
S. Mantry,<sup>56</sup> W. Melnitchouk,<sup>1</sup> C. Mezrag,<sup>57</sup> S. Moch,<sup>58</sup> H. Moutarde,<sup>57</sup> Swagato Mukherjee,<sup>8,†</sup> F. Murgia,<sup>3</sup>  
B. Nachman,<sup>59,60</sup> P. M. Nadolsky,<sup>61</sup> J.D. Nam,<sup>62</sup> D. Neill,<sup>63</sup> E.T. Neill,<sup>53</sup> E. Nocera,<sup>64</sup> M. Nycz,<sup>21</sup>  
F. Olness,<sup>61</sup> F. Petriello,<sup>46,47</sup> D. Pitonyak,<sup>65</sup> S. Plätzer,<sup>66</sup> Stefan Prestel,<sup>67,\*</sup> Alexei Prokudin,<sup>68,1,\*</sup> J. Qiu,<sup>1</sup>  
M. Radici,<sup>6</sup> S. Radhakrishnan,<sup>69,18</sup> A. Sadofyev,<sup>70</sup> J. Rojo,<sup>71,72</sup> F. Ringer,<sup>73,19</sup> Farid Salazar,<sup>37,38,74,75,\*</sup>  
N. Sato,<sup>1</sup> Björn Schenke,<sup>8,\*</sup> Sören Schlichting,<sup>76,\*</sup> P. Schweitzer,<sup>77</sup> S. J. Sekula,<sup>78,\*</sup> D. Y. Shao,<sup>79</sup>  
N. Sherrill,<sup>80</sup> E. Sichtermann,<sup>18</sup> A. Signori,<sup>6</sup> K. Şimşek,<sup>81</sup> A. Simonelli,<sup>9</sup> P. Sznajder,<sup>82</sup> K. Tezgin,<sup>83</sup>  
R. S. Thorne,<sup>17</sup> A. Tricoli,<sup>8</sup> R. Venugopalan,<sup>8</sup> A. Vladimirov,<sup>84</sup> Alessandro Vicini,<sup>22,\*</sup> Ivan Vitev,<sup>85,\*</sup>  
D. Wiegand,<sup>86</sup> C.-P. Wong,<sup>48</sup> K. Xie,<sup>87</sup> M. Zaccheddu,<sup>2,3</sup> Y. Zhao,<sup>88</sup> J. Zhang,<sup>89</sup> X. Zheng,<sup>21</sup> and P. Zurita<sup>84</sup>



EIC's versatility, resolving power and intensity (luminosity) open new windows of opportunity to address some of the crucial and fundamental scientific questions in particle physics. The paper summarizes the EIC physics from the perspective of the HEP community participating in Snowmass 2021

- Beyond the Standard Model Physics at the EIC
- Tomography (1,3,5 d PDFs) of Hadrons and Nuclei at the EIC
- Jets at EIC
- Heavy Flavors at EIC
- Small-x Physics at the EIC

- High luminosity wide CM range
- Polarized e, p, and ion beams
- All nuclei



# Detector technologies EIC & LHC:

*Potential for overlap and collaboration: Many EIC collaborators already part of RD51 (and family) at CERN & vice-versa.*

- MAPS  $\mu$ Vertex for primary/secondary vtx: barrel & end-caps (ALICE ITS3)
- Micro Pattern Gas Detectors: large rapidity, spatial resolution  $\sim 100 \mu\text{m}$
- Electromagnetic Calorimetry for kinematic reconstruction, precise energy measurements  $e, \gamma$ ;  $e/\pi$  &  $\pi^0/\gamma$  separation. Various technologies at various locations:
  - W/SciFi w/o PMT, PbWO<sub>4</sub>, SiGlass; AstroPix & Pb/SciFi
  - High resolution Crystal Cal for e-endcap
  - Barrel EMCal 6 layers AstroPix and Pb/SciFi
- Particle Identification – extremely important for most EIC physics
  - K/pi separation over a wide range 1-20 GeV/c
  - Hadron ID: hpDIRC in Barrel, forward EndCap: dual RICH, backward Endcap: modular RICH or pF RICH, also TOF for short lever arm : LGAD, LAPPD
- Streaming Readout

# Vision for the 2<sup>nd</sup> detector: C<sup>3</sup>

- **Complementary** (IR, detector technologies & design)
  - Continue to explore complementary ready and not-yet-ready technologies
  - Generic detector R&D program – Run through JLab
- **Complementary** (physics)
  - A significant list of physics topics (some-exclusive to 2<sup>nd</sup> IR, some-overlapping) exists: drill down and see which of those can *develop into strong pillars of science for the 2<sup>nd</sup> detector*.
  - New physics developing around the world : we need to monitor constantly
- **Complementary** (people)
  - New non-US/outside groups who may bring new interests & funding in future
  - New US groups – other than those with significant responsibilities in ePIC

## Path forward 2<sup>nd</sup> Detector: focused workshops and detector studies on new physics topics:

- ✓ Look at **complementary detector technologies** (to ePIC) and attract groups that are experts in them to the EICUG
- ✓ Focused discussions on **new physics topics** (not just listed in this talk but also beyond) to try to make a unique case complementary to ePIC/EIC White Paper
- ✓ **Build community** – new groups/faces/resources needed to contribute and become part of new detector effort

### Resources:

Generic detector R&D – supported by DOE administered from JLab

Center for Frontiers in Nuclear Science @ Stony Brook (& EIC – Theory Institute at BNL) and the EIC<sup>2</sup> at JLab

## Remarks:

- EIC project's path is well understood. **Its success is paramount.**
- 2<sup>nd</sup> detector is essential for completing the Vision of EIC
  - **C<sup>3</sup>** : **Complementary physics, technology and people**
- It is time to march forward developing a design and case for the 2<sup>nd</sup> detector: Detailed studies through **series of workshops, outreach and critical evaluation** for each developing argument → **Plan an INT-Program in ~2025** like we had in 2010

I look forward to discussions on any and all fronts for the 2<sup>nd</sup> detector development.

# Summary

1. EIC Project (1 machine + 1 detector) timeline & criticality of its success  
Consolidate the collaboration, science and detector : Its success is paramount
2. Some thoughts on EIC early operations & planning
  - Historical precedent and early operational experience: inclusive – to – exclusive, e-A to polarized e-p/A
3. Is “theory” ready to meet EIC data? NO
4. Ultimate Success of EIC → Physics with a second detector :  
(Complementarity)<sup>3</sup> – physics, detector and people.



# Thank you

# EIC Design Parameters

**Table 3.3:** EIC beam parameters for different center-of-mass energies  $\sqrt{s}$ , with strong hadron cooling. High divergence configuration.

Species	proton	electron	proton	electron	proton	electron	proton	electron	proton	electron
Energy [GeV]	275	18	275	10	100	10	100	5	41	5
CM energy [GeV]	140.7		104.9		63.2		44.7		28.6	
Bunch intensity [ $10^{10}$ ]	19.1	6.2	6.9	17.2	6.9	17.2	4.8	17.2	2.6	13.3
No. of bunches	290		1160		1160		1160		1160	
Beam current [A]	0.69	0.227	1	2.5	1	2.5	0.69	2.5	0.38	1.93
RMS norm. emit, h/v [ $\mu\text{m}$ ]	5.2/0.47	845/71	3.3/0.3	391/26	3.2/0.29	391/26	2.7/0.25	196/18	1.9/0.45	196/34
RMS emittance, h/v [nm]	18/1.6	24/2.0	11.3/1.0	20/1.3	30/2.7	20/1.3	26/2.3	20/1.8	44/10	20/3.5
$\beta^*$ , h/v [cm]]	80/7.1	59/5.7	80/7.2	45/5.6	63/5.7	96/12	61/5.5	78/7.1	90/7.1	196/21.0
IP RMS beam size, h/v [ $\mu\text{m}$ ]	119/11		95/8.5		138/12		125/11		198/27	
$K_x$	11.1		11.1		11.1		11.1		7.3	
RMS $\Delta\theta$ , h/v [ $\mu\text{rad}$ ]	150/150	202/187	119/119	211/152	220/220	145/105	206/206	160/160	220/380	101/129
BB parameter, h/v [ $10^{-3}$ ]	3/3	93/100	12/12	72/100	12/12	72/100	14/14	100/100	15/9	53/42
RMS long. emittance [ $10^{-3}$ , eV·s]	36		36		21		21		11	
RMS bunch length [cm]	6	0.9	6	0.7	7	0.7	7	0.7	7.5	0.7
RMS $\Delta p/p$ [ $10^{-4}$ ]	6.8	10.9	6.8	5.8	9.7	5.8	9.7	6.8	10.3	6.8
Max. space charge	0.007	neglig.	0.004	neglig.	0.026	neglig.	0.021	neglig.	0.05	neglig.
Piwinski angle [rad]	6.3	2.1	7.9	2.4	6.3	1.8	7.0	2.0	4.2	1.1
Long. IBS time [h]	2.0		2.9		2.5		3.1		3.8	
Transv. IBS time [h]	2.0		2		2.0/4.0		2.0/4.0		3.4/2.1	
Hourglass factor $H$	0.91		0.94		0.90		0.88		0.93	
Luminosity [ $10^{33}\text{cm}^{-2}\text{s}^{-1}$ ]	1.54		10.00		4.48		3.68		0.44	

# eRHIC Hadron Polarization

## Measured RHIC Results:

- Proton Source Polarization 83 %
- Polarization at extraction from AGS 70%
- Polarization at RHIC collision energy 60%

## Planned near term improvements:

**AGS:** Stronger snake, skew quadrupoles, increased injection energy

→ expect 80% at extraction of AGS

**RHIC:** Add 2 snakes to 4 existing no/reduce polarization loss

→ expect 80% in Polarization in RHIC and EIC

Expected simulations results benchmarked against RHIC operations

## <sup>3</sup>He in eRHIC with six snakes

Achieved 85% polarization in <sup>3</sup>He ion source



Polarization preserved with 6 snakes for up to twice the design emittance

## Deuterons in eRHIC:

Requires tune jumps in the AGS, then

benchmarked simulation show 100% Spin transparency

**No polarization loss** expected in the EIC hadron ring



# EIC achieves high luminosity $L = 10^{34} \text{ cm}^{-2}\text{s}^{-1}$

- **Large bunch charges**  $N_e \leq 1.7 \cdot 10^{11}$ ,  $N_p \leq 0.69 \cdot 10^{11}$
- **Many bunches**,  $n_b=1160$ 
  - crossing angle collision geometry
  - large total beam currents
  - limited by installed RF power of 10 MW
- **Small beam size** at collision point achieved by
  - small emittance, requiring either:
    - strong hadron cooling to prevent emittance growth
    - or frequent hadron injection
  - and strong focusing at interaction point (small  $\beta_y$ )
  - flat beams  $\sigma_x/\sigma_y \approx 10$
- **Strong, but previously demonstrated beam-beam interactions**  
 $\Delta v_p = 0.01$  demonstrated in RHIC  
 $\Delta v_e = 0.1$  demonstrated in HERA, B-factories

