

B Physics Observables for New Physics Discoveries

Update from Valencia

Tim Gershon
University of Warwick


SuperB meeting in Elba,
31st May 2008

Outline


- B physics measurements
 - Precise CKM determination: V_{ub}
 - Theoretically clean rare decays (inclusive $b \rightarrow s\gamma$, $b \rightarrow sll$, exclusive $B \rightarrow K(^*)\nu\nu$, $B \rightarrow l\nu$)
- “Golden modes”
- High energy interplay and Snowmass points
- New physics models

SuperB UT fit scenarios

“the nightmare”



“the dream”



- Possible NP discovery from precise CKM metrology
- Precise knowledge of SM parameters essential in any scenario

How to make the dream reality?

- Need very precise measurements
- Is 2% error on V_{ub} feasible?
 - Experimentally: yes (B.Viaud)
 - Theoretically: maybe (P.Gambino)
 - Note
 - “maybe” is a very positive answer at this stage
 - need to discuss with other experts
 - (endorsed already by I.Bigi)

Towards precise V_{ub}

- Control weak annihilation and shape function
 - Possible from data
- Control error from m_b
 - Precise measurements of radiative decays
 - Improvements in lattice and other theory
 - Inclusive m_X analysis \Rightarrow reduced sensitivity
- Control perturbative corrections
 - Homework for theorists


A new theoretical analysis

- **kinetic scheme**. Wilsonian infrared cutoff $\mu \sim 1$ GeV: contribution of soft gluons absorbed into definition of OPE parameters AND distribution function(s)
- **Fermi motion**: finite m_b SF, includes all available subleading corrections
- local OPE breaks down at **high q^2** : need to model the tail, consistent with positivity, **WA** naturally emerge.
- Triple differential distribution including all known pert and nonpert effects, c++ code

published in PG, P.Giordano, G.Ossola, N.Uraltsev, JHEP10(2007)058

Theoretical errors

- Parametric errors generally dominant, in particular m_b , 3-4%
- Perturbative corrections 2-3%
- Functional form 1-2%
- Modelling of the q^2 tail and WA depending on cut from 0 to 7%. WA tends to decrease V_{ub}



cuts	$ V_{ub} \times 10^3$	f	exp	par	pert	tail model	q_*^2	X	ff	tot th
A [28]	3.87	0.71	6.7	3.5	1.7	1.6	2.0	$+0.0$ -2.7	$+2.4$ -1.1	± 4.7 -3.8
B [28, 29]	4.44	0.38	7.3	3.5	2.6	3.0	4.0	$+0.0$ -5.0	$+1.4$ -0.5	± 6.6 -5.5
C [30]	4.05	0.30	5.7	4.2	3.3	1.8	0.9	$+0.0$ -6.2	$+1.2$ -0.7	± 5.7 -6.9

A= M_x cut Belle, B= (M_x, q^2) cut Belle+Babar, C= E_l cut Babar

Overall theory errors are 5-9%, depending on the cuts.

Main theoretical desiderata

- know the b mass and the OPE pars precisely
lattice, $b \rightarrow c$ and $bs\gamma$ moments, pert calculations, goal 15-20 MeV for m_b
- study all the spectra of $b \rightarrow u \ell \nu$
to constrain WA and the SFs, complementary to OPE constraints
- be as inclusive as possible
to minimize dependence on functional forms

Present parametric is 3.5% with $\delta m_b \sim 40$ MeV, dominates cleanest cuts.
From $b \rightarrow c$ experience, duality violation should be small
Therefore, a 2% goal on $|V_{ub}|$ seems to be realistic.

Rare Decays


- $b \rightarrow s\gamma$ serves as prototype FCNC
 - Extremely important NP constraints (see later)
 - Precision of BF \sim theory errors ($\sim 7\%$)
 - Prospects to improve the theoretical precision?
 - Also, isospin & CP asymmetries, photon polarization, etc.
 - Discussed in the CDR, not updated

- Nonperturbative corrections $\Lambda^2/m_{b,c}^2$ to $\Gamma(\bar{B} \rightarrow X_s \gamma)$ are well under control
- However: Estimation of power corrections of $O(\alpha_s \Lambda/m_b)$ should be improved:
Largest uncertainty (5%) in our new NNLL prediction (see Lee et al)
- Further uncertainties: parametric (3%), higher-order (3%), mc-interpolation (3%)

$$BR(\bar{B} \rightarrow X_s \gamma)_{E_\gamma > 1.6 \text{ GeV}} = BR(\bar{B} \rightarrow X_c e \bar{\nu})^{\text{exp}} \left[\frac{\Gamma(b \rightarrow s \gamma)}{\Gamma(b \rightarrow ce \bar{\nu})} \right]_{\text{LO EW}} f \left(\frac{\alpha_s(M_W)}{\alpha_s(m_b)} \right) \times$$

$$\times \left\{ 1 + \underbrace{O(\alpha_s)}_{\text{NLO } \sim 25\%} + \underbrace{O(\alpha_s^2)}_{\text{NNLO } \sim 7\%} + \underbrace{O(\alpha_{\text{em}})}_{\sim 4\%} + \underbrace{O\left(\frac{\Lambda^2}{m_b^2}\right)}_{\sim 1\%} + \underbrace{O\left(\frac{\Lambda^2}{m_c^2}\right)}_{\sim 3\%} + \underbrace{O\left(\frac{\alpha_s \Lambda}{m_b}\right)}_{\sim 5\%} \right\}$$

- Additional sensitivities to nonperturbative physics due to necessary cuts in the photon energy spectrum to suppress the $B\bar{B}$ background:
Shape function methods and multi-scale SCET analysis
⇒ Additional theoretical uncertainties





Main source of theoretical uncertainty in $B(b \rightarrow s \gamma)$ at present.
If this can be removed, theory error could be 3%.

Inclusive, lepton tag: after selection




- BB background reduced with π^0 and η veto
- Still significant background remains at low E_γ
- Study inclusive π^0 and η production to tune MC of BB background

$\gamma\gamma$ invariant mass in bins of π^0 energy





Fits to data and MC

Inclusive, hadronic tags: E_γ Spectrum



efficiency correction

also takes care of E_γ resolution





Recoil Technique (II)

EXPECTED IMPROVEMENTS at SUPER-B

- The Super-B detector design would introduce significant improvement in the recoil performances;
- HERMITICITY:
 - helps to reduce the background when applying a cut on the track multiplicity in the recoil (see M. Mazur talk in Paris) → *30% bkg reduction is realistic*
 - modify the distribution of Eextra (the most important cut) → *effect to be established, see next slide*
- VERTEXING:
 - vertexing informations poorly used at present;
 - bkg reduction is probably possible applying vertexing requirements and secondary vertex informations;
- OTHER (PID, K_1^0 velo, etc.).



Recoil Technique (II)

EXPECTED IMPROVEMENTS at SUPER-B

- The Super-B detector design would introduce significant improvement in the recoil performances;

- HERMITICITY:

helps to
the tra
talk in

- modify
cut) →

- VERTEXING:

- vertexing informations poorly used at present;
- bkg reduction is probably possible applying vertexing requirements and secondary vertex informations;

- OTHER (PID, K_1^0 velo, etc.).

**Detailed Simulations
Needed
to have precise
estimates of improvements**

ing a cut on
M. Mazur

listic


important
slide

Caveats



Searches for modes with missing energy rely heavily on the low multiplicity of the signal to keep backgrounds manageable

- higher multiplicity requires harder cuts or additional kinematic handles
- Detector non-hermiticity and/or non-physics “junk” is a killer
- Tag reconstruction is usually CLEANER in these events, so can potentially use tag modes that appear to be too messy in “generic” B decays



Caveats



Tag B reconstruction efficiency scales non-linearly with tracking and calorimeter acceptance

- Not clear what the impact of reduced beam energy asymmetry will be

Need to balance reconstruction efficiency against cleanliness of the reconstruction by careful choice of what modes to reconstruct

- Potential for significant efficiency gain by including e.g. $D^0 \rightarrow K^+ \pi^- 2\pi^0$, or purity gain by including e.g. J/ψ - seeded decays

Devil is in the details!

- performance of vertexing, kinematic fitting, track-cluster matching, presence of tracking and/or calorimeter artifacts etc etc.

⇒ Important to study impact of proposed detector design

Simulation studies

- Sensitivity estimates for many interesting channels require simulation studies
- Huge effort by fastsimu group since Valencia
- Physics studies can and will benefit from this important work
 - Improved, detailed, dedicated studies necessary for TDR phase

Rare decays (cont.)

- $b \rightarrow sll$
 - Situation very different to $b \rightarrow sy$
 - Exclusive modes will be precisely measured by LHCb
 - Inclusive analyses theoretically much cleaner
 - Not yet possible at B factories
 - (semi-inclusive analyses have been done)
 - Requires SuperB statistics
 - $b \rightarrow sTT$ provides another interesting possibility

Recent proposal: normalization to semileptonic $B \rightarrow X_u \ell \nu$ decay rate **with the same cut** reduces the impact of $1/m_b$ corrections in the high- q^2 region significantly. [Ligeti, Tackmann, hep-ph/0707.1694](#)

Numerical results [Huber, Hurth, Lunghi](#)


$$\begin{aligned} \mathcal{R}(s_0)_{\mu\mu}^{\text{high}} &= 2.29 \times 10^{-3} \left(1 \pm 0.04_{\text{scale}} \pm 0.02_{m_t} \pm 0.01_{C, m_c} \pm 0.006_{m_b} \pm 0.005_{\alpha_s} \pm 0.09_{\text{CKM}} \right. \\ &\quad \left. \pm 0.003_{\lambda_2} \pm 0.05_{\rho_1} \pm 0.03_{f_u^0 + f_s} \pm 0.05_{f_u^0 - f_s} \right) \\ &= 2.29 \times 10^{-3} (1 \pm 0.13) \end{aligned}$$

Theoretical uncertainty in $B(b \rightarrow s \ell \ell)$ quite large at present.
But, in this approach, dominated by CKM factors.

- Zero of the forward-backward asymmetry q_0^2 :

$$(q_0^2)_{\mu\mu} = \left[3.50 \pm 0.10_{\text{scale}} \pm 0.002_{m_t} \pm 0.04_{m_c, C} \right. \\ \left. \pm 0.05_{m_b} \pm 0.03_{\alpha_s(M_Z)} \pm 0.001_{\lambda_1} \pm 0.01_{\lambda_2} \right] \text{GeV}^2 = (3.50 \pm 0.12) \text{GeV}^2$$

$$(q_0^2)_{ee} = \left[3.38 \pm 0.09_{\text{scale}} \pm 0.002_{m_t} \pm 0.04_{m_c, C} \right. \\ \left. \pm 0.04_{m_b} \pm 0.03_{\alpha_s(M_Z)} \pm 0.002_{\lambda_1} \pm 0.01_{\lambda_2} \right] \text{GeV}^2 = (3.38 \pm 0.11) \text{GeV}^2$$



Forward-backward asymmetry in inclusive $b \rightarrow sll$
Theoretically very clean! (Much better than exclusive)


More rare decays

- $b \rightarrow svv$ and $B \rightarrow lv$ provide further examples of rare decay modes where the full statistics of SuperB is required to reach the interesting sensitivity region
 - Expect observation of $B \rightarrow K(*)vv$ above $10/ab$
 - Precision of $B \rightarrow \mu\nu$ overtakes that of $B \rightarrow \tau\nu$ (systematics limited) around the same time



$B^+ \rightarrow K^+ \nu \nu$ (II)

RESULTS at SUPER-B



Observation between 10 and 20 ab^{-1} ;


Exp. error ~ theoretical error around 30 ab^{-1} ;

18% error at 50 ab^{-1} in the most conservative scenario.

Tagged $B^+ \rightarrow l^+ \nu$ ($l = e, \mu$)



- Signal lies at kinematic endpoint in lepton p^* for B decays, hence essentially no B background
- Continuum background can produce high p^* leptons, but this background can be directly determined from data using the tag B m_{ES} sideband
- Narrow signal peak would lead to a very compelling signal with a very small number of events



	$e^+ \nu$	$\mu^+ \tilde{\nu}$
$\epsilon_{\text{tot}} \times 10^5$	135 ± 4	120 ± 4
$n_b^* \text{ MC}$	2.66 ± 0.13	5.74 ± 0.25
n_b^*	2.67 ± 0.19	5.67 ± 0.34
n_s^*	-0.07 ± 0.03	-0.11 ± 0.05
$\mathcal{B} \times 10^{-6}$	$-0.1^{+2.6}_{-1.7}$	$-0.2^{+2.7}_{-1.8}$
$\mathcal{B}^{90\% \text{ C.L.}}$	5.2×10^{-6}	5.6×10^{-6}

Leptonic Decays





- $B \rightarrow \tau \nu$ will be limited by systematics at some stage
 - CDR estimate of precision may be optimistic
- $B \rightarrow \mu \nu$ does **not** suffer the same problem
 - Will become “golden channel” for leptonic B decays at some luminosity (likely above $10/\text{ab}$)
- Also allows universality test

Golden modes

	H^+	Minimal	Non-Minimal	Non-Minimal	NP	Right-Handed
	high $\tan\beta$	FV	FV (1-3)	FV (2-3)	Z-penguins	currents
$\mathcal{B}(B \rightarrow X_s \gamma)$		X		O		O
$A_{CP}(B \rightarrow X_s \gamma)$				X		O
$\mathcal{B}(B \rightarrow \tau \nu)$	<i>X-CKM</i>					
$\mathcal{B}(B \rightarrow X_s l^+ l^-)$				O	O	O
$\mathcal{B}(B \rightarrow K \nu \bar{\nu})$				O	X	
$S(K_S \pi^0 \gamma)$						X
β			<i>X-CKM</i>			O

- Which channel is “golden” for new physics discovery depends on the particular brand of NP
 - Many others can be added to the list
 - [add your favourite here]
 - ... however, the request was to make a selection

Improvements in $B \rightarrow l\nu$ with Luminosity



Comparison with KEKB upgrade

Mode	Sensitivity		
	Current	10 ab^{-1}	75 ab^{-1}
$\mathcal{B}(B \rightarrow X_s \gamma)$	7%	5%	3%
$A_{CP}(B \rightarrow X_s \gamma)$	0.037	0.01	0.004–0.005
$\mathcal{B}(B^+ \rightarrow \tau^+ \nu)$	30%	10%	3–4%
$\mathcal{B}(B^+ \rightarrow \mu^+ \nu)$	X	20%	5–6%
$\mathcal{B}(B \rightarrow X_s l^+ l^-)$	23%	15%	4–6%
$A_{FB}(B \rightarrow X_s l^+ l^-)_{s_0}$	X	30%	4–6%
$\mathcal{B}(B \rightarrow K \nu \bar{\nu})$	X	X	16–20%
$S(K_S^0 \pi^0 \gamma)$	0.24	0.08	0.02–0.03

Benchmarks and LHC Interplay



- Work done mainly by F.Ronga & S.Heinemayer
- Many thanks for their efforts and responses to our requests ...
- Work on interplay between flavour and LHC is ongoing



Flavour as a Window to New Physics at the LHC 5 May - 13 June 2008

<http://ph-dep-th.web.cern.ch/ph-dep-th/content2/THInstitutes/2008/flavour/TH-Flavour.html>
<http://indico.cern.ch/conferenceOtherViews.py?view=standard&confId=31959>

Thursday 29 May 2008 [top](#)



10:00->12:00 **Week IV: B@LHC Focus Week: Charm, Tau, New States, ...** (Location: [13-2-005](#))



10:00 Charm Physics at LHCb (30')  ) Raluca Muresan

10:30 Charm Physics (Theory) (30')  ) Alexei Petrov

11:00 Discussion (1h00')

14:00->18:00 **Week IV: B@LHC Focus Week: Spectroscopy, Lambda_b, |V(xb)|, ...**

14:00 Non-CP Physics at LHCb (30')  ) Michael Schmelling

14:30 Determination of Vcb and Vub (Theory) (30')  ) Nikolai Uraltsev



15:00 Discussion (30')



15:30 **Coffe Break**



16:00 Special Seminar: Where to expect New Physics in flavour-violating processes? (1h00') Andrzej Buras

Friday 30 May 2008 [top](#)




09:00->12:00 **Week IV: B@LHC Focus Week: Long-Term Future**



09:00 LHCb upgrade (20')  ) Jim Libby

09:20 Status and Prospects of sKEKB (20')  ) Yutaka Ushiroda

09:40 The SuperB Project: an overview (20')  ) Marcello Giorgi

10:00 **Coffee Break**

10:30 Non-Perturbative QCD Calculations for B Physics (and related topics) (30')   ) Guido Martinelli

11:00 Theory Outlook (30')  ) Zoltan Ligeti

11:30 Final Discussion (30')

Done 5.658s Tor Disabled

The SPS1a benchmark point

A (too) good point for LHC!

$$M_0 = +100 \text{ GeV}/c^2$$

$$M_{1/2} = +250 \text{ GeV}/c^2$$

$$A_0 = -100 \text{ GeV}/c^2$$

$$\tan \beta = +10$$

$$\text{sign}(\mu) = +1$$

Allows cascade decay


$$\tilde{q}_L \rightarrow \tilde{\chi}_2^0 q \rightarrow \tilde{\ell}_R \ell q \rightarrow \tilde{\chi}_1^0 \ell \ell q$$

for "edge" measurements:

$$(m_{\tilde{t}_1}^2)^{\text{edge}} = \frac{(m_{\tilde{\chi}_2^0}^2 - m_{\tilde{l}_R}^2)(m_{\tilde{l}_R}^2 - m_{\tilde{\chi}_1^0}^2)}{m_{\tilde{l}_R}^2}$$

$$(m_{\tilde{q}_L}^2)^{\text{edge}} = \frac{(m_{\tilde{q}_L}^2 - m_{\tilde{\chi}_2^0}^2)(m_{\tilde{\chi}_2^0}^2 - m_{\tilde{\chi}_1^0}^2)}{m_{\tilde{\chi}_2^0}^2}$$

$$(m_{\tilde{q}_L}^2)_{\text{min}} = \frac{(m_{\tilde{q}_L}^2 - m_{\tilde{\chi}_2^0}^2)(m_{\tilde{\chi}_2^0}^2 - m_{\tilde{l}_R}^2)}{m_{\tilde{\chi}_2^0}^2}$$



SUSY spectrum at SPS1a

Note: SPS1a is close to the overall preferred minimum with today's data.

LHC performance at SPS1a [hep-ph/0410364]

Performance based on 300/fb (2014)

SUSY spectrum [GeV]

	Mass	Error
$\tilde{\chi}_1^0$	96.9	4.8
$\tilde{\chi}_2^0$	179.8	4.7
$\tilde{\chi}_4^0$	375.6	5.1
\tilde{e}_R	144.1	4.8
\tilde{e}_L	202.6	5.0
$\tilde{\mu}_R$	144.1	4.8
$\tilde{\mu}_L$	202.6	5.0
$\tilde{\tau}_1$	134.7	8.0
\tilde{q}_R	547.5	12.0
\tilde{q}_L	565.0	8.7
\tilde{b}_1	514.9	7.5
\tilde{b}_2	544.1	7.9
\tilde{g}	608.0	8.0
h^0	112.9	0.25

Edge measurements [GeV]

$(m_{\ell\ell})^{\text{edge}}$	58.878	0.085
$(m_{q\ell\ell})^{\text{edge}}$	451.1	4.5
$(m_{q\ell})_{\text{min}}^{\text{edge}}$	317.5	3.1

⇒ Impact on CMSSM parameters

- include this spectrum as constraints;
- combine with today's constraints;
- get best fit values and errors:

$$M_0 = 100.0 \pm 1.5$$

$$M_{1/2} = 250.0 \pm 1.1$$

$$A_0 = 100 \pm 30$$


$$\tan \beta = 9.8 \pm 1.2$$

All ideal masses generated by SoftSusy

Flavour Physics predictions


Strong impact of LHC constraints on Flavour Sector!

$$\begin{aligned}
 R(B \rightarrow s\gamma) &= 1.063 \pm 0.022 \\
 R(\Delta m_s) &= 1.0582 \pm 0.0007 \\
 R(B \rightarrow \tau\nu) &= 0.970 \pm 0.007 \\
 R(B \rightarrow X_s \ell\ell) &= 0.910 \pm 0.003 \\
 R(\Delta m_s / \Delta m_d) &= 0.99988 \pm 0.00005 \\
 B_s \rightarrow \mu\mu &= 2.736e-09 \pm 0.066e-9 \\
 B_d \rightarrow \mu\mu &= 1.580e-10 \pm 0.038e-10
 \end{aligned}$$



But...

- this point is especially good for L(H)C;
- we assumed MFV;
- correlations are not taken into account.



SPS 4: mSUGRA scenario with large $\tan \beta$

The large value of $\tan \beta$ in this scenario has an important impact on the phenomenology in the Higgs sector. The couplings of A, H to $b\bar{b}$ and $\tau^+\tau^-$ as well as the $H^\pm t\bar{b}$ couplings are significantly enhanced in this scenario, resulting in particular in large associated production cross sections for the heavy Higgs bosons.

Point:

$$m_0 = 400 \text{ GeV}, \quad m_{1/2} = 300 \text{ GeV}, \quad A_0 = 0, \quad \tan \beta = 50, \quad \mu > 0.$$

This point equals mSUGRA point 3 of the “Points d’Aix” and is similar to BDEGMOPW point 1.

Expect observable flavour signatures for these MFV points

SPS 5: mSUGRA scenario with relatively light scalar top quark

This scenario is characterized by a large negative value of A_0 , which allows consistency of the relatively low value of $\tan \beta$ with the constraints from the Higgs search at LEP, see Ref. [34].

Point:

$$m_0 = 150 \text{ GeV}, \quad m_{1/2} = 300 \text{ GeV}, \quad A_0 = -1000, \quad \tan \beta = 5, \quad \mu > 0.$$

This point equals mSUGRA point 4 of the “Points d’Aix”.

More Snowmass Points

	SPS1a	SPS4	SPS5
$\mathcal{R}(B \rightarrow X_s \gamma)$	0.919 ± 0.038	0.248	0.848 ± 0.081
$\mathcal{R}(B \rightarrow \tau \nu)$	0.968 ± 0.007	0.436	0.997 ± 0.003
$\mathcal{R}(B \rightarrow X_s l^+ l^-)$	0.916 ± 0.004	0.917	0.995 ± 0.002
$\mathcal{R}(B \rightarrow K \nu \bar{\nu})$	0.967 ± 0.001	0.972	0.994 ± 0.001
$\mathcal{B}(B_d \rightarrow \mu^+ \mu^-)/10^{-10}$	1.631 ± 0.038	16.9	1.979 ± 0.012
$\mathcal{R}(\Delta m_s)$	1.050 ± 0.001	1.029	1.029 ± 0.001
$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/10^{-9}$	2.824 ± 0.063	29.3	3.427 ± 0.018
$\mathcal{R}(K \rightarrow \pi^0 \nu \bar{\nu})$	0.973 ± 0.001	0.977	0.994 ± 0.001

No LHC sensitivity estimates \Leftrightarrow no uncertainties

Already excluded!

How to Add Flavour Structure?

- Explicit models (T.Shindou & collaborators)
 - mSUGRA (MFV)
 - MSSM + r-h-neutrinos
 - SU(5) SUSY with r-h-neutrinos
 - MSSM with U(2) flavour symmetry
- Alternative, general, approach:
 - MSSM with mass-insertions
 - M.Ciuchini & L.Silvestrini, work in CDR, update ongoing

MSSM

SuperB


$$M_{\tilde{d}}^2 = \begin{pmatrix} m_{\tilde{d}_L}^2 & m_{\tilde{d}_L} m_{\tilde{d}_R} (A_d - \mu \tan \beta) & (\Delta_{12}^d)_{LL} & (\Delta_{12}^d)_{LR} & (\Delta_{13}^d)_{LL} & (\Delta_{13}^d)_{LR} \\ m_{\tilde{d}_L} m_{\tilde{d}_R} (A_d - \mu \tan \beta) & m_{\tilde{d}_R}^2 & (\Delta_{12}^d)_{RL} & (\Delta_{12}^d)_{RR} & (\Delta_{13}^d)_{RL} & (\Delta_{13}^d)_{RR} \\ (\Delta_{12}^d)_{LL} & (\Delta_{12}^d)_{RL} & m_{\tilde{s}_L}^2 & m_{\tilde{s}_L} m_{\tilde{s}_R} (A_s - \mu \tan \beta) & (\Delta_{23}^d)_{LL} & (\Delta_{23}^d)_{LR} \\ (\Delta_{12}^d)_{LR} & (\Delta_{12}^d)_{RR} & m_{\tilde{s}_L} m_{\tilde{s}_R} (A_s - \mu \tan \beta) & m_{\tilde{s}_R}^2 & (\Delta_{23}^d)_{RL} & (\Delta_{23}^d)_{RR} \\ (\Delta_{13}^d)_{LL} & (\Delta_{13}^d)_{RL} & (\Delta_{23}^d)_{LL} & (\Delta_{23}^d)_{LR} & m_{\tilde{b}_L}^2 & m_{\tilde{b}_L} m_{\tilde{b}_R} (A_b - \mu \tan \beta) \\ (\Delta_{13}^d)_{LR} & (\Delta_{13}^d)_{RR} & (\Delta_{23}^d)_{LR} & (\Delta_{23}^d)_{RR} & m_{\tilde{b}_L} m_{\tilde{b}_R} (A_b - \mu \tan \beta) & m_{\tilde{b}_R}^2 \end{pmatrix}$$

LHC, ILC - HE frontier



Mass Insertions
 $(\delta_{ij}^d)_{AB} = (\Delta_{ij}^d)_{AB} / m_{\tilde{q}}^2$

3σ from 0 sensitivity plot

- i) sensit. to $\Lambda < 20$ TeV
- ii) sensit. to $|(\delta_{23}^d)_{LR}| > 10^{-2}$ for $\Lambda < 1$ TeV





BR(B → X_sγ)
 BR(B → X_sll)
 A_{CP}(B → X_sγ)
 all together




Im(δ_{23}^d)_{LR} vs Re(δ_{23}^d)_{LR}
 Reconstruction of
 $(\delta_{23}^d)_{LR} = 0.028 e^{i\pi/4}$ for
 $\Lambda = m_{\tilde{g}} = m_{\tilde{q}} = 1$ TeV

More comparison with KEKB upgrade



Re vs Im parts of MSSM mass insertion parameter $(\delta_{13}^d)_{LL}$


$$S_{CP}(B_d \rightarrow K^* \gamma)$$


mSUGRA

U(2)

SU(5) Non-Degen. (I)

Correlation between ϕ_3 and $\Delta m_{B_s}/\Delta m_{B_d}$



mSUGRA

U(2)

Need $<1^\circ$
precision on γ
in UT fit

Summary

- Thanks, once again, to the local organizers of the Valencia workshop
- The physics case has been made:
 - Conceptual Design Report
 - Flavour in the LHC Era write-up (arXiv:0801.1833,26)
 - Report from Valencia
 - Several other documents
- Further refinements always possible, but what we really need now is a machine, detector, and lots and lots of lovely luminosity!