

SECOND ECFA WORKSHOP on e^+e^- Higgs / Electroweak / Top Factories

DISCUSSION STARTER

FOCUS TOPIC “BSM TOP (DECAY)”

FOCUS GROUP: NUÑO CASTRO, MARINA COBAL, GAUTHIER DURIEUX, RF,
MARÍA TERESA, NÚÑEZ PARDO DE VERA, KIRILL SKOVPEN, MARCEL VOS


OCT 11TH 2023, PAESTUM

ECFA-WG1-SRCH: ALEKSANDER FILIP ZARNECKI,
REBECA GONZALEZ SUAREZ, ROBERTO FRANCESCHINI




The most obvious way to study the top quark is to produce it in large numbers

⇒ Make a Top factory



pp

$\sigma_{pp \rightarrow t\bar{t}} \simeq 1 \text{ nb}$ at 14 TeV means 10^9 top quarks per ab^{-1} at HL-LHC



e^+e^-

$\sigma_{e^+e^- \rightarrow t\bar{t}} \simeq 1 \text{ pb}$ at 0.365 TeV means 10^6 top quarks per ab^{-1} at the top factory



obvious complementarities,

- where the HL-LHC suffers too busy events the top factory can shine
- where the top factory lacks statistics the HL-LHC can supply

also “single” production modes can bring very useful info

$pp \rightarrow t + X$ e.g. can measure weak couplings

$e^+e^- \rightarrow tq$ can probe top quark contact interactions (even at energies below the top quark pair threshold)

also “boosted” production modes can bring very useful info

$pp \rightarrow tt$ still O(pb) for $p_{T,top} \geq 500 \text{ GeV}$ at LHC14

$pp \rightarrow ttj$ still O(pb) for $p_{T,jet} \geq 500 \text{ GeV}$ at LHC14

also “associated” production modes can bring very useful info

$pp \rightarrow ttZ, tth$ unique routes to access couplings
weak and Yukawa couplings


very articulated field with lots of cross-talk between pp and e^+e^-

Top quark decay


Traditional topic for top factories is the possibility of BSM decays

Especially when the direct reach was limited (e.g. LEP and TeVatron times) the top quark might have been our “window” to new physics

e.g. $t \rightarrow H^+ b$ in MSSM or general 2HDM



e.g. $t \rightarrow Vq$ in MSSM or general 2HDM and many other models



Such light H^+ is hardly tenable* these days. The possibility of flavor violation, instead, is more subtle.

Charged Higgs direct searches




Fig. 4. Upper limit on $B(t \rightarrow H^+ b)$ for the simultaneous fit of $B(t \rightarrow H^+ b)$ and $\sigma_{t\bar{t}}$ versus M_{H^+} . The yellow band shows the ± 1 SD band around the expected limit. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)







Fig. 4 The extracted 95% CL upper limits on $\mathcal{B}(t \rightarrow H^+ b)$ are shown in the range of the charged Higgs mass from 90 GeV to 150 GeV. The limits shown are calculated using the CL_s limit-setting procedure.



Good top bad top



Good top bad top



Top quark decay at the Top Factory


$t \rightarrow BSM$

1311.2028 - Agashe et al. - Snowmass 2013 Top quark working group report

Table 1-7. SM and new physics model predictions for branching ratios of top FCNC decays. The SM predictions are taken from [119], on 2HDM with flavor violating Yukawa couplings [119, 120] (2HDM (FV) column), the 2HDM flavor conserving (FC) case from [121], the MSSM with 1TeV squarks and gluinos from [122], the MSSM for the R-parity violating case from [123, 124], and warped extra dimensions (RS) from [125, 126].

Process	SM	2HDM(FV)	2HDM(FC)	MSSM	RPV	RS
$t \rightarrow Zu$	7×10^{-17}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow Zc$	1×10^{-14}	$\leq 10^{-6}$	$\leq 10^{-10}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-5}$
$t \rightarrow gu$	4×10^{-14}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow gc$	5×10^{-12}	$\leq 10^{-4}$	$\leq 10^{-8}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-10}$
$t \rightarrow \gamma u$	4×10^{-16}	–	–	$\leq 10^{-8}$	$\leq 10^{-9}$	–
$t \rightarrow \gamma c$	5×10^{-14}	$\leq 10^{-7}$	$\leq 10^{-9}$	$\leq 10^{-8}$	$\leq 10^{-9}$	$\leq 10^{-9}$
$t \rightarrow hu$	2×10^{-17}	6×10^{-6}	–	$\leq 10^{-5}$	$\leq 10^{-9}$	–
$t \rightarrow hc$	3×10^{-15}	2×10^{-3}	$\leq 10^{-5}$	$\leq 10^{-5}$	$\leq 10^{-9}$	$\leq 10^{-4}$

2HDM help needed



- Most references are on complete models, some from 1990s
- Correct formulae do not expire.
- But phenomenological interpretations of correct formulae do expire when high energy (LHC) and high-intensity (flavor) experiments work hard to search NP!
- Most recent searches of NP express results in “simplified models”, small brothers of the full microscopic models, used to can capture salient feature of a relatively broad class of models.
- Translation and merging of info is needed to really update these specific models

Decays

FCNC EW

2301.11605 - ATLAS






Table 8: Observed and expected 95% CL limits on the FCNC $t \rightarrow Zq$ branching ratios and the effective coupling strengths for different vertices and couplings (top eight rows). For the latter, the energy scale is assumed to be $\Lambda_{\text{NP}} = 1$ TeV. The bottom rows show, for the case of the FCNC $t \rightarrow Zu$ branching ratio, the observed and expected 95% CL limits when only one of the two SRs, either SR1 or SR2, and all CRs are included in the likelihood.

Observable	Vertex	Coupling	Observed	Expected
SRs+CRs				
$\mathcal{B}(t \rightarrow Zq)$	tZu	LH	6.2×10^{-5}	$4.9^{+2.1}_{-1.4} \times 10^{-5}$
$\mathcal{B}(t \rightarrow Zq)$	tZu	RH	6.6×10^{-5}	$5.1^{+2.1}_{-1.4} \times 10^{-5}$
$\mathcal{B}(t \rightarrow Zq)$	tZc	LH	13×10^{-5}	$11^{+5}_{-3} \times 10^{-5}$
$\mathcal{B}(t \rightarrow Zq)$	tZc	RH	12×10^{-5}	$10^{+4}_{-3} \times 10^{-5}$
$ C_{uW}^{(13)*} $ and $ C_{uB}^{(13)*} $	tZu	LH	0.15	$0.13^{+0.03}_{-0.02}$
$ C_{uW}^{(31)} $ and $ C_{uB}^{(31)} $	tZu	RH	0.16	$0.14^{+0.03}_{-0.02}$
$ C_{uW}^{(23)*} $ and $ C_{uB}^{(23)*} $	tZc	LH	0.22	$0.20^{+0.04}_{-0.03}$
$ C_{uW}^{(32)} $ and $ C_{uB}^{(32)} $	tZc	RH	0.21	$0.19^{+0.04}_{-0.03}$
SR1+CRs				
$\mathcal{B}(t \rightarrow Zq)$	tZu	LH	9.7×10^{-5}	$8.6^{+3.6}_{-2.4} \times 10^{-5}$
$\mathcal{B}(t \rightarrow Zq)$	tZu	RH	9.5×10^{-5}	$8.2^{+3.4}_{-2.3} \times 10^{-5}$
SR2+CRs				
$\mathcal{B}(t \rightarrow Zq)$	tZu	LH	7.8×10^{-5}	$6.1^{+2.7}_{-1.7} \times 10^{-5}$
$\mathcal{B}(t \rightarrow Zq)$	tZu	RH	9.0×10^{-5}	$6.6^{+2.9}_{-1.8} \times 10^{-5}$



Present $BR < 10^{-4}$, HL-LHC might leave little room to improve (projection for HL-LHC should be around few 10^{-5} for $Z \rightarrow \ell\ell$ according to ATL-PHYS-PUB-2019-001)

FCNC Higgs

1707.01404 - ATLAS 2112.09734, 2111.02219 CMS



No significant excess is observed and an upper limit is set on the $t \rightarrow cH$ branching ratio of 2.2×10^{-3} at the 95% confidence level, while the expected limit in the absence of signal is 1.6×10^{-3} . The corresponding limit on the $t\bar{c}H$ coupling is 0.090 at the 95% confidence level. The observed upper limit on the $t \rightarrow uH$ branching ratio is 2.4×10^{-3} .




Present $BR(t \rightarrow cH) < 2.2 \cdot 10^{-3}$ @ 36 fb^{-1} , $9.4 \cdot 10^{-4}$ @ 137 fb^{-1} , HL-LHC might go to $3 \cdot 10^{-4}$ (TBC), which leaves little room for the top factory, e.g. 500 fb^{-1} at CLIC380 gives is $BR < 2 \cdot 10^{-4}$

[1810.05487] Expected 95% C.L. limits for 500 fb⁻¹ collected at 380 GeV CLIC are: $BR(t \rightarrow c\gamma) < 4.7 \cdot 10^{-5}$, $BR(t \rightarrow cH) \times BR(H \rightarrow bb) < 1.2 \cdot 10^{-4}$ and $BR(t \rightarrow c\text{Emiss}) < 1.2 - 4.1 \cdot 10^{-4}$.




$t \rightarrow au_j, a \rightarrow uu, cc, uc$

2101.07803, 2202.09371 - Carmona, Elahi, Scherb, Schwaller



$$\mathcal{L} = \frac{1}{2}(\partial_\mu a)(\partial^\mu a) - \frac{m_a^2}{2}a^2 + \frac{\partial_\mu a}{f_a} [(c_{uR})_{ij} \bar{u}_R \gamma^\mu u_{Rj}]$$

$$C_{uR} \sim \begin{pmatrix} 2 & 3\epsilon & 3\epsilon^2 \\ 3\epsilon & 1 & \epsilon \\ 3\epsilon^2 & \epsilon & \epsilon^2 \end{pmatrix}$$




focus on displaced flavor-conserving decay so far

Ample room for improvement with top factory studies for prompt decays $a \rightarrow jj, uc, cc$ and invisible

$t \rightarrow \phi q, \phi\phi q, \phi \rightarrow \ell\ell$








2005.09594 - Castro, Chala, Peixoto, Ramos

- BP 1 : $\mathbf{Y}_{i3}^q = \mathbf{Y}_{3i}^q = 0.01$, $\Lambda = 5 \text{ TeV} \implies \mathcal{B}(t \rightarrow Sq) \sim 10^{-8} - 10^{-7}$,
- BP 2 : $\mathbf{Y}_{i3}^q = \mathbf{Y}_{3i}^q = 0.10$, $\Lambda = 5 \text{ TeV} \implies \mathcal{B}(t \rightarrow Sq) \sim 10^{-6} - 10^{-5}$,
- BP 3 : $\mathbf{Y}_{i3}^q = \mathbf{Y}_{3i}^q = 0.10$, $\Lambda = 1 \text{ TeV} \implies \mathcal{B}(t \rightarrow Sq) \sim 10^{-4} - 10^{-3}$,
- BP 4 : $\tilde{\mathbf{Y}}_{i3}^q = \tilde{\mathbf{Y}}_{3i}^q = 1.00$, $\Lambda = 5 \text{ TeV} \implies \mathcal{B}(t \rightarrow SSq) \sim 10^{-11} - 10^{-8}$,
- BP 5 : $\tilde{\mathbf{Y}}_{i3}^q = \tilde{\mathbf{Y}}_{3i}^q = 0.20$, $\Lambda = 1 \text{ TeV} \implies \mathcal{B}(t \rightarrow SSq) \sim 10^{-10} - 10^{-7}$,
- BP 6 : $\tilde{\mathbf{Y}}_{i3}^q = \tilde{\mathbf{Y}}_{3i}^q = 1.00$, $\Lambda = 1 \text{ TeV} \implies \mathcal{B}(t \rightarrow SSq) \sim 10^{-8} - 10^{-5}$,



focus on lepto-philic so far

hadro-philic would require some model building



LHC

Associated productions

e^+e^- at $\sqrt{s} = 350, 365, 380$ GeV

Associated $t\bar{t}a, a \rightarrow \mu\mu$

2304.14247 - ATLAS






Table 2: Definition of the mass requirements on m






	Signal Regions		on-Z Control Region		$t\bar{t}$ Control Region
Channel	$e\mu\mu$	$\mu\mu\mu$	$e\mu\mu$	$\mu\mu\mu$	$e\mu\mu$
Binning	$m_{\mu\mu}^a$	$m_{\mu\mu}^a$	$n_{\text{jets}}, n_{b\text{-jets}}$	$n_{\text{jets}}, n_{b\text{-jets}}$	$p_T^{\mu, \text{fake}}$
$n_{\text{electrons}}$	1	0	1	0	1
n_{muons}	2	3	2	3	2
$m_{\mu\mu}$ [GeV]	$12 < m_{\mu\mu}^a < 77$	$12 < m_{\mu\mu}^a < 77$ and $m_{\mu\mu}^{\text{other}} < 77$ or > 107	$77 < m_{\mu\mu}^a < 107$	$77 < m_{\mu\mu}^a < 107$ or $77 < m_{\mu\mu}^{\text{other}} < 107$	$12 < m_{\mu\mu}^a < 77$
n_{jets}	≥ 3				1 or 2
$n_{b\text{-jets}}$	≥ 1				1

so far limited to muon final states and moderate mass 12-77 GeV

ample room for improvement outside the probed mass range and in other a decay final states, e.g. hadrons

$t\bar{t}a, a \rightarrow bb$

1507.07004 - Casolino, Farooque, Juste, Liu, Spannowsky



LHC bounds need to be interpreted

95% CL limit on \tilde{g}_t assuming $BR(a \rightarrow bb) = 1$

$\mathcal{L} \text{ (fb}^{-1}\text{)}$	$m_A \text{ (GeV)}$					
	20	30	40	60	80	100
1	2.73	2.14	2.18	2.48	2.82	3.65
30	1.31	0.94	0.95	1.10	1.23	1.62
100	1.06	0.72	0.71	0.83	0.93	1.25
300	0.88	0.57	0.55	0.65	0.72	1.00
3000	0.54	0.35	0.34	0.39	0.43	0.67

e^+e^- study missing

$e^+ e^- \rightarrow \gamma \rightarrow t\bar{t}$

$j j c b$
 $i c c b$
invisible

ample room for improvement outside the probed mass range and in other a decay final states, e.g. hadrons

Thank you!

BSM decays of the top quark


Focus Topic “*EXtt*”

- Update of the tenable FCNC BR incorporating direct searches for the microscopic BSM states that mediate flavor changing transitions, e.g. squarks
- Identification of more general patterns of BSM decay of the top quark in light of recent model building and of the new constraints
- Identification of blindspots for the LHC, ideally pursuing both signature-driven and model-driven routes
- Establishing the sensitivity of experiments at the top factory with realistic detector treatment (Delphes via KEY4HEP)
- Establishing connection with other stages of the HET program (e.g. single top production at 240 GeV)

Top quark decay at the Top Factory

$$t \rightarrow BSM$$


- BSM means SM EW final states



Top quark decay at the Top Factory

$$t \rightarrow BSM$$

Kaustubh Agashe and Sagar Airen at U. of Maryland agreed to help with the update





- Last refresh of BSM benchmark is quite old (2013)
- Regardless of the focus topics a refresh seems needed for the final report
- Quick assessment concluded that most results may need a retouch
- Inquiry for update/revalidation of the BSM benchmarks has started
- Relation to EFT to be investigated further

Top quark decay at the Top Factory

$$t \rightarrow BSM$$

1311.2028 - Agashe et al. - Snowmass 2013 Top quark working group report



Top quark decay at the Top Factory

$t \rightarrow BSM$

1311.2028 - Agashe et al. - Snowmass 2013 Top quark working group report


Table 1-7. SM and new physics model predictions for branching ratios of top FCNC decays. The SM predictions are taken from [119], on 2HDM with flavor violating Yukawa couplings [119, 120] (2HDM (FV) column), the 2HDM flavor conserving (FC) case from [121], the MSSM with 1TeV squarks and gluinos from [122], the MSSM for the R-parity violating case from [123, 124], and warped extra dimensions (RS) from [125, 126].

Process	SM	2HDM(FV)	2HDM(FC)	MSSM	RPV	RS
$t \rightarrow Zu$	7×10^{-17}	-	-	$\leq 10^{-7}$	$\leq 10^{-6}$	-
$t \rightarrow Zc$	1×10^{-14}	$\leq 10^{-6}$	$\leq 10^{-10}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-5}$
$t \rightarrow gu$	4×10^{-14}	-	-	$\leq 10^{-7}$	$\leq 10^{-6}$	-
$t \rightarrow gc$	5×10^{-12}	$\leq 10^{-4}$	$\leq 10^{-8}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-10}$
$t \rightarrow \gamma u$	4×10^{-16}	-	-	$\leq 10^{-8}$	$\leq 10^{-9}$	-
$t \rightarrow \gamma c$	5×10^{-14}	$\leq 10^{-7}$	$\leq 10^{-9}$	$\leq 10^{-8}$	$\leq 10^{-9}$	$\leq 10^{-9}$
$t \rightarrow hu$	2×10^{-17}	6×10^{-6}	-	$\leq 10^{-5}$	$\leq 10^{-9}$	-
$t \rightarrow hc$	3×10^{-15}	2×10^{-3}	$\leq 10^{-5}$	$\leq 10^{-5}$	$\leq 10^{-9}$	$\leq 10^{-4}$

observable at top factory

not observable at top factory

Use MSSM instead




typical 2HDM contribution

$$BR \sim (1/M_{2HDM})^4$$

Even a mere factor 2 stronger bounds on the particles originating flavor violation makes a factor 16 in the FCNC BR. This can take a “border-line observable at top factory” BR=10⁻⁵ down to 10⁻⁶ and ruin the party.

Top quark decay at the Top Factory

$t \rightarrow BSM$



$$BR \sim \left(1/M_{NP}\right)^4$$

Even a mere factor 2 stronger bounds on the particles originating flavor violation makes a factor 16 in the FCNC BR. This can take a “border-line observable at top factory” $BR=10^{-5}$ down to 10^{-6} and ruin the party.

Top quark decay at the Top Factory

$t \rightarrow BSM$

1311.2028 - Agashe et al. - Snowmass 2013 Top quark working group report

Table 1-7. SM and new physics model predictions for branching ratios of top FCNC decays. The SM predictions are taken from [119], on 2HDM with flavor violating Yukawa couplings [119, 120] (2HDM (FV) column), the 2HDM flavor conserving (FC) case from [121], the MSSM with 1TeV squarks and gluinos from [122], the MSSM for the R-parity violating case from [123, 124], and warped extra dimensions (RS) from [125, 126].

Process	SM	2HDM(FV)	2HDM(FC)	MSSM	RPV	RS
$t \rightarrow Zu$	7×10^{-17}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow Zc$	1×10^{-14}	$\leq 10^{-6}$	$\leq 10^{-10}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-5}$
$t \rightarrow gu$	4×10^{-14}	–	–	$\leq 10^{-7}$	$\leq 10^{-6}$	–
$t \rightarrow gc$	5×10^{-12}	$\leq 10^{-4}$	$\leq 10^{-8}$	$\leq 10^{-7}$	$\leq 10^{-6}$	$\leq 10^{-10}$
$t \rightarrow \gamma u$	4×10^{-16}	–	–	$\leq 10^{-8}$	$\leq 10^{-9}$	–
$t \rightarrow \gamma c$	5×10^{-14}	$\leq 10^{-7}$	$\leq 10^{-9}$	$\leq 10^{-8}$	$\leq 10^{-9}$	$\leq 10^{-9}$
$t \rightarrow hu$	2×10^{-17}	6×10^{-6}	–	$\leq 10^{-5}$	$\leq 10^{-9}$	–
$t \rightarrow hc$	3×10^{-15}	2×10^{-3}	$\leq 10^{-5}$	$\leq 10^{-5}$	$\leq 10^{-9}$	$\leq 10^{-4}$

observable at top factory


not observable at top factory

Use MSSM instead

- Most references are on complete models, some from 1990s
- Correct formulae do not expire.
- But phenomenological interpretations of correct formulae do expire when high energy (LHC) and high-intensity (flavor) experiments work hard to search NP!
- Most recent searches of NP express results in “simplified models”, small brothers of the full microscopic models, used to can capture salient feature of a relatively broad class of models.
- Translation and merging of info is needed to really update these specific models

MSSM RPC squarks


$$\tilde{q} \rightarrow q \text{ mET}$$



BRs in the MSSM may undergo a reduction due to squarks limits at 2+ TeV

MSSM-RPV squarks


$$\tilde{q} \rightarrow qq$$



RPV MSSM seems to have less stringent bounds, so it might re-enable the MSSM (RPV columns untouched)

MSSM RPC squarks


$$\tilde{q} \rightarrow q \text{ mET}$$



BRs in the MSSM may undergo a reduction due to squarks limits at 2+ TeV

MSSM-RPV squarks

$$\tilde{q} \rightarrow qq$$



RPV MSSM seems to have less stringent bounds, so it might re-enable the MSSM (RPV columns untouched)


Randall-Sundrum \rightarrow Heavy Vector Triplet scenario

$$\rho \rightarrow VV$$

hep-ph/0709.0007

	A_1		\tilde{Z}_1		\tilde{Z}_{X1}	
	$\Gamma(\text{GeV})$	BR	$\Gamma(\text{GeV})$	BR	$\Gamma(\text{GeV})$	BR
$\bar{t}t$	55.8	0.54	18.3	0.16	55.6	0.41
$\bar{b}b$	0.9	8.7×10^{-3}	0.12	10^{-3}	28.5	0.21
$\bar{u}u$	0.28	2.7×10^{-3}	0.2	1.7×10^{-3}	0.05	4×10^{-4}
$\bar{d}d$	0.07	6.7×10^{-4}	0.25	2.2×10^{-3}	0.07	5.2×10^{-4}
$\ell^+\ell^-$	0.21	2×10^{-3}	0.06	5×10^{-4}	0.02	1.2×10^{-4}
$W_L^+W_L^-$	45.5	0.44	0.88	7.7×10^{-3}	50.2	0.37
$Z_L h$	-	-	94	0.82	2.7	0.02
Total	103.3		114.6		135.6	

- RS FCNC computed for **3 TeV** new resonances in 2013
- Quite generous/safe assumption back then
- No specific RS searches found so far for the relevant color-singlet new resonances for the FCNC BR
- Closest searches expressed as “Heavy Vector Triplet” model B (established map between models and appropriate rescaling in progress)
- Limits potentially touching the 3+ TeV at 140/fb and may reach 6 TeV at HL-LHC (prelim. estimate)





Randall-Sundrum \rightarrow beyond Heavy Vector Triplet scenario

$$\rho \rightarrow t\bar{t}$$

	A_1		\tilde{Z}_1		\tilde{Z}_{X1}	
	$\Gamma(\text{GeV})$	BR	$\Gamma(\text{GeV})$	BR	$\Gamma(\text{GeV})$	BR
$t\bar{t}$	55.8	0.54	18.3	0.16	55.6	0.41
$b\bar{b}$	0.9	8.7×10^{-3}	0.12	10^{-3}	28.5	0.21
$u\bar{u}$	0.28	2.7×10^{-3}	0.2	1.7×10^{-3}	0.05	4×10^{-4}
$d\bar{d}$	0.07	6.7×10^{-4}	0.25	2.2×10^{-3}	0.07	5.2×10^{-4}
$\ell^+\ell^-$	0.21	2×10^{-3}	0.06	5×10^{-4}	0.02	1.2×10^{-4}
$W_L^+W_L^-$	45.5	0.44	0.88	7.7×10^{-3}	50.2	0.37
$Z_L h$	-	-	94	0.82	2.7	0.02
Total	103.3		114.6		135.6	

- RS FCNC computed for **3 TeV** new resonances in 2013
- Quite generous/safe assumption back then
- No specific RS searches found so far for the relevant color-singlet new resonances for the FCNC BR
- Closest searches expressed as Z'_{TC2} or dark matter mediator Z' (established map between models and appropriate rescaling in progress)
- Limits potentially touching the 3+ TeV at 140/fb and may reach 6 TeV at HL-LHC (prelim. estimate)




appropriate rescaling of couplings necessary, in progress

Top quark decay at the Top Factory

$t \rightarrow BSM$

- can we find a (light) state in the mass range not currently investigated by the LHC?
- can we find a new state in the final states not currently investigated by the LHC?





Top quark decay at the Top Factory

$$t \rightarrow \phi q, \phi \rightarrow bb$$




ATLAS 2301.03902, 1806.02836 Banerjee, Chala, Spannowsky,

One extra singlet: $SO(6) \rightarrow SO(5)$

2. Rare top decays. About $O(10^6)$ ttbar events at 350 GeV FCC-ee, ILC, ...



Banerjee, MC, Spannowsky '18 23



Ample room for improvement with top factory

New terrain of search for m_X outside the range studied by LHC (e.g. $m_X \in [10, 20] \text{ GeV}$)


Top quark decay at the Top Factory

$$t \rightarrow \phi q, \phi \rightarrow b\bar{b}$$

Kevin Mota and Kirill Skovpen started to work on IDEA projections for this decay

- Delphes simulation using **IDEA** detector card
- First look at semi-leptonic final states
- So far simple analysis pipeline already shows potential for signal-vs-background discrimination
- N=4 Durham exclusive jet clustering so far, other possibilities to be investigated
- Still can exploit jet flavor tagging, angular variables, ...


$$\frac{1}{\sigma} \frac{d\sigma}{m_{jj}}$$



NP in top quark production

$$e^+e^- \rightarrow tt\phi$$

- can we find a light new state produced in association with $t\bar{t}$ during a **threshold scan** at $\sqrt{s} = 350$ GeV?
- can we find a a light new state in the “**above**” $t\bar{t}$ **threshold** operation of the top quark factory $\sqrt{s} = 365$ or 380 GeV ?



threshold studies with BSM

invisible