Contribution ID: 38

Light by Light Scattering at NLO in QCD+QED

Tuesday, 10 September 2024 11:25 (25 minutes)

In this talk I plan to discuss the QCD+QED corrections to Light by Light scattering (LbL) process at next-toleading order (NLO). The recent experimental observation of this fundamental process at the Large Hadron Collider has revived the interest to precisely predict its cross-section. We discuss two radically different computational approaches, both exact in the fermion mass dependence, thus offering a strong cross-check of our results. The first approach is a fully analytic method to calculate compact and well-organized twoloop helicity amplitudes. The second one is entirely numerical and leverages the Local Unitarity construction. Our two calculations agree with each other and conclude that including the exact fermion mass contribution typically increases the size of the NLO corrections. Moreover, we find that the exact result converges slowly to the massless limit of the high-energy regime, thus emphasizing the importance of including the full mass dependence at NLO. We also compare our results with the ATLAS measurement of LbL in ultra-peripheral lead-lead collisions, and find that the inclusion of exact NLO corrections reduces, but does not eliminate, the existing tension with theoretical predictions.

Primary authors: Dr ABDUL HAMEED, Ajjath (LPTHE (CNRS), Paris); CHAUBEY, Ekta (Istituto Nazionale di Fisica Nucleare); SHAO, Hua-Sheng (LPTHE Paris); FRAAIJEC, Mathijs (Institute for Theoretical Physics, University of Bern); Dr HIRSCHI, Valentin (Institute for Theoretical Physics, University of Bern)

Presenter: Dr ABDUL HAMEED, Ajjath (LPTHE (CNRS), Paris)

Session Classification: Electroweak and Higgs Physics, EFT and BSM

Track Classification: Electroweak and Higgs Physics, EFT and BSM