

Overview of the Baikal-GVD neutrino telescope: status 2024



UNIVERZITA
KOMENSKÉHO
V BRATISLAVE

Rastislav Dvornický
for the Baikal-GVD collaboration

Comenius University in Bratislava, Slovakia
dvornicky@fmph.uniba.sk



BAIKAL-GVD



Platform “Ivanovskaya” of Circum-Baikal railway

Telescope is located 3.6 km away from shore



Constant lake depth: 1366 - 1367 m

Location

Water transparency:

- Absorption length: 21 - 23 m
- Scattering length: 60 - 80 m


Stable ice cover over 7 - 8 weeks in February - April: detector deployment and maintenance





Basic components

String:



Each string carries 36 optical modules (OMs)

- 10-inch high Q eff. PMT
- 15 m vertical spacing
- OM facing the lake bottom

Time calibration systems

- LED in each OM
- LED beacons
- Isotropic lasers between clusters
- Calibration precision ~ 2 ns


Geometry calibration system

- Acoustic modems on each string
- OM positioning precision ~ 20 cm

- buoy
- string master module
- section master module

- optical module
- acoustic modem
- anchor

Optical module (OM):





Detector status

Presently detector consists of 110 strings arranged into 14 independent detectors - **clusters**


- 3960 OMs in total

Baikal-GVD cluster:

- 8 regular strings, 525 m is instrumented with optical modules (OM)
- 60m radius
- Inter-cluster string carrying lasers, some instrumented with OMs
- Has its own control, trigger and readout systems

Additional cluster “EXP”:

- 4 strings with experimental high-speed DAQ






Expedition 2024

Successful 2024 deployment campaign 16/02 - 07/04


- 14 regular strings carrying 36 OMs installed
- 2 strings added to experimental (“optical”) cluster
- Pilot string for HUNT project



HUNT - next generation neutrino telescope project [\[PoS\(ICRC2023\)1080\]](#)

OMs based on
20-inch PMT

Pilot string with 12 OMs
deployed as a part of
experimental cluster in joint
IHEP (Beijing) and Baikal-GVD
effort






Data flow

Each cluster is connected to the **shore center** with opto-electric cable

- Power distribution
- Data transmission



Baikal shore center:

- Power distribution
- Data readout hardware/software
- Data-taking management (shifter)
- Data quality control
- Long-term storage of raw data
- Alert system (to be deployed)




Data flow



Raw data are transferred from the Shore center to JINR

- Shore center → Baikalsk: 300 Mbit/s radiochannel
- Baikalsk → JINR: Ethernet
- Compressed data volume ~10-40 GB per day per cluster
- Full-scale reconstruction at JINR
- **Delay due to shore → JINR data transfer: < 1 min**



JINR computing farm:

- Long-term storage of raw data
- Event reconstruction, storage
- Databases
- Alert workflow
- User analysis



Event reconstruction

Cluster event is read-out if coincident signal is found on neighbouring OM
An event frame is 5 μ s

Most of pulses (or hits) in the event frame are noise from lake water luminescence:



- Typical pulse rate 20-100 kHz
- ~ 1 photoelectron (p.e.) charge deposition
- Substantial seasonal variations
- Rate is larger on top layers

Challenge for our MC simulation

Variety of algorithms for noise suppression

Machine learning -based algorithm in development:
[\[arXiv:2210.04653\]](https://arxiv.org/abs/2210.04653)

track-like event before the noise cleaning, data 2019






Event reconstruction




Time, location and deposited charge of each pulse are used for the reconstruction

Track angular resolution:
 $\sim 0.8^\circ$ - $\sim 0.2^\circ$ for tracks longer than 200 m

Cascade angular resolution:
 $2-4^\circ$ depending on energy and cascade location



track-like,
data 2019




cascade-like,
data 2022




Event reconstruction

Track energy resolution:
Factor 3 for 100 TeV muon



Cascade energy resolution:
 $\delta E/E \sim 10\text{-}30\%$





Results in cascade channel




Search for diffuse astrophysical neutrino flux

Most of the Baikal-GVD data were processed with HE cascade analysis algorithms

Four years dataset: 04.2018 - 03.2022

14328 events $E_{sh} > 10$ TeV, $N_{hit} > 11$ after quality cuts



All-sky analysis:

- $E_{sh} > 70$ TeV, $N_{hit} > 19$
- 16 events were selected
- 8.2 background ev. expected
 - $7.4 \mu_{atm}$, $0.8 \nu_{atm}$
- $5.8 \nu_{astro}$ ev. expected
- Largest energy event: ~ 1.2 PeV

All-sky diffuse flux significance: 2.22σ

[Phys.Rev. D 107, 042005 (2023)]



Search for diffuse astrophysical neutrino flux


Analysis of upward-going events

- Zenith angle cut: $\cos(\theta) < -0.25$
- **Loosened cuts: $E_{sh} > 15$ TeV, $N_{hit} > 11$**
- 11 events selected
- 3.2 ± 1.0 atm. background ev. are expected
 - $0.5 \mu_{atm}$, $2.7 \nu_{atm}$
- Highest energy: 224 TeV


Significance of diffuse flux in upward-going events: 3.05σ !

Main uncertainties

- Absorption length $\pm 5\%$
- OM sensitivity $\pm 10\%$
- ν_{atm} flux normalisation $\pm 15\%$



[Phys.Rev. D 107, 042005 (2023)]






Diffuse spectrum

Extraction of spectrum power and flux normalisation: $\Phi_{astro}^{\nu+\bar{\nu}} = 3 \times 10^{-18} \phi_{astro} \left(\frac{E_\nu}{E_0} \right)^{-\gamma_{astro}}$

$$\gamma_{astro} = 2.58$$

$$\phi_{astro} = 3.04$$



Results are in agreement with previous measurements by IceCube and ANTARES


First “non-IceCube” evidence for diffuse ν_{astro} flux at above 3σ !

[Phys.Rev. D 107, 042005 (2023)]



HE cascade sky map

[MNRAS 526 (2023) 942]




Three events close to the Galactic plane (grey line)

The red plus and circle – IC hotspot
[Aartsen & et al. ApJ, 835,151 (2017)]

Intriguing coincidence in view of recent IC statement on diffuse flux from galactic plane [Science 380, 6652, 1338-1343 (2023)]

Best fit positions and 90% angular uncertainty regions



color represents energy:

$E_{rec} < 100 \text{ TeV}$
 $100 \text{ TeV} < E_{rec} < 200 \text{ TeV}$
 $200 < E_{reco} < 1000 \text{ TeV}$
 $E_{rec} > 1 \text{ PeV}$



Cascade diffuse flux update

Preliminary: An update of analysis adding data from 04.2022 - 03.2023 (10 cluster detector)

Comparison of statistical significances for old and new samples

All-sky analysis

Seasons	N_{data}	N_{bckg}	P-value	$\sigma(\text{stat.})$
18-21	16	8.2	2.09×10^{-2}	2.31
18-22	28	14.5	1.06×10^{-3}	3.07

Upgoing analysis

Seasons	N_{data}	N_{bckg}	P-value	$\sigma(\text{stat.})$
18-21	11	3.2	1.7×10^{-3}	3.13
18-22	19	5.7	1.11×10^{-5}	4.24


Significance of excess over atmospheric background increases



HE cascades and the galaxy plane


Hint on alert events concentration near galactic plane

Baikal-GVD: 25 all sky alerts for **04/2018-03/2022**



Baikal-GVD alerts compared to IC galaxy plane analysis

Extended dataset of
45 all-sky alerts **04/2018 - 03/2023**




Analysis continues




Cascades: TXS0506 coincidence

[MNRAS 527 (2024) 8784]



Upgoing cascade analysis, highest energy event (18.04.2021):

- 224 TeV, 24 hits
- Neutrino source candidate TXS 0506+056 is within 90% containment circle
- Signalness: 97.1% (probability of astro origin)
- Chance coincidence probability ($E > 200$ TeV): 0.0074



Analysis of RATAN-600 radiotelescope data (11GHz) showed increased activity

- IC event registered during γ flare
- Baikal event during radio flare
- Consistency with IC observations: 8% or 13% depending on ν spectrum assumption



Track-like channel




Track-like events


Two modes of analysis

- Single-cluster: each cluster is treated as an independent detector
- Multi-cluster: common reconstruction for simultaneously triggered single-cluster events


Single-cluster
upgoing
event:



Multi-cluster events:



Lake and PMT noise hits




late

early




Track trigger-level sensitivity, 12 clusters

Effective area



Effective volume:
measure of sensitive volume



Absorption in Earth is not taken into account

At the reconstruction level sensitivity will be lower (estimation is in progress)



First track-like neutrino candidate event sample



First set of single-cluster muon neutrino candidates based on 2019 data

- Cut-based analysis optimized for low-energy (atmospheric) neutrino, $\langle E_\nu \rangle \sim 500$ GeV
- Runs from April 1st until June 30th 2019
- Results are compared to atmospheric neutrino simulation

MC expected: 43.6

- atm. neutrino :43.6
- atm. muon: 0

Observed: 44



Excellent agreement of MC expectation and data

[Eur. Phys. J. C 81, 1025 (2021)]

Successful Baikal-GVD performance validation



Progress in single-cluster track-like analysis

Large-scale data and MC track channel reprocessing campaign is ongoing

Improved track MC with more detailed detector description

- Switch to CORSIKA 7.741 for muon bundle simulation
- Realistic time-dependent detector configuration

Improved muon reconstruction


- New noise suppression algorithm
- More precise track fit algorithm
- Improved neutrino selection capabilities

Improvement in tools for muon background suppression


BDT discriminant as a main variable for neutrino selection

Good data-MC agreement
→ background is under control

Low-E BDT



High-E BDT





Increasing ν_μ candidate dataset

Seasons 2020-2021 were reprocessed in single-cluster regime


- 3845 days single-cluster livetime equivalent
- Validation of reconstruction results is ongoing
- Optimisation of high-energy ν selection is ongoing

Demonstration sample of ν_μ candidates dominated by atmospheric neutrino

671 neutrino candidates selected in 3845 days

- atm. μ : 3.5
- atm. ν : 565.1
- **data: 671**

Total rate is 15% larger than MC expectation





High-energy track event candidate

Preliminary: spectacular single-cluster event
with high probability of astrophysical origin

Season 2019, Cluster 3, run 590

θ_z = 153.4°

N_{hits} = 30


E_{rec} = **103.4 TeV**

[68% CI: 24.9 < E < 266.3 TeV]

Track length: 332.4 m


Angular resolution: 0.45° (50%)

0.67° (68%)





Track-like multi-cluster analysis



Track-like multi-cluster analysis
unlocks the full Baikal-GVD potential
in angular resolution

First multi-cluster neutrino candidate
events start to appear

Example of ν candidate event:

Summer 2019
Clusters 1 & 4

$$\theta_z = 125.6^\circ$$

$$N_{\text{hits}} = 10$$

$$\text{track length} = 399 \text{ m}$$


$$E_{\text{rec}} < 1 \text{ TeV}$$



Track-like event multi-cluster analysis


In total 5 ν candidates selected from 150 days of 2019
(5-cluster detector)
Dominated by atmospheric neutrino

Multi-cluster analysis is in the development phase





Track-like events skymap



Multi-cluster neutrino candidate events, very preliminary, **dominated by atmospheric events**

Single cluster 100 TeV event - high probability of astrophysical origin



Alert program



Alert workflow


Getting ready to full-scale participation in real-time multi-messenger alert exchange

Automated alert generation and follow-up system

- Baikal-GVD alerts: distribution of our own alerts for events with high probability of astro origin
- Follow-up: follow-up analysis of external alert events


Baikal-GVD alert generation

- Simplified extrapolated calibrations
- Processing delay 3-10 minutes
- Planned to be deployed at the shore to reduce delay
- Presently internal distribution of alerts





Global Coordinate Network (GCN) alert follow-up



Fermi-GBM/LAT:

[$T_0 - 1 \text{ day}, T_0$],
 [$T_0 - 1 \text{ day}, T_0 + 12 \text{ hours}$],
 [$T_0 - 1 \text{ day}, T_0 + 1 \text{ day}$]

LIGO-Virgo-KAGRA:


IGWN reception: “significant” = 1
 [$T_0 - 1000 \text{ s}, T_0 + 1000 \text{ s}$],
 [$T_0 - 1000\text{s}, T_0 + 14 \text{ days}$]

IceCube:

[$T_0 - 1 \text{ h}, T_0 + 1 \text{ h}$]
 [$T_0 - 1 \text{ day}, T_0 + 1 \text{ day}$]

Search for online coincidences:


- ON/OFF method
- ON includes 90% localization error and Baikal-GVD median angular resolution
- OFF is extended within a ± 5 declination band
- OFF is evaluated using real data from previous seasons






Follow-up of IceCube and ANTARES alerts

60 ANTARES alerts followed,
3 correlated cascades
[PoS(ICRC2021)1121]



[PoS(ECRS2022)096]

Baikal-GVD neutrino alerts and follow-up for neutrinos



Follow-up of IceCube “astrotracks” events (~20 per year)

- On 8.12.2021 detected cascade from the direction of blazar PKS0735+17 in coincidence with IC211208A
- Delay wrt. IC: 3.95 hrs., $E \sim 43$ TeV
- Pre-trial significance: 2.85σ , later reduced to 1.13σ
- Astrotelegram published:

<https://www.astronomerstelegam.org/?read=15112>



Multi-messenger follow-ups


Baikal-GVD follows reported multimessenger high-energy events, e.g.:

GW170817 (LIGO/VIRGO) - neutron star merger, first gravitational waves detection associated with γ /optical/radio signal: time-integrated flux (fluence) limit is set

[Phys. Rev. Lett. 119, 161101]
[JETP Letters, v.108, issue 12]

Radio-burst from magnetar **SGR 1935+2154** (28.04.20)

- IceCube fluence limit: $5.2 \cdot 10^{-2} \text{ GeV} \cdot \text{cm}^{-2}$
- ANTARES fluence limit: $14 \text{ GeV} \cdot \text{cm}^{-2}$
- Baikal-GVD fluence limit: $2 \text{ GeV} \cdot \text{cm}^{-2}$ [PoS(ICRC2021)946]





Summary

Baikal-GVD has reached $\sim 0.6 \text{ km}^3$ detector volume:
110 strings carrying 3960 OMs

- Also: 4 strings with experimental high-bandwidth DAQ

Baikal-GVD is joining the astrophysical neutrino origin quest

- Telescope performance was validated with the atmospheric neutrino flux observation
- First high-energy events are selected in track-like event analysis
- HE cascade event analysis confirms the diffuse flux observation at the level above 3σ
- Experiment participates in high-energy alert follow-up and alert exchange



Thank you for your attention!