

RICAP - Roma International Conference for AstroParticle Physics


Rome 23rd-27th September 2024

Gamma-ray emission from Embedded Star Clusters

Giada Peron — INAF Osservatorio Astrofisico di Arcetri

Cosmic rays

The role of Star Clusters



* SCs have strong winds able to accelerate particles at their wind termination shock; $\mathcal{P}_w \approx 0.1 \mathcal{P}_{SN}$

* The acceleration lasts for ~ 1 Myr, favoring the acceleration to the highest energies;

* The ambient is naturally rich of ^{22}Ne ;

$$\epsilon_{winds} X_{winds} + (1 - \epsilon_{winds}) X_{ISM} = X_{CR} \rightarrow \epsilon_{winds} \simeq 16\%$$


* Several detections of regions towards SCs in gamma rays up to the highest energies;

e.g. Bykov et al. 2020, Morlino et al. 2021, Vieu et al. 2022, Gabici et al. 2023

Optical


Star clusters

In the gamma-ray sky are only a handful




Anderson+2013


Infrared



Fermi-4FGL



HESS-HGPS



Gamma




Star clusters

Matching with gamma-ray sources

The match is defined on a geometrical basis:
 match if their distance $< R^*$

R_* ()

- R of the HII region
- R of the SCs (50% of stars)
- R of the termination shock
- R of the bubble







Star clusters


Matching significance

- i. Calculate the number of matches with SC catalogs;
- ii. Calculate the number and distribution of matches with 1000 randomly generated catalogs;
- iii. Evaluate matching significance as:


$$\Sigma = \frac{N_{match}^{real} - \langle N_{match}^{sim} \rangle}{\sigma(N_{match}^{sim})}$$



Star clusters Matching with gamma-ray sources





- * Very good correspondence between **WISE** and **Fermi-LAT** sources
- * The significance decreases for larger objects like Gaia's and TeV sources.
- * Detection extended sources is challenged both by faintness and by source confusion





Star clusters from WISE

Matching with gamma-ray sources: **Fermi-LAT (50 MeV – 1 TeV)**



Matching significance and number of matching sources **increase** with **luminosity** and **radius**

Star clusters seen by Fermi-LAT





Map: Fermi-LAT TS (>1 GeV)

Regions: Fermi-LAT 4FGL sources

Region: Gaia RTS

Region : WISE (22 micron) HII regions

Contours: WISE (22 micron) emission



The morphology of the **Fermi** emission well matches with the **HII region** contours

..and more examples

Peron et al. 2024, in prep



Star clusters seen by Fermi-LAT

In the Vela Molecular cloud Ridge

Star forming complex at ~1-2 kpc

* SC of Age ~ 1 Myr → No Supernovae

* Embedded in dense medium ~ 1000 cm⁻³



Star clusters




Acceleration efficiency

$$\eta = \frac{L_{CR}}{L_w}$$

L_{CR} from full confinement $L_{CR}[measured] \ll L_{CR}[real]$

A fraction, ξ , of L_w goes to blow the bubble of radius R

Maurin et al. 2016



Canto' et al. 2000 [Simulations]

$$\dot{M} \sim 10^{-4} M_{\odot} \text{yr}^{-1}$$

$$v \sim 1000 \text{ km s}^{-1}$$

$$R \sim R_{HII} \sim R_{\gamma}$$

$$R \simeq R_{HII} = \left(\frac{\xi L_w}{n} \right)^{\frac{1}{5}} t^{\frac{3}{5}}$$

Weaver et al. 1977

Star clusters

Acceleration efficiency

$$\eta = \frac{L_{CR}}{L_w}$$

$$L_{CR}[measured] \ll L_{CR}[real]$$

$$\langle \eta \rangle \sim 0.5\% \ll \eta[real]$$



Source	Age [Myr]	N [1E3 cm-3]	Efficiency [%]
RCW 32	2	1.9	0.85
RCW 36	1.1	2.6	0.79
RCW 38	0.5	2.1	4E-03

$$\epsilon_w = \frac{\mathcal{P}_{CR}^w}{\mathcal{P}_{CR}^{tot}} = \frac{0.0005 \times 10^{41} \text{erg s}^{-1}}{7 \times 10^{40} \text{erg s}^{-1}} \simeq 1\% \ll \epsilon_w[real]$$

From isotopic ratio:

$$\epsilon_{winds} X_{winds} + (1 - \epsilon_{winds}) X_{ISM} = X_{CR} \rightarrow \epsilon_{winds} \simeq 16\%$$

Star clusters Prospect for future observations




Celli & Peron, A&A, 2024, 689, A258

Star clusters



Prospect for future observations

Visibility of superbubbles challenged by their extension



Sensitivity curves from Celli & Peron, 2024

HII regions are much more compact



Conclusions

- * We detected **GeV gamma-ray** emission in correspondence of a few **embedded star clusters**;
- * We showed that the coincidence is **statistically significant** and is interpreted as a consequence of the enhanced target around these regions and their compactness; Conversely, no significant emission with Gaia SCs.
- * Embedded clusters are **young** therefore they are not influenced by SN explosions;
- * The derived efficiency suggests that a small **part of galactic cosmic rays is contributed by stellar winds**, consistently with the estimate based on composition;
- * The real **efficiency is larger** as it should account also for escaped particles;
- * More observations will come to help constraining these values and to shed light on the TeV energy band.