



Contribution ID: 299

Type: **oral**

Non-thermal lobe of the Milky Way powered by the Galactic Center outflows

Wednesday, 25 September 2024 14:51 (17 minutes)

The large-scale structures such as Fermi Bubbles and eROSITA Bubbles provide a unique opportunity to study our Milky Way. However, the nature and origin of these large structures are still under debate. In this talk, I will present the identification of several kpc-scale magnetised structures based on their polarized radio emission and their gamma-ray counterparts, which can be interpreted as the radiation of relativistic electrons in the Galactic magnetic halo. These non-thermal structures extend far above and below the Galactic plane and are spatially coincident with the thermal X-ray emission from the eROSITA Bubbles. The morphological consistency of these structures suggests a common origin, which can be sustained by Galactic outflows driven by the active star-forming regions located at 3 – 5 kpc from the Galactic Centre. These results reveal how X-ray-emitting and magnetised halos of spiral galaxies can be related to intense star formation activities and suggest that the X-shaped coherent magnetic structures observed in their halos can stem from galaxy outflows.

Primary author: ZHANG, Heshou (INAF - OA Brera)

Co-authors: Dr PONTI, Gabriele (INAF-OAB); Dr CARRETTI, Ettore (INAF-IRA); Prof. LIU, Ruo-Yu (Nanjing University); Prof. MORRIS, Mark (UCLA); Prof. HAVERKORN, Marijke (Radboud University); Dr LOCATELLI, Nicola (INAF-OAB); Ms ZHENG, Xueying (MPE); Prof. AHARONIAN, Felix (MPIK); Dr ZHANG, Hai-Ming (Guangxi University); Ms ZHANG, Yi (MPE); Mr STEL, Giovanni (INAF-OAB); Prof. STRONG, Andrew (MPE); Mr YEUNG, Micheal (MPE); Dr MERLONI, Andrea (MPE)

Presenter: ZHANG, Heshou (INAF - OA Brera)

Session Classification: Searches for Galactic astrophysical sources