


We present the results of a laser-driven plasma evolution experiment at the POLARIS laser system at the Institute for Optics and Quantum Electronics at the Friedrich-Schiller-University Jena and the Helmholtz-Institute Jena [1]. The interaction of a high intensity main laser pulse with water microdroplet targets was investigated with off-harmonic optical probe pulses. In contrast to experiments with thin foils, the spherical symmetry of droplets facilitates a direct imaging of the plasma expansion process using shadowgraphy. By changing the temporal delay of the probe laser with respect to the main laser, the expansion process was probed in a temporal window between -4.7 to +258 ps relative to the arrival of the main laser pulse. The strong emission of light from the laser-induced plasma at the fundamental and second harmonic frequency was suppressed with a bandpass filter, a polarization filter and a spatial filter (coronagraph). A detailed analysis of the shadowgraphic images allowed us to estimate the plasma expansion velocity of the front and rear side of the droplets at early times of the interaction.

## Experimental setup and diagnostics



- **POLARIS** at 1030 nm, contrast enhanced with plasma mirror,  $\tau \approx 150$  fs (FWHM),  $I = 4 \times 10^{19}$  W / cm<sup>2</sup>,  $a_0 \approx 5.5$ , linear polarization
- **Water droplets** with  $d \approx 20$   $\mu$ m, commercial nozzle (Micro Jet Components), synchronized with laser
- **Optical probe system** [2]: single pass NOPA,  $\mu$ J-level energy, broad bandwidth between 750-950 nm, spectrally filtered after interaction to 800 nm (40 nm FWHM).
- **Imaging setup**: Mitutoyo NIR 10x objective (d), wedged glass plate (Brewster angle) + spatial filter (coronagraph) to suppress the strong plasma emission and scattered light from the main laser pulse, reference setup (probe 1) for comparison
- **Coronagraph**: circular mask of diameter  $d = 300$   $\mu$ m placed in the intermediate image plane of the droplets

## Shadowgraphy measurement




Fig. 1: Shadowgraphic images of water droplets with coronagraph for different pump-probe delays  $\tau$  (top left). The laser energy  $E$  on the target is also shown (bottom left) [1].

## Plasma Expansion

- $\tau = 0$ : blackening of the neighbouring droplets due to generation of plasma  $\Rightarrow n_e \gg n_c = 1.74 \times 10^{21}$  / cm<sup>3</sup> at 800 nm
- Growing spatial extent of the dark volume (overcritical plasma) for increased delays
- Edge of the dark region corresponds to  $n_e \ll n_c$  [3], still, the expansion of the shadow can be measured
- Delays  $\tau < 20$  ps: front and rear side grow along the laser axis linearly: velocity  $v_{\text{front}} \approx 1.27$   $\mu$ m / ps and  $v_{\text{rear}} \approx 0.77$   $\mu$ m / ps  $\Rightarrow$  comparable to similar experiments with liquid targets [4, 5]
- Estimated **ion sound speed**:  $c_s = 0.25$   $\mu$ m / ps (for  $n_e = 153n_c$ )
- after the interaction: decreasing electron temperature, change of expansion geometry to spherical expansion  $\Rightarrow$  smaller expansion velocity for larger times
- Ponderomotive scaling for hot electrons ( $k_B T_e \approx 2.4$  MeV) leads to an ion sound speed of  $c_s \approx 15$   $\mu$ m / ps (protons)  $\Rightarrow$  disagreement of measured values may indicate a different intensity scaling or that the hot electron plasma model cannot be applied




Fig. 2: Expansion of the front (blue) and rear (orange) of the plasma parallel to the laser propagation axis as a function of the delay  $\tau$  [1].

## Summary

- Implementation of a pump-probe setup in a water-based microdroplet experiment to study the temporal evolution of the plasma using shadowgraphy
- Successful suppression of the plasma emission and main laser light scattering by using an off-harmonic spectral filter, a polarization filter and coronagraph
- Measurement of the evolution of the plasma expansion up to delays of +258 ps,
- Linear expansion in the first 20 ps with an expansion velocity  $v_{\text{front}} = 1.27$   $\mu$ m / ps (plasma front) and different behaviour on the rear side
- Estimated ion sound speed  $c_s = 0.25$   $\mu$ m / ps

## Acknowledgements

This poster presentation has received support from the European Union's Horizon 2020 Research and Innovation programme under Grant Agreement No 101004730.

## References:

- [1] M. Beyer, Development of Laser Sources and Diagnostics for Probing Relativistic Laser-Matter Interaction, Master thesis, 2022
- [2] I. Tamer et al., Few-cycle fs-pumped NOPA with passive ultrabroadband spectral shaping, 2020
- [3] M. Nolte, Characterization of liquid micro-droplets for laser-driven proton acceleration, Master thesis, 2023
- [4] G. Becker et al., Characterization of laser-driven proton acceleration from water microdroplets, 2019
- [5] C. Bernert et al., Off-harmonic optical probing of high intensity laser plasma expansion dynamics in solid density hydrogen jets, Scientific reports, 2022