

The Interaction of Intense Light with Wavelength-Scale Objects

Tuesday, 19 September 2023 17:05 (20 minutes)

Intense laser fields interact very differently with micrometric rough surfaces than with flat objects. The interaction features high laser energy absorption and increased emission of MeV electrons, ions, and of hard x-rays.

I will report on how we revealed the underlying reason for this phenomenon by irradiating isolated, micrometric, translationally-symmetric objects by 20 TW laser pulses. The interaction resulted in the emission of two forward-directed electron jets having a small opening angle, a narrow energy spread in the MeV range. PIC simulations show that electrons that are ionized and pulled into vacuum through the vacuum heating mechanism near the edge of the object, manage to circumvent it because of the combined action of the transverse electric field and their cyclotron motion under the magnetic field of the laser. Then, after they pass the object, the electrons form attosecond duration bunches and interact with the laser field over large distances in vacuum. The diffraction of the laser fields obscured by the target creates confined volumes that trap and accelerate electrons within a narrow range of initial momentum. The preservation of the attosecond duration of the electron bunches over large distances, may be applied to the design of future laser-based light sources.

Primary authors: Prof. AREFIEV, Alexey (UC San Diego); Dr LEVANON, Assaf (Tel-Aviv University); Dr BLACKMAN, David (UC San Diego); POMERANTZ, Ishay; Mr COHEN, Itamar (Tel-Aviv University); Dr PERELMUTTER, Lior (Tel-Aviv University); Ms ELKIND, Michal (Tel-Aviv University); Prof. GLENZER, Siegfried (SLAC National Accelerator Laboratory); Ms MEIR, Talia (Tel-Aviv University); Mr CATABI, Tomer (Tel-Aviv University)

Presenter: POMERANTZ, Ishay

Session Classification: WG1: Plasma-based accelerators and ancillary components

Track Classification: WG1: Plasma-based accelerators and ancillary components