6th European Advanced Accelerator Concepts workshop

ID contributo: 282

Fast laser field reconstruction method based on a Gerchberg-Saxton algorithm with modes decomposition

mercoledì 20 settembre 2023 19:00 (1O 30m)

Numerical simulations of laser-plasma acceleration in the non-linear regime require an accurate modelling of the laser driving the plasma waves. To reconstruct the transverse laser field distribution from fluence measurements, a fast and flexible field reconstruction method based on the Gerchberg-Saxton Algorithm with Hermite-Gauss Modes Decomposition (GSA-MD) has been developped [2].

The minimisation of the reconstruction error via tuning of the Hermite-Gauss modes origins is used to improve the accuracy of the GSA-MD.

This algorithm was used to reconstruct the asymmetries of the energy distribution in the focal volume [1,2,3] from fluence images collected during two experimental campaigns at the APOLLON and Lund Laser Centre facilities.

It will be shown how using the GSA-MD-reconstructed laser field as input for quasi-cylindrical PIC simulations allowed to obtain a better agreement with experimental results compared to using an ideal laser field [1,3]. This led to a better understanding of the effects of laser asymmetries on the quality of the output electrons bunch in the injector stage.

References

[1] I. Moulanier et al., Physics of Plasmas 30, 053109 (2023)

[2] I. Moulanier et al., Submitted paper to JOSA B (2023)

[3] L. T. Dickson et al., Physical Review Accelerators and Beams 25, 101301 (2022)

Autori principali: MOULANIER, Ioaquin (Laboratoire de Physique des Gaz et Plamas); DICKSON, Lewis; MAS-SIMO, Francesco (LPGP - CNRS); CROS, Brigitte (LPGP-CNRS-UP11)

Relatore: MOULANIER, Ioaquin (Laboratoire de Physique des Gaz et Plamas)

Classifica Sessioni: Poster session

Classificazione della track: WG3: Theory and simulations