

G

S

S

I



# THE NUSES MISSION

**P. Savina** (GSSI and INFN-LNGS)  
on behalf of the NUSES Collaboration

# THE NUSES MISSION

Italian led mission conceived as a **pathfinder** for **new observation methods and technologies** in the study of high and low energy radiations from space enabling new sensors and tools

Flagship initiative to relaunch economy of L'Aquila Area

Joint **GSSI**-Thales Alenia Space Italy (**TAS-I**) project.



Funded by the Italian government and the Abruzzo regional government.



Joint **GSSI-INFN** effort currently ongoing for the **design and construction** of the NUSES payloads.



The NUSES mission has been approved by ASI: funds for launch and ground segment.

## Industrial Partners:



# THE NUSES COLLABORATION

60+ persons from many institutions.

Large **expertise** (and **sinergies**) from space missions/R&D:  
AMS, DAMPE, eASTROGAM, Fermi, LIMADOU, GAPS,  
HERD, PAMELA, POEMMA, SPB2, ...

## Italian Institutes:

- Gran Sasso Science Institute
- Laboratori Nazionali del Gran Sasso
- Università dell'Aquila
- Università di Roma "Tor Vergata" and INFN-Roma2
- Università di Torino and INFN Torino
- Università di Trento and INFN-TIFPA
- Università di Bari and INFN Bari
- Università di Padova and INFN Padova
- Università "Federico II" and INFN Napoli
- Università del Salento and INFN Lecce

## Other Institutes:


- University of Geneva
- University of Chicago
- Interests from other US institutions, ...



# NUSES: TWO PAYLOADS


## Zirè

- Measure the flux ( $E < 300$  MeV) of cosmic  $e^-$ ,  $p$  and light nuclei of solar/galactic origin;
- Study of the cosmic radiation variability (Van Allen belt system);
- Possible correlation with seismic activity due to **Magnetosphere-Ionosphere-Lithosphere Coupling (MILC)**;
- Detection of 0.1 - 30 MeV photons for **study of transient and stable gamma sources**;
- Paving the way for future **applications of new technology (SiPM, ...)**;



## Terzina

**Pathfinder** for future missions devoted to UHE cosmic rays and neutrino astronomy through **space-based atmospheric Cherenkov light** detection.



## New Technologies and approaches



Development of new observational techniques, testing new sensors (e.g. **SiPM**) and related electronics/DAQ for space missions. New solutions for the satellite platform.

# NUSES: THE SATELLITE



## New Italian Micro BUS


New platform concept which foresees a modular approach relying on standard trays.





AOCS, Telemetry and Telecommand (TT&C) and GPS Receiver units





AOCS (Altitude and Orbit Control System): units/actuators



EPS (Electric Power system)





# NUSES: THE ORBIT



**Low Earth Orbit (LEO)** with high inclination, sun synchronous orbit on the day-night border.

- ❖ Altitude ~550 km
- ❖ Inclination = 97.8 deg
- ❖ LTAN = 18:00



Measuring CR fluxes with energy  $E < 300$  MeV:


Energy spectrum of low energy charged cosmic particles is different with respect to the Local Interstellar Spectra (the spectra that would be measured outside the heliospheric boundaries)

Magnitude is strictly dependent on the time of the measurements

## Goals:


- Monitoring near-Earth space environment
- Study of fluxes of high- & low-energy charged particles from the Van Allen radiation belts
- Measurements of magnetospheric and solar activity
- Analysis of the ionospheric and plasmaspheric fluctuations & possible correlations with seismo-electromagnetic phenomena

# ZIRÈ: (VAN ALLEN) RADIATION BELTS




Van Allen radiation belt is a zone of **energetic charged particles**.

- Inner belt contains
- electrons [0.01, 1] MeV
  - protons [0.1, 100] MeV



Protons with  $E > 50$  MeV in the lower belts at lower altitudes are mostly the result of the beta decay of neutrons created by cosmic ray collisions with nuclei of the upper atmosphere.

The outer belt consists mainly of high-energy (0.1–10 MeV) electrons trapped by the Earth's magnetosphere. It is more variable than the inner belt, as it is more easily influenced by solar activity.






# ZIRÈ MOTIVATION: MILC and TRANSIENT SOURCES


Acoustic-Gravity Wave (AGW) generated around epicenter.

AGW interacts with the ionosphere generating plasma density variations.

Hunting for possible correlations of the electron and proton fluxes with seismic activities through magnetosphere ionosphere lithosphere coupling (MILC) phenomena




Study of the astrophysical phenomena in the 100 keV up 50 MeV region which is to date not extensively measured.



# ZIRÈ: LAYOUT

## Fiber TracKer (FTK)

3 double layer XY modules of fibers to be used for track identification.



## Plastic Scintillator Tower (PST)


Tower of 32 Plastic Scintillator layers. Each layer is composed by 3 bars.

6 layers:  
4x12x1 cm<sup>3</sup>  
26 layers:  
4x12x0.5 cm<sup>3</sup>





## Anti-Coincidence System (ACS)

9 PS layers surrounding the detector.





## Calorimeter (CALOg)

Matrix 4x4x2 of LYSO/GAGG crystals. Segmented in cubes to be Crystal Eye pathfinder.




# ZIRÈ: e, p ACCEPTANCE AND $\gamma$ EFFECTIVE AREA



CALOG will be also used for the study of **low energy  $\gamma$ -rays** between 10 keV and 50 MeV.


**Two windows** surrounding the CALOG are included for this purpose.

# ZIRÈ: A PATHFINDER FOR CRYSTAL EYE



- Wide FOV: ~ 6 sr.
- Full sky coverage.
- Very large effective area: ~ 5 times Fermi-GBM at 1 MeV.
- High localization capability: few degrees.
- Use LYSO/GAGG scintillator with SiPM for the signal readout.

See poster from R. Sarkar



# ZIRÈ: PARTICLE IDENTIFICATION



**Particle Identification** by studying the correlation between the energy deposit inside FTK+PS0 and the inverse of the total energy deposition in the whole detector.

$$\Delta_0 = \Delta E_{\text{FTK+PS0}} \simeq \frac{dE}{dX} \propto \frac{z^2}{\beta^2} F(\gamma\beta)$$

$$E_{\text{dep}} \simeq E_k \propto \beta^2$$




$$\Delta_0 \propto \frac{z^2 F(\gamma\beta)}{E_{\text{dep}}}$$

For non-relativistic nuclei




# ZIRÈ: LOW ENERGY MODULE


Contained Event: **Selected**





Through-going:  
**MIP - Calibrations**



Low-energy:  
**Single silicon spectra**




Shielded/Vetoed:  
**Rejected**




# ZIRETTINO (TEST AND CALIBRATIONS)


4 LYSO and 4 GAGG crystals





Test Beam Setup



8 layers of PST Bars





Single FTK layer



# TERZINA: Astrophysical neutrinos and CR


The observation of **astrophysical neutrinos** at energies  $>$  **few PeV** can be achieved only by detecting EAS produced by Earth skimming events. The Cherenkov emission of these cascades provides a **unique signal for space based (LEO) instruments**.

Similar signals are produced by high energy **CR** ( $E > 1$  PeV) impinging the atmosphere from **above the Earth's limb**. Thus, also CR with  $E > 1$  PeV can be efficiently observed through EAS's Cherenkov emission from space.








# TERZINA: CHERENKOV SIGNAL



Most contributing atmosphere layers @ 550 km:  
between 20 - 40 km altitude





# TERZINA: LAYOUT




- ✓ Equivalent focal length 925 mm
- ✓ Field of View (FoV) :  $7.2^\circ$
- ✓ Point spread function (PSF) :  $<1.0$  mm
- ✓ Effective area of the telescope :  $0.1$  m<sup>2</sup>
- ✓ M1 paraboloid, M2 hyperbole

Point spread function for different inclination angles




Terzina total weight ~35 kg

# TERZINA: FOCAL PLANE



Upside-down image in the telescope

Camera plane with projection on the Earth (total area **360 x 140 km<sup>2</sup>**)



SiPM arrays: 8 x 8 pixels

5 x 2 arrays (640 pixels)

Pixel: 3 x 3 mm<sup>2</sup>


Pixel FoV: ~0.18°

Array dimension: 25.3 x 25.3 mm<sup>2</sup>


Array Effective Area: 24 x 24 mm<sup>2</sup>

See Poster from: [S. Davarpanah](#)

# TERZINA: APERTURE



Night Glow Background



**More than 20 CR events per year with  $E > 100$  PeV expected**

# CONCLUSIONS


## Science Goals:

- **Measure electrons, protons and nuclei up to hundreds MeV;**
- Study particle flux correlation with seismic activity and space weather phenomena;
- Monitor very low energy (0.5-5 MeV) electron flux;
- **Photon detection from 100 keV up to 30 MeV; (Crystal eye pathfinder)**
- Cross correlations among low-energy-electrons, protons-alpha, photons in coincidence with (high-intensity) GRBs;
- First observation of high energy cosmic ray showers from space through Cherenkov signal.
- Certify HE neutrino detection feasibility through Cherenkov emission in the Earth skimming geometry.

## New Technologies:

- Use of SiPM in space;
- Use a scintillating fiber tracker ( $\sim 300\mu\text{m}$ ) readout by SiPM arrays;
- Optimize a LYSO/GAGG crystal array to act as a (astrophysical)  $\gamma$  detector (0.1-30 MeV);
- Design/use low power electronics (try to go down to  $\sim$ few mW/ch );
- Test / Optimize onboard (Standard and/or Machine Learning) techniques for data reduction;
- Test new approaches for the satellite platform.

See poster from R. Sarkar



G

S

S

I



# Thank you

**P. Savina** (GSSI and INFN-LNGS)  
on behalf of the NUSES Collaboration