



Contribution ID: 17

Type: Oral

## Compton Scattering Monochromatic Tunable X-ray Source based on X-band Multi-bunch linac at the University of Tokyo

*Tuesday, 9 September 2008 10:30 (30 minutes)*

**Primary author:** Dr SAKAMOTO, Fumito (The University of Tokyo)

**Co-authors:** Dr HASHIMOTO, Eiko (The University of Tokyo); Mr NOSE, Hiroyuki (IHI); Dr SAKAE, Hisaharu (IHI); Prof. URAKAWA, Junji (KEK); Mr LEE, KiWoo (The University of Tokyo); Dr YAMAMOTO, Masashi (Accutera); Dr YOSHIDA, Mitsuhiro (KEK); Prof. UESAKA, Mitsuru (The University of Tokyo); Mr KANEKO, Namio (IHI); Dr NAKAMURA, Naoki (Accutera); Prof. FUKUDA, Shigeki (KEK); Mr NATSUI, Takuya (The University of Tokyo); Mr YAMAMOTO, Tomohiko (The University of Tokyo); Prof. HIGO, Toshiyasu (KEK); Mr TANIGUCHI, Yoshihiro (The University of Tokyo)

**Presenter:** Dr SAKAMOTO, Fumito (The University of Tokyo)

**Session Classification:** Morning