

Nonfactorizable charm loop in rare $B_s \rightarrow \gamma l^+ l^-$ decay

Tuesday, 30 July 2024 15:40 (20 minutes)

We present the first theoretical calculation of nonfactorizable charm-quark loop contributions to the $B_s \rightarrow \gamma l^+ l^-$ amplitude. This contribution involves the B -meson three-particle Bethe-Salpeter amplitude, $\langle 0 | \bar{s}(y) G_{\mu\nu}(x) b(0) | \bar{B}_s(p) \rangle$, for which we take into account constraints from analyticity and continuity. We calculate the relevant form factors, $H_{A,V}^{\text{NF}}(k'^2, k^2)$, and provide convenient parametrizations of our results, applicable in the region below hadron resonances, $k'^2 < M_{J/\psi}^2$ and $k^2 < M_\phi^2$. We report that factorizable and nonfactorizable charm contributions to the $B_s \rightarrow \gamma l^+ l^-$ amplitude have opposite signs. To compare the charm and the top contributions, the nonfactorizable charming loop contribution is expressed as a non-universal (i.e., dependent on the reaction) q^2 -dependent correction $\Delta^{\text{NF}} C_7(q^2)$ to the Wilson coefficient C_7 . For the $B_s \rightarrow \gamma l^+ l^-$ amplitude the correction is found to be positive, $\Delta^{\text{NF}} C_7(q^2)/C_7 > 0$. Our numerical results for the form factors $H_i^{\text{NF}}(k'^2, k^2)$ depend sizeably on the precise value of the parameter λ_{B_s} , and for a fixed value of λ_{B_s} , $H_i^{\text{NF}}(k'^2, k^2)$ may be calculated with about 10% accuracy.

Primary author: BELOV, Ilia (Istituto Nazionale di Fisica Nucleare)

Co-authors: BEREZHNOY, Alexander; MELIKHOV, Dmitri (Faculty of Physics, University of Vienna)

Presenter: BELOV, Ilia (Istituto Nazionale di Fisica Nucleare)

Session Classification: Heavy Flavours

Track Classification: Heavy Flavours