

THE THEORY OF PTOLEMY

Angelo Esposito



SAPIENZA
UNIVERSITÀ DI ROMA



Istituto Nazionale di Fisica Nucleare

New York University, Nov. 8th 2023

MANY FIELDS

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!

Particle physics

What's the neutrino mass?
What does this tell us about new physics?

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!

Particle physics

What's the neutrino mass?
What does this tell us about new physics?

Cosmology

Can we observe the $C\nu B$?
What lessons about the very early Universe?

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!

Particle physics

What's the neutrino mass?
What does this tell us about new physics?

Cosmology

Can we observe the $C\nu B$?
What lessons about the very early Universe?

Nuclear physics

Find the best β -emitted
Accurately calculation of nuclear
matrix elements

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!

Particle physics

What's the neutrino mass?
What does this tell us about new physics?

Cosmology

Can we observe the $C\nu B$?
What lessons about the very early Universe?

Nuclear physics


Find the best β -emitted
Accurately calculation of nuclear matrix elements

Condensed matter

Spectral distortions induce by condensed matter d.o.f.
Best substrate for $C\nu B$ capture?

MANY FIELDS

The PTOLEMY project touches many different fields. An extremely fascinating problem!



TWO AVENUES

TWO AVENUES


PTOLEMY will study the [spectrum of the electron](#) emitted by the β -processes of tritium, the lightest unstable nucleus

TWO AVENUES

PTOLEMY will study the **spectrum of the electron** emitted by the β -processes of tritium, the lightest unstable nucleus

Measure the neutrino mass

β -decay part of the spectrum





TWO AVENUES

PTOLEMY will study the **spectrum of the electron** emitted by the β -processes of tritium, the lightest unstable nucleus

Measure the neutrino mass

β -decay part of the spectrum





TWO AVENUES

PTOLEMY will study the **spectrum of the electron** emitted by the β -processes of tritium, the lightest unstable nucleus

Measure the neutrino mass

β -decay part of the spectrum




TWO AVENUES

PTOLEMY will study the **spectrum of the electron** emitted by the β -processes of tritium, the lightest unstable nucleus



Measure the neutrino mass

β -decay part of the spectrum



Observe relic neutrinos

Absorption part of the spectrum




TWO AVENUES

PTOLEMY will study the **spectrum of the electron** emitted by the β -processes of tritium, the lightest unstable nucleus


Measure the neutrino mass

β -decay part of the spectrum



Observe relic neutrinos

Absorption part of the spectrum



CONDENSED MATTER

CONDENSED MATTER

The key to the success of the project is **graphene**:

CONDENSED MATTER

The key to the success of the project is **graphene**:

*we can store **large quantities of tritium** in a small volume and achieve **small energy resolutions***

CONDENSED MATTER

The key to the success of the project is **graphene**:

*we can store **large quantities of tritium** in a small volume and achieve **small energy resolutions***

It also makes the theory side of the problem **more challenging**:

CONDENSED MATTER

The key to the success of the project is **graphene**:

*we can store **large quantities of tritium** in a small volume and achieve **small energy resolutions***

It also makes the theory side of the problem **more challenging**:

*we need to account for a **plethora of condensed matter effects**, making the situation qualitatively different from vacuum*

CONDENSED MATTER

The key to the success of the project is **graphene**:

*we can store **large quantities of tritium** in a small volume and achieve **small energy resolutions***

It also makes the theory side of the problem **more challenging**:

*we need to account for a **plethora of condensed matter effects**, making the situation qualitatively different from vacuum*

The **condensed matter d.o.f.** actively participate to the reactions!

CONDENSED MATTER


The key to the success of the project is **graphene**:

we can store *large quantities of tritium* in a small volume and achieve *small energy resolutions*

It also makes the theory side of the problem **more challenging**:

we need to account for a *plethora of condensed matter effects*, making the situation qualitatively different from vacuum

The **condensed matter d.o.f.** actively participate to the reactions!



CONDENSED MATTER


The key to the success of the project is **graphene**:

we can store *large quantities of tritium* in a small volume and achieve *small energy resolutions*

It also makes the theory side of the problem **more challenging**:

we need to account for a *plethora of condensed matter effects*, making the situation qualitatively different from vacuum

The **condensed matter d.o.f.** actively participate to the reactions!



CONDENSED MATTER


The key to the success of the project is **graphene**:

we can store *large quantities of tritium* in a small volume and achieve *small energy resolutions*

It also makes the theory side of the problem **more challenging**:

we need to account for a *plethora of condensed matter effects*, making the situation qualitatively different from vacuum

The **condensed matter d.o.f.** actively participate to the reactions!



CONDENSED MATTER


The key to the success of the project is **graphene**:

we can store *large quantities of tritium* in a small volume and achieve *small energy resolutions*

It also makes the theory side of the problem **more challenging**:

we need to account for a *plethora of condensed matter effects*, making the situation qualitatively different from vacuum

The **condensed matter d.o.f.** actively participate to the reactions!



CONDENSED MATTER


The key to the success of the project is **graphene**:

we can store *large quantities of tritium* in a small volume and achieve *small energy resolutions*

It also makes the theory side of the problem **more challenging**:

we need to account for a *plethora of condensed matter effects*, making the situation qualitatively different from vacuum

The **condensed matter d.o.f.** actively participate to the reactions!



NEUTRINO MASS

NEUTRINO MASS

The β -decay part of the spectrum depends parametrically on the effective neutrino mass:

$$\frac{dN_\beta}{dK_e} \equiv f(E_e, m_\nu),$$

$$m_\nu^2 = \sum_{i=1}^3 |U_{ei}|^2 m_i^2$$

NEUTRINO MASS

The β -decay part of the spectrum depends parametrically on the effective neutrino mass:

$$\frac{dN_\beta}{dK_e} \equiv f(E_e, m_\nu), \quad m_\nu^2 = \sum_{i=1}^3 |U_{ei}|^2 m_i^2$$

The effect is much more pronounced near the electron's maximum energy, the so-called *end-point*


NEUTRINO MASS

The β -decay part of the spectrum depends parametrically on the effective neutrino mass:

$$\frac{dN_{\beta}}{dK_e} \equiv f(E_e, m_{\nu}),$$

$$m_{\nu}^2 = \sum_{i=1}^3 |U_{ei}|^2 m_i^2$$

The effect is much more pronounced near the electron's maximum energy, the so-called *end-point*




NEUTRINO MASS

The β -decay part of the spectrum depends parametrically on the effective neutrino mass:

$$\frac{dN_{\beta}}{dK_e} \equiv f(E_e, m_{\nu}),$$

$$m_{\nu}^2 = \sum_{i=1}^3 |U_{ei}|^2 m_i^2$$

The effect is much more pronounced near the electron's maximum energy, the so-called *end-point*



NEUTRINO MASS

The β -decay part of the spectrum depends parametrically on the effective neutrino mass:

$$\frac{dN_\beta}{dK_e} \equiv f(E_e, m_\nu),$$


$$m_\nu^2 = \sum_{i=1}^3 |U_{ei}|^2 m_i^2$$

The effect is much more pronounced near the electron's maximum energy, the so-called *end-point*

Measure the end-point region



Extract m_ν



NEUTRINO MASS

NEUTRINO MASS

To correctly extract the neutrino mass we need a **reliable model for the event rate, including all relevant effects**

NEUTRINO MASS


To correctly extract the neutrino mass we need a **reliable model for the event rate, including all relevant effects**

The simplest are those due to the **initial ${}^3\text{H}$ and the final ${}^3\text{He}^+$ states**

NEUTRINO MASS

To correctly extract the neutrino mass we need a **reliable model for the event rate, including all relevant effects**


The simplest are those due to the **initial ${}^3\text{H}$ and the final ${}^3\text{He}^+$ states**



NEUTRINO MASS

To correctly extract the neutrino mass we need a **reliable model for the event rate, including all relevant effects**

The simplest are those due to the **initial ${}^3\text{H}$ and the final ${}^3\text{He}^+$ states**



Tritium-graphene binding potential obtained from *Density Functional Theory (DFT)*

[PTOLEMY – PRD 2022]

NEUTRINO MASS

NEUTRINO MASS

The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

NEUTRINO MASS


The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

$$\frac{d\Gamma_\beta}{d\mathbf{k}} = 2\pi \sum_{f, \mathbf{p}} \left| e \langle \mathbf{k} | \nu \langle \mathbf{p} | {}_{He} \langle f | H_w | i \rangle_T \right|^2 \delta(E_i - E_f)$$

NEUTRINO MASS

The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

$$\frac{d\Gamma_\beta}{d\mathbf{k}} = 2\pi \sum_{f, \mathbf{p}} \left| e \langle \mathbf{k} | \nu \langle \mathbf{p} | {}_{He} \langle f | H_w | i \rangle_T \right|^2 \delta(E_i - E_f)$$





[Andrea Casale's Master's thesis]

NEUTRINO MASS

The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

$$\frac{d\Gamma_\beta}{d\mathbf{k}} = 2\pi \sum_{f, \mathbf{p}} \left| e \langle \mathbf{k} | \nu \langle \mathbf{p} | {}_{He} \langle f | H_w | i \rangle_T \right|^2 \delta(E_i - E_f)$$





[Andrea Casale's Master's thesis]

NEUTRINO MASS

The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

$$\frac{d\Gamma_\beta}{d\mathbf{k}} = 2\pi \sum_{f, \mathbf{p}} \left| e \langle \mathbf{k} | \nu \langle \mathbf{p} | {}_{He} \langle f | H_w | i \rangle_T \right|^2 \delta(E_i - E_f)$$





[Andrea Casale's Master's thesis]

NEUTRINO MASS

The electron spectrum depends on the initial ${}^3\text{H}$ state, and must be summed over all possible final ${}^3\text{He}^+$ states,

$$\frac{d\Gamma_\beta}{d\mathbf{k}} = 2\pi \sum_{f, \mathbf{p}} \left| e \langle \mathbf{k} | \nu \langle \mathbf{p} | {}_{He} \langle f | H_w | i \rangle_T \right|^2 \delta(E_i - E_f)$$



[Andrea Casale's Master's thesis]

NEUTRINO MASS

NEUTRINO MASS

Many degrees of freedom yet to be included:

NEUTRINO MASS

Many degrees of freedom yet to be included:

- **vibrational** degrees of freedom

NEUTRINO MASS

Many degrees of freedom yet to be included:

- **vibrational** degrees of freedom
- **electronic** degrees of freedom

NEUTRINO MASS

Many degrees of freedom yet to be included:

- vibrational degrees of freedom
 - electronic degrees of freedom
- } help from correlation functions methods

NEUTRINO MASS

Many degrees of freedom yet to be included:

- vibrational degrees of freedom
 - electronic degrees of freedom
- } help from correlation functions methods


To pin them down with sufficient accuracy will require a lot of feedback from different corners of the field

NEUTRINO MASS

Many degrees of freedom yet to be included:

- vibrational degrees of freedom
 - electronic degrees of freedom
- } help from correlation functions methods

To pin them down with sufficient accuracy will require a lot of feedback from different corners of the field




CνB

$C\nu B$

To hunt for the $C\nu B$ one must try to observe the tiny peak coming from the absorption process


$C\nu B$

To hunt for the $C\nu B$ one must try to observe the tiny peak coming from the absorption process




$C\nu B$

To hunt for the $C\nu B$ one must try to observe the tiny peak coming from the absorption process



$C\nu B$

To hunt for the $C\nu B$ one must try to observe the tiny peak coming from the absorption process



It crucial to be able to distinguish the peak from the “background” coming from the β -decay part of the spectrum

$C\nu B$

$C \not{=} B$

But good old Heisenberg tells us that:

CvB

But good old Heisenberg tells us that:

localization in **x** \longleftrightarrow delocalization in **p**

CνB

But good old Heisenberg tells us that:

localization in \mathbf{x} \longleftrightarrow delocalization in \mathbf{p}

An uncertainty in the tritium momentum implies an **intrinsic quantum distortion** in the electron energy,

$$\Delta K_e \sim \frac{k_e}{m_T} \frac{1}{\Delta x_T}$$

CvB

But good old Heisenberg tells us that:

localization in \mathbf{x} \longleftrightarrow delocalization in \mathbf{p}

An uncertainty in the tritium momentum implies an **intrinsic quantum distortion** in the electron energy,

$$\Delta K_e \sim \frac{k_e}{m_T} \frac{1}{\Delta x_T}$$

for graphene $\Delta x_T \simeq 0.08 \text{ \AA}$

CνB

But good old Heisenberg tells us that:

localization in \mathbf{x} \longleftrightarrow delocalization in \mathbf{p}

An uncertainty in the tritium momentum implies an **intrinsic quantum distortion** in the electron energy,

$$\Delta K_e \sim \frac{k_e}{m_T} \frac{1}{\Delta x_T} \simeq 1 \text{ eV} \gtrsim 2m_\nu$$

for graphene $\Delta x_T \simeq 0.08 \text{ \AA}$

CνB


But good old Heisenberg tells us that:

localization in \mathbf{x} \longleftrightarrow delocalization in \mathbf{p}

An uncertainty in the tritium momentum implies an **intrinsic quantum distortion** in the electron energy,

$$\Delta K_e \sim \frac{k_e}{m_T} \frac{1}{\Delta x_T} \simeq 1 \text{ eV} \gtrsim 2m_\nu$$

for graphene $\Delta x_T \simeq 0.08 \text{ \AA}$



[PTOLEMY – PRD 2022]

$C\nu B$


$C\nu B$

Flat graphene is not ideal for $C\nu B$ detection. What should we do?

CνB

Flat graphene is not ideal for CνB detection. What should we do?

Localize tritium more
right-most peak gets enhanced




[Andrea Casale's Master's thesis]

CνB

Flat graphene is not ideal for CνB detection. What should we do?

Localize tritium less

left-most peak emerges from β -decay



[Andrea Casale's Master's thesis]

$C\nu B$

CνB

How much should we delocalize the initial tritium?

CνB

How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV}$$

CνB

How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$

CνB

How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$


Can we find a substrate that does this job?

CνB

How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$

Can we find a substrate that does this job?




nanotubes?

CνB


How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$

Can we find a substrate that does this job?



nanotubes?




fullerenes?

CνB


How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$

Can we find a substrate that does this job?



nanotubes?



fullerenes?


...what else?

CνB


How much should we delocalize the initial tritium?

$$\Delta K_e \sim 100 \text{ meV} \implies \Delta x_T \sim 1 \text{ \AA}$$

Can we find a substrate that does this job?



nanotubes?



fullerenes?

...what else?

We need to be creative and get to work!

OUTLOOK

OUTLOOK

There's a lot theory involved in the PTOLEMY project. It's challenging, interesting... and worth doing it!

OUTLOOK

There's a lot theory involved in the PTOLEMY project. It's challenging, interesting... and worth doing it!


It mandates for an [interplay between high and low energy physics](#)

OUTLOOK

There's a lot theory involved in the PTOLEMY project. It's challenging, interesting... and worth doing it!

It mandates for an **interplay between high and low energy physics**

There is the potential to close the window on the **neutrino mass**...
but we need accurate theory models!




OUTLOOK

There's a lot theory involved in the PTOLEMY project. It's challenging, interesting... and worth doing it!

It mandates for an **interplay between high and low energy physics**

There is the potential to close the window on the **neutrino mass**... but we need accurate theory models!




To hunt for the $C\nu B$ we need to study **possible substrates to win against Heisenberg**. Lots of studies to be done!

OUTLOOK

There's a lot theory involved in the PTOLEMY project. It's challenging, interesting... and worth doing it!

It mandates for an **interplay between high and low energy physics**

There is the potential to close the window on the **neutrino mass**... but we need accurate theory models!



To hunt for the $C\nu B$ we need to study **possible substrates to win against Heisenberg**. Lots of studies to be done!

Thanks for your attention!