





First Full Dataset Search for a ν_e Low Energy Excess


For the first time, the full set of 1.11×10^{21} POT is used to search for the low-energy excess (LEE) observed in MiniBooNE. This search tests whether the excess is due to an increase in the number of electron neutrinos. Two models are tested: one that models the excess as a function of the neutrino energy and one that preserves the distribution of the shower kinematics in MiniBooNE. Results using the first 6.86×10^{20} POT of this dataset were published in [1].



MicroBooNE Public Note 1127


All public notes are available at: <https://microboone.fnal.gov/public-notes/>

Single Photon Searches



To search the whole photon phase-space for signs of an excess, an **inclusive single photon** analysis has been developed. The analysis targets any final state consistent with what would be observed as a single photon in MiniBooNE, rather than testing a particular model. A selection using Wire-Cell reconstruction [2] tools and targeted BDTs achieves an efficiency of 7.0% and a purity of 40.2%.

MicroBooNE Public Note 1125



Coherent single photon production is a very rare process, and no search has ever been performed. Signal events appear as low energy photons in the forward direction. We use new tools to help reject low energy hadronic activity.

MicroBooNE Public Note 1131


Previous results showed no excess of $NC \Delta \rightarrow N\gamma$ events with protons. This expanded analysis will have more sensitivity to $NC \Delta \rightarrow N\gamma$ events with no protons, compared to [4].

MicroBooNE Public Note 1126

The LEE could consist of a larger scaling factor for op events ($x_{\Delta,op}$) and a smaller scaling factor for Np events ($x_{\Delta,Np}$). These possibilities can be represented as a 2D phase space, and we expect to exclude a larger region than in our previous result.

Electrons? e^-

Photons? γ




WANTED
Solution to the MiniBooNE Anomaly?!


$e^- e^+$ BSM $e^- e^-$ Pairs?

Dark-Sector $e^+ e^-$ Solutions

Sufficiently overlapping or asymmetric $e^- e^-$ pairs from the decay of BSM particles can also mimic potential MiniBooNE signals.



Simulation of dark sector $e^- e^-$ pairs





In one class of such models neutrinos act as a portal to the **dark sector**. Active neutrinos upscatter via a dark photon (Z') off an Argon nucleus to produce an unstable heavy sterile neutrino (N). This heavy sterile neutrino then decays back to a visible $e^- e^-$ pair which can be detected.


MicroBooNE Public Note 1124

Targeted boosted decision trees can select these dark sector $e^- e^-$ events at high efficiencies of 30 to 40%, depending on model parameter space.

MicroBooNE has two ongoing dark sector $e^- e^-$ searches using both the Wire-Cell [2] and Pandora [3] reconstruction frameworks.



95% CLs sensitivity to a targeted 3+1 light Z' model, with $m_{Z'} = 30$ MeV. Model and MiniBooNE allowed region taken from E. Bertuzzo et al [5].



95% CL Sensitivity to a broad class of heavy and light Z' model, highlighting seven masses of Z' ranging from 30 MeV to 1.25 GeV. Model and MiniBooNE allowed regions taken from [6] and using the DarkNews BSM generator [7].