

# Constraints on ultralight dark matter with the European Pulsar Timing Array

Clemente Smarra

SISSA, Trieste

V Gravi-Gamma-Nu Workshop, Bari, 09/10/2024

*'A pacienza vale cchiù d'a scienza*

# Outline

- ▶ Ultra-Light Dark Matter (ULDM)



- ▶ PTAs constraints on ULDM

- ▶ Gravitationally coupled ULDM: CS+ (EPTA, 2023), Afzal+ (NANOGrav, 2023)
  - ▶ Conformally coupled ULDM: CS+ (2024)
  - ▶ ULDM couplings with SM: Kaplan et al. (2022), Afzal+ (NANOGrav, 2023)


# ULDM: where it stands

The CDM paradigm has some well-known issues, for example (Robles et al. 2018):

- ▶ *cusps/core problem*
- ▶ *Missing satellite problem*
- ▶ *Lower-than-expected central densities*

These problems might be alleviated invoking baryonic physics, e.g. gravitational stirring by SNe. But in dwarf spheroidal galaxies?

- ▶ Need to invoke another mechanism → ULDM




"After the discovery of 'antimatter' and 'dark matter', we have just confirmed the existence of 'doesn't matter', which does not have any influence on the Universe whatsoever."

## ULDM: which one?

We focus on a scalar field with mass  $m \sim 10^{-22} \text{eV}$ , whose de Broglie wavelength acts as a *quantum pressure* that suppresses power on very small scales. In formulae:

$$\frac{\lambda_{\text{dB}}}{2\pi} = \frac{\hbar}{mv} \approx 60 \text{pc} \left( \frac{10^{-22} \text{eV}}{m} \right) \left( \frac{10^{-3} c}{v} \right)$$

# PTAs constraints on ULDM




## PTAs: identikit!

- ▶ PTA experiments observe collections of pulsars and search for "special" signatures in their pulse time of arrivals (TOAs).
- ▶ They observe milli-second pulsars (MSPs), the most precise celestial clocks.
- ▶ Challenge: relate the observed time of arrivals (TOAs) **at the observatory** to the time of emission **at the pulsar**. (Edwards et al. 2006)

Convenient to single out three main contributions:

$$t_e^{psr} = t_a^{obs} - \Delta_{\odot} - \Delta_{IS} - \Delta_B$$



# How to look for a signal in PTAs

The main observable in a PTA experiment is the timing residuals,  $\vec{\delta t}$ , which measure the discrepancy between the observed times of arrival (TOAs) and the ones predicted by the pulsar timing model. In general, each process will affect the timing residuals in a peculiar way. Qualitatively,

$$\vec{\delta t} = \mathbf{M}\vec{\epsilon} + \overrightarrow{W.N.} + \overrightarrow{R.N.} + \text{boh?...} \quad (1)$$

In order to look for a signal in PTAs, we should model how it affects the timing residuals!

## ULDM: Classical Wave

In the following, we will think of ULDM as a free scalar field. Due to the huge occupation number (Khmelnitsky and Rubakov, 2013), the ULDM field can be thought as a collection of *classical waves*

$$\phi(\mathbf{x}, t) = A(\mathbf{x})\cos(mt + \alpha(\mathbf{x}))$$

Skipping a little bit of details (Khmelnitsky and Rubakov, 2013), it turns out that(adapted from Porayko, 2018):



$$\Delta t(t) = \pi \frac{G\rho_{DM}}{2m_\phi^3} \left[ \hat{\phi}_E^2 \sin [2m_\phi t + 2\alpha(x_e)] - \hat{\phi}_P^2 \sin \left[ 2m_\phi \left( t - \frac{d_p}{c} \right) + 2\alpha(x_p) \right] \right]$$

- ▶ *correlated limit*
- ▶ *pulsar-correlated limit*
- ▶ *uncorrelated limit*





# ULDM: Gravitational interaction



EPTA paper VI, CS+ (2023) PRL

# ULDM: Coupling to matter

- ▶ The previous results are based on the *gravitational* interaction of ULDM with matter.
- ▶ However, it might also happen that ULDM features some interaction with the Standard Model particles. Why? Why not? What will we be able to say in this case?

**Qualitative understanding of the main point of this part:** the coupling to matter modifies the moment of inertia of the pulsar. By conservation of angular momentum, this produces a change in the spin frequency of the pulsar.

# Conformally coupled ULDM

We will consider here *universal* interactions of ULDM with the SM. To characterize the coupling we define a field-dependent function  $\mathcal{A}(\phi)$  and assume ULDM couples universally with the SM through a Jordan-Fierz metric  $\tilde{g}_{\mu\nu} = \mathcal{A}^2(\phi) g_{\mu\nu}$ .

$$S = M_{\text{P}}^2 \int d^4x \sqrt{-g} \left[ \frac{R}{2} - g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + m^2 \phi^2 \right] + S_m[\psi_m, \tilde{g}_{\mu\nu}]$$

- ▶ FJBD theory:  $\mathcal{A}(\phi) = e^{\alpha\phi} \sim 1 + \alpha\phi$
- ▶ DEF theory:  $\mathcal{A}(\phi) = e^{\beta\phi^2/2} \sim 1 + \frac{1}{2}\beta\phi^2$

## Conformal ULDM: sensitivity

With the help of the code presented in (A.Kuntz, E.Barausse 2024), we compute the angular momentum sensitivity, defined as:

$$s_I = - \frac{1}{2\alpha(\phi)} \frac{d \ln I}{d\phi} \Big|_{N,J} = \frac{1}{2\alpha(\phi)} \frac{d \ln \Omega_{\text{obs}}}{d\phi} \Big|_{N,J},$$

where  $\alpha(\phi) = \mathcal{A}'(\phi)/\mathcal{A}(\phi)$ ,  $N$  is the pulsar's baryon number and  $J$  is the Einstein-frame angular momentum.

# Conformal ULDM

In order to look for a signal in PTAs, we should model how it affects the timing residuals.

$$\text{FJBD: } \Delta t(t) = \frac{\Psi}{m} s_I \hat{\phi}(\mathbf{x}) \sin(mt + \theta(\mathbf{x})) \quad \Psi = 2\alpha \frac{\sqrt{\rho}}{M_{\text{Pl}} m}$$


$$\text{DEF: } \Delta t(t) = \frac{\Psi}{2m} \beta s_I \hat{\phi}^2(\mathbf{x}) \sin(2mt + \theta(\mathbf{x})) \quad \Psi = \frac{\rho}{M_{\text{Pl}}^2 m^2}$$

- ▶ *correlated limit*
- ▶ *pulsar-correlated limit*
- ▶ *uncorrelated limit*




# Conformal bounds vs ULDM mass

FJBD



Outcompete Cassini bounds in the relevant mass range!

DEF



Improve on spontaneous scalarization bounds!


## Bonus: Other ways to study ULDM

In the same way, one can also study ULDM couplings to the SM (Kaplan+ 2022).

The philosophy is the same:

- ▶ Define the Lagrangian
- ▶ Compute the induced timing residuals

$$\mathcal{L} \supset \frac{\phi}{M_{\text{pl}}} \left( \frac{d_\gamma}{4e^2} F_{\mu\nu} F^{\mu\nu} - \sum_{f=e,\mu} d_f m_f \bar{f} f \right)$$




## A few concluding remarks

- ▶ PTAs are wonderful laboratories to test signatures in signals coming from pulsars;
- ▶ It is possible to constrain ULDM density **below** the predicted abundance;
- ▶ If ULDM is non-minimally coupled to the SM, PTA searches can outcompete previous bounds (e.g. Cassini GR bounds or spontaneous scalarization) by several orders of magnitude in the relevant mass range;
- ▶ It is in general possible to set competitive constraints on ULDM couplings to the SM.



# Final Considerations

“San Nicola proteggici da le  
rizz vacand”



# APPENDIX

# ULDM: some formulae


This behaviour can be easily seen by solving the relevant set of equations, namely the

*Schrödinger-Poisson* system:

$$i\hbar\frac{\partial\psi}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\psi + mV\psi$$
$$\nabla^2 V = 4\pi G(\rho - \bar{\rho})$$

Solving this numerically, we have a **soliton-like** behaviour at the centre and a **NFW-like** behaviour in the outskirts.

Robles et al. (2018)



# ULDM: Classical Wave

In the following, we will think of ULDM as a free scalar field. Due to the huge occupation number (Khmelnitsky and Rubakov, 2013), the ULDM field can be thought as a collection of *classical waves*

$$\phi(\mathbf{x}, t) = A(\mathbf{x})\cos(mt + \alpha(\mathbf{x}))$$

Energy momentum tensor:

$$T_{\mu\nu} = \partial_\mu\phi\partial_\nu\phi - \frac{1}{2}g_{\mu\nu}((\partial\phi)^2 - m^2\phi^2)$$

from which

$$\rho_{DM} \equiv T_{00} = \frac{1}{2}m^2A^2$$

## ULDM: Gravitational interaction

To find the gravitational field produced by ULDM, we can write (Newtonian gauge)

$$ds^2 = (1 + 2\Phi(\mathbf{x}, t))dt^2 - (1 - 2\Psi(\mathbf{x}, t))\delta_{ij}dx^i dx^j$$

We can split the potentials in  $t$ -independent and  $t$ -dependent part

$$\Psi(\mathbf{x}, t) \simeq \Psi_0(\mathbf{x}) + \Psi_c(\mathbf{x}) \cos(\omega t + 2\alpha(\mathbf{x})) + \Psi_s(\mathbf{x}) \sin(\omega t + 2\alpha(\mathbf{x}))$$

From the trace of the  $ij$  components of Einstein equations

$$-6\ddot{\Psi} + 2\Delta(\Psi - \Phi) = 8\pi G T_{kk}$$

we get:

- ▶  $\Psi_0 = \Phi_0$ ;
- ▶  $\Psi_c = \frac{1}{2}\pi G A(\mathbf{x})^2 = \pi \frac{G \rho_{DM}(\mathbf{x})}{m_\phi^2}$
- ▶  $\Psi_s = 0$

# ULDM: Gravitational delay

Now, remember what we wrote before:

"In order to look for a signal in PTAs, we should model how it affects the timing residuals!"

$$\Delta t(t) = - \int_0^t \frac{\Omega(t') - \Omega_0}{\Omega_0} dt'$$

Skipping a little bit of details (Khmelnitsky and Rubakov, 2013), it turns out that (adapted from Porayko, 2018):

$$\Delta t(t) = \frac{\Psi_c}{2m_\phi} \left[ \hat{\phi}_E^2 \sin [2m_\phi t + 2\alpha(x_e)] - \hat{\phi}_P^2 \sin \left[ 2m_\phi \left( t - \frac{d_p}{c} \right) + 2\alpha(x_p) \right] \right] \quad (2)$$

- ▶ *correlated limit*
- ▶ *pulsar-correlated limit*
- ▶ *uncorrelated limit*

