

# Raggi cosmici: risultati, interpretazioni, problemi aperti

Fiorenza Donato  
Torino University & INFN

IFAE - Firenze 03/04/2024




# The measured Cosmic Ray (CR) spectrum

CR database: D. Maurin+ 2306:08901

C. Evoli at <https://agenda.infn.it/event/21891/>  
See also N. Tomassetti 2301.10255

Gabici, Evoli, Gaggero, Lipari, Mertsch,  
Orlando, Strong, Vittino 1903.11584



Direct  
measures

Air showers





Fig. 1. The individual CR flux for nuclear species up to Oxygen as measured by PAMELA and AMS02. Shadow regions correspond to 1 sigma total errors (systematic and statistical added in quadrature).



CRs at zero-th order, or  
In the old times there were power laws





1. The bulk of the energy of CRs comes from SNR explosions in the galactic disk

The power of  $\sim$  GeV CRs can be computed (Strong+ApJL 2010) from  $\gamma$  rays as  $P_{CR} \sim 10^{41}$  erg/s. It is equivalent to the power of observed SNRs in the Galaxy

2. CRs are accelerated through diffusive shock acceleration in SNRs

SNRs provide the right energy needed for CRs (Baade&Zwicky 1934)

Classical test is through  $\gamma$ -rays observations of SNRs (O'Drury+ A&A1994)

Still some ambiguities on hadron acceleration by SNRs which, could be explained by leptonic emission (i.e. SNR RX J1713.7-3946)

See Bell MNRAS 1978, MNRAS2004, Bell+MNRAS2013; Caprioli+ MNRAS2009; Blasi+ApJ2012 ; Recchia&Gabici MNRAS2018

Probe: detection of the maximum energy at 67.5 MeV in the  $\pi^0$  decay rest frame;  $\gamma$  rays from molecular clouds illuminated by nearby, freshly accelerated protons




### 3. Composition: primary, secondaries, both

Primaries: produced in the sources (SNR and Pulsars): H, He, CNO, Fe;  $e^-$ ,  $e^+$ ; possibly  $e^+$ ,  $p^-$ ,  $d^-$  from Dark Matter annihilation/decay

Secondaries: produced by spallation of primary CRs (p, He, C, O, Fe) on the interstellar medium (ISM): Li, Be, B, sub-Fe, [...], (radioactive) isotopes ;  $e^+$ ,  $p^-$ ,  $d^-$

N. Tomassetti 2301.10255



Solar System abundances, similar to interstellar ones, are deprived of nuclei such as Li, Be, B, sub-Fe, believed to be of secondary origin

All species are, at some extent, both primary and secondary



## 4. CRs are diffusively confined in an extended magnetic halo

CRs must be confined a region much thicker than the Galactic disk. Radioactive isotopes such as  $^{10}\text{Be}$  indicate the existence of a magnetic diffusive halo several kpc thick (L or H)

$$D(R) \sim D_0 \times f(R) \sim D_0 \times R^\delta$$

$$D_0 \sim 3 \times 10^{28} \left( \frac{H}{5 \text{ kpc}} \right) \left( \frac{\Lambda}{10 \text{ g/cm}^2} \right)^{-1} \text{ cm}^2/\text{s} .$$

Radio haloes observed in external galaxies.

A very extended halo,  $> 100 \text{ kpc}$ , has been observed across **M31** (Karwin+ApJ2019).

DM annihilation has been explored (Karwin+2020).

Non-standard propagation of CRs can explain it (Recchia+ApJ2021)



# Propagation equation

$$\frac{\partial \psi_i(\mathbf{x}, p, t)}{\partial t} = q_i(\mathbf{x}, p) + \nabla \cdot (D_{xx} \nabla \psi_i - \mathbf{V} \psi_i) + \frac{\partial}{\partial p} p^2 D_{pp} \frac{\partial}{\partial p} \frac{1}{p^2} \psi_i - \frac{\partial}{\partial p} \left( \frac{dp}{dt} \psi_i - \frac{p}{3} (\nabla \cdot \mathbf{V}) \psi_i \right) - \frac{1}{\tau_{f,i}} \psi_i - \frac{1}{\tau_{r,i}} \psi_i.$$

**Diffusion:**  $D(\mathbf{x}, R)$  a priori

usually assumed isotropic in the Galaxy:  $D(R) = D_0 R^\delta$  ( $R = pc/Ze$ )  
 $D_0$  and  $\delta$  preferably fixed by B/C (Kappl+15; Genolini+15 (K15))

**Sources:** injection from stellar relics (SNRs, PWN)

Spallation from nuclei scattering off the interstellar medium (ISM)

**Energy losses:** Nuclei: ionisation, Coulomb (spallations)

Leptons: Synchrotron on the galactic  $B \sim 3 \mu\text{G}$

Inverse Compton on photon fields (stellar, CMB, UV, IR)


**Geometry** of the Galaxy: cylinder with half-height  $L \sim \text{kpc}$

**Solution of the eq.:** semi-analytic (Maurin+ 2001, Donato+ 2004, Maurin 2018 ...), USINE codes or fully numerical: GALPROP (Strong & Moskalenko 1998), DRAGON (Evoli+ 2008; 2016), PICARD (Kisskammann, 2014, Kisskammann+ 2015)




# Propagation models vs data

Weinrich+ A&A 2020



Di Mauro, FD+ 2023



See also Evoli+ PRD 2020; Schroer+ PRD 2021; Cuoco&Korsmeier PRD 2021, 2022

Data on nuclear species are well described by propagation models with diffusion coefficient power index  $\delta = 0.50 \pm 0.03$ .



Convection or reacceleration models both work.

Interpretation hampered by cross sections



# Hardening of nuclear spectra

PAMELA Coll. Science 2011; AMS Coll Phys Rept 2021; PRL2017; PRL2018



A general hardening is observed at  $\sim 300$  GV

The rigidity dependence of Li, Be and B measured by PAMELA and AMS are nearly identical, and different from the primary He, C and O (and also p).

The spectral index of secondaries hardens  $\sim 0.13$  more than for primaries




# Hardening of nuclear spectra: diffusion

Most credited explanation is a DIFFUSION effect at  $\sim 300$  GV, naturally with a twice power law for secondaries.

(Geholini+ PRL 2017;; Evoli+ PRD2019)

Tomasetti ApJL 2012


$$K(z, \rho) = \begin{cases} k_0 \beta \rho^\delta & \text{for } |z| < \xi L \text{ (inner halo)} \\ k_0 \beta \rho^{\delta+\Delta} & \text{for } |z| > \xi L \text{ (outer halo)} \end{cases}$$



The diffusion coefficient close to the disk is different than in outer diffusive halo

Interpretations still hampered by spallation cross sections

Evoli+ PRL 2018 - Blasi, Serpico, Amato PRL 2012




CRs diffuse on external turbulence (mainly above the break) and on the waves generated by CRs themselves



# P and He spectra: shifts, breaks and bumps

1. p spectrum is distinctly softer ( $\Delta\gamma \sim 0.1$ ) than He at all energies (**shift**): Not understood yet
2. R dependence of He, C, O are very similar; all (also p) **break** at 300 GV:  $\sim$  understood
3. The p and He spectra  $>$  TeV show a bump: suggestions

Dampe Coll




See also CALET Coll, PRL 2022 and @ ICRC2023

**Bump:** probably an effect in acceleration or escape from the sources




# Light isotopes (D, $^3\text{He}$ )

Coste+ A&A 2012



Gomez-Coral+ PRD 2023



D and  $^3\text{He}$  may **not** share identical slopes.  
Is D harder than  $^3\text{He}$ ?


Fractional contribution of Parent nuclei to D and  $^3\text{He}$  are different



# Radioactive Light isotopes

Radioactive isotopes ( $^{10}\text{Be}$ ,  $^{26}\text{Al}$ ) can track the diffusive halo size  
 Important to test origin and propagation of CRs


Maurin et al, A&A 2002



Weinrich et al. A&A 2020

Jacobs, Mertsch, Pahn 2305.10337

AMS Coll. ICRC2023



Need of precise data on light radioactive isotopes ( $^{10}\text{Be}$  mainly)  
 up to 100 GeV/n - and cross sections.






# Cross sections for Galactic CRs

Production cross sections (source of CRs), and to a lesser extent inelastic cross sections (Loss of CRs)

Data driven parameterizations (Silberberg & Tsao), semi-empirical formulae (Webber+), parametric formulae/direct fit to the data (Galprop), MonteCarlo codes (Fluka, Geant, ...)

Genolini, Moskalenko, Maurin, Unger PRC 2018




Now probably the most limiting aspect now for a clear interpretation of precise CR data coming from space



# Cross sections: the most relevant ones

First: Improve Boron production cross sections

Genolini, Moskalenko, Maurin, Unger PRC 2018 and 2307.06798




Dedicated campaigns at ACCELERATORS are needed.  
Some already started or planned measurements.  
(LHCf, LHCb, NA61, Amber/Compass, ...)




# Antiprotons in CRs

AMS-02 antiprotons are consistent with a secondary astrophysical origin

M. Boudaud+ PRD 2020



Feng, Tomassetti, Oliva PRD2016



- Secondary pbar flux is predicted consistent with AMS-02 data
- Transport and cross section uncertainties are comparable
- A tiny dark matter contribution cannot be excluded
- Precise predictions are mandatory

See also Korsmeier, FD, Di Mauro PRD 2018, Reinert&Winkler JCAP2018





# The observed electron spectrum

Dominated by radiative cooling

AMS Coll Phys.Rept. 2021

CALET Coll. PRL 2023



Data on total electron not fully compatible among them  
 A prominent break is observed at  $\sim$  TeV, (see Dampe talk by De Mitri)  
 still too uncertain to fix models. Pulsars can do the job




# Detected $e^+$ and $e^-$ are local

$$\lambda^2(E, E_S) = 4 \int_E^{E_S} dE' \frac{D(E')}{b_{\text{loss}}(E')}$$

Typical propagation length in the Galaxy

Manconi, Di Mauro, FD JCAP 2017



Sources of  $e^+$  &  $e^-$  in the Galaxy

- Inelastic **hadronic collisions** (asymm.)
- **Pulsar** wind nebulae (PWN) (symm.)
- **Supernova** remnants (SNR) (only  $e^-$ )
- Particle **Dark Matter** annihilation ( $e^+, e^-$ )?

$e^-, e^+$  suffer strong radiative cooling and arrive at Earth if produced within few kpc around it.


Local sources very likely leave their imprints in the spectra




# $e^+$ & $e^-$ spectra, a natural explanation

$e^+$  and  $e^-$  AMS-02 spectra fitted with a multi-component model:  
secondary production,  $e^-$  from SNR,  $e^+$  from PWN

Di Mauro, FD, Manconi PRD 2021



Di MAuro, FD, Korsmeier, Manconi, Orusa 2304.01261



The break at 42 GeV in  $e^-$  is explained by interplay between SNR and PWN  
Secondary  $e^+$  depend strongly on  $L$ . Deficit from  $\sim 1$  GeV




# Antideuteron in cosmic rays

FD, Fornengo, Salati PRD2000

See also Baer & Profumo JCAP2008, FD, Fornengo, Maurin PRD2008, Ibarra & Wild JCAP2012, PRD2013, Fornengo, Maccione, Fitting JCAP2013, Serksnyte et al, PRD 2022, Gomez-Coral PRD2018, Kachelriess+ JCAP2020, CPC2023

P. Von Doetinchem et al. Phys. Rep. 2021  
FD, Fornengo, Korsmeier, PRD 2018



AMS-02 antiproton data

Antideuteron predictions for DM model indicated by pbar AMS-02 data

Bands are for coalescence uncertainty

GAPS in 2024


Antideuterons will be a unique window to probe nuclear fusion in secondary events, and to search for Dark Matter annihilation or decay below  $\sim 1 \text{ GeV/n}$



# Perspectives with antihelium

Cirelli+JHEP2014; Carlson+ PRD2014

FD, Fornengo, Korsmeier, PRD 2018



- Good signal-to-bkgd ratios
- Predictions for most DM models much lower than experimental reach
- Nuclear physics brings relevant effects through  $(p_{\text{coal}})^6$


Challenging for present day experiments  
Looking at antimatter is fundamental for exotic physics




# The Galactic plane seen in neutrinos

The ICECUBE Coll., Science 2023

The galactic plane view strongly established in  $\gamma$  rays



Gaggero+ ApJL 2015; De La Torret+ Frontiers AASS 2023



Galactic cosmic rays interact with atoms of the interstellar medium:



Also contribution from unresolved sources could contribute




# The $\gamma$ -ray counterpart of the sky

Fermi-LAT (0.1-300 GeV)


Courtesy of Silvia Manconi, TMEX 2023

[Fermi-LAT 5 years, energy > 1 GeV]



LHAASO Galactic plane  
(10-1000 TeV)

Caot PRL 2023



A prediction of the emission from all diffuse, point and extended sources, at all latitudes, is possible.

Data at very high energy seem to overshoot predictions from local source extrapolations



## Osservazioni finali

Il quadro teorico attuale sui raggi cosmici risponde a un certo numero di **domande fondamentali** all'ordine zero. Le caratteristiche generali (i.e. leggi di potenza) sono giustificate.

Nuovi dati, e molto precisi, richiedono una nuova, più complessa modellizzazione teorica.


La comprensione dei dati di raggi cosmici (carichi, ma anche raggi gamma e neutrini) non può prescindere da un approccio a **multi-frequenza e multi-messaggero**, e da campagne di misure agli **acceleratori**.



# The GeV excess at the Galactic center

Goodenough+'09, Vitale+'09, Abazajian+PRD'12, Hooper+PDU'13, Daylan+PDU'16, Calore+JCAP'15, Cholis+JCAP'15, Calore+PRD'15, Ajello+2015, Linden+PRD'16, Ackermann+ApJ'17, ...500+papers

Found with template fitting (Calore+JCAP2015), adaptive template fitting (Storms+ 2017), weighted Likelihood (Di Mauro PRD2021, Abdollahi AJS2020) photon counts statistics (1pPDF: Calore, FD,+ PRL2021; NPTF Lee+2016), machine Learning (List+PRL20, Mishra-JCAPSharma+PRD21, Caron+22), wavelet transforms (Bartels+PRL16)



MurgiaAR 2020


No matter the method, the GC excess is statistically significant



# GAPS detector to fly in Antarctic by 2023

Dedicated to antideuteron searches

F. Rogers et al. *Astrop. Phys.* 2023





Secure results on very low energy antiprotons



# Fit of Galactic pulsar populations to AMS-02 $e^+$ data

Orusa, Di Mauro, FD, Manconi JCAP 2021



The contribution of pulsars to  $e^+$  is dominant above 100 GeV and may have different features.  
 $E > 1$  TeV: unconstrained by data.  
 Secondaries forbid evidence of sharp cut-off.  
 No need for Dark Matter, indeed



# Possible origin of anti-helium: anti-clouds, anti-stars

V. Poulin et al. PRD 2019




FIG. 4. Abundance of  $\bar{\text{H}}$ ,  $\bar{\text{D}}$  and  $\bar{^4\text{He}}$  with respect to that of  $\bar{^3\text{He}}$  as a function of the (anti-)baryon-to-photon ratio  $\bar{\eta}$ . The *Planck* value is represented by the grey band. The value required by the *AMS-02* experiment is shown by the orange band.

Anti-clouds: require anisotropic BBN  
for the right  $\bar{^3\text{He}}/\bar{^4\text{He}}$

*AMS-02* measures are local, *Planck*'s  
ones averaged over the Universe

Exotic mechanism for segregation of  
anti-clouds is needed


Traces in  $\bar{p}$  and  $\bar{\text{D}}$

One anti-star could make the job.  
How did they survive?



# Antideuteron perspectives

Serksnyte et al, PRD 2022



Low energy window keeps being a discovery field  
 Uncertainties on  $P_c$  is  $\pm 70\%$

See also Korsmeier, FD, Fornengo PRD 2018




# Hardening of nuclear spectra

If it were acceleration, the hardening would be the same for primaries and secondaries


Recchia & Gabici MNRAS 2014; Pluskin & Zirakashvili ApJ 2013; Zatsepin & Sokolskaya A&A 2006; Yuan+ PRD 2011

Tomassetti & FD A&A 2021



An hardening is expected from fragmentation in the SNRs

Tomassetti & Oliva ApJL 2017



An hardening is expected from reacceleration in the SNRs

Also Tomassetti & FD ApJL 2015

Interpretations of current data is not clear,  
and still hampered by spallation cross sections



# Antideuterons from relic WIMPS

FD, Fornengo, Salati PRD 62 (2000)043003

In order for fusion to take place, the two antinucleons must have low kinetic energy

Kinematics of spallation reactions prevents the formation of very low antiprotons (antineutrons).

At variance, dark matter annihilates almost at rest


$$\frac{dN_{\bar{D}}}{dE_{\bar{D}}} = \left( \frac{4 P_{\text{coal}}^3}{3 k_{\bar{D}}} \right) \left( \frac{m_{\bar{D}}}{m_p m_n} \right) \sum_{F,h} B_{\chi h}^{(F)} \left\{ \frac{dN_p^h}{dE_{\bar{p}}} \left( E_{\bar{p}} = \frac{E_{\bar{D}}}{2} \right) \right\}^2$$

Background and DM have different kinematics and source spectra



# Perspectives with antideuterons

Bess Polar-II @ ICRC2023



GAPS - dedicated to antineutron searches -  
will fly from Antarctica Dec 2024