




Finanziato
dall'Unione europea
NextGenerationEU



Ministero
dell'Università
e della Ricerca



Italiadomani
PIANO NAZIONALE
DI RIPRESA E RESILIENZA



New BSM ideas

M. Nardecchia
Sapienza University of Rome



New BSM ideas?

 You

I have been invited to a conference to talk about "New Beyond the Standard Model ideas", what I should talk about?

 ChatGPT

That sounds like an exciting opportunity! When discussing "New Beyond the Standard Model ideas," you could explore various theoretical and experimental avenues that extend beyond the current framework of particle physics. Here are some potential topics you could cover:

1. Supersymmetry (SUSY)
2. String Theory and Extra Dimensions
3. Grand Unified Theories (GUTs)
4. Dark Matter and Dark Energy
5. Neutrino Physics
6. Experimental Searches for New Particles
7. Alternative Quantum Field Theories
8. Astroparticle Physics

 You

What about light new physics?

 ChatGPT

Light new physics refers to theoretical frameworks and experimental searches for new phenomena that could manifest at accessible energy scales or through subtle effects in observable phenomena. Here are some potential avenues to explore when discussing light new physics:


1. Axions and Axion-Like Particles (ALPs)
2. Light Scalars and Pseudoscalars
3. Hidden Photons
4. New Gauge Bosons
5. Neutrino Mass Mechanisms
6. Composite Particles
7. Fifth Forces and Modifications to Gravity



Finanziato
dall'Unione europea
NextGenerationEU



Ministero
dell'Università
e della Ricerca



Italiadomani
PIANO NAZIONALE
DI RIPRESA E RESILIENZA



Light-Heavy New Physics connections

Some biased comments about NP physics + recent results based on

“Light vectors coupled to anomalous currents with harmless Wess-Zumino terms”,

L. Di Luzio, MN, C.Toni,
JHEP, arXiv:2204.05945

“Closing in on new chiral leptons at the LHC”,

D. Barducci, L. Di Luzio, MN, C.Toni
JHEP, arXiv: 2311.10130

New Physics



Experimental evidences:

- Neutrino masses
- Dark Matter
- Baryon Asymmetry of the Universe
- (Gravity)

Theoretical problems/puzzles/hints:

- Hierarchy or Naturalness problem
- Flavour puzzle
- Strong CP problem
- Family replication
- GUT
-


What is the energy scale of New Physics at its coupling to the Standard Model?
Can we study it on shell?




$$m_H^2 = m_{\text{tree}}^2 + \delta m_H^2$$

$$\delta m_H^2 = \frac{3}{\sqrt{2}\pi^2} G_F m_t^2 \Lambda^2 \approx (0.3 \Lambda)^2$$

Coupling vs Mass Range




Why? Why not



- Didn't find NP at high energy
- Some problems can be addressed in this regime: axions, portal to DM sector, etc.
- Ideas for new experiments on smaller scale than LHC
- No "No-Lose Theorem"

Why?




- SUSY
- Composite Higgs

UV completing the light NP physics could be phenomenologically important!

Light New Vector and (Anomalous) Currents

Consider, for example, a light vector coupled to the baryon number current:




$$\mathcal{L} \supset g_X \frac{1}{3} Z'_{B\mu} (\bar{q} \gamma^\mu q)$$

(Notation: X=Z' interchangeably in what follows)

Naively, NP physics at low energy implies two parameters: the coupling and the mass

SM + X EFT is non-renormalizable and the current is **anomalous** at quantum level:



$$\partial^\mu J_\mu^{\text{baryon}} = \frac{A}{16\pi^2} \left(g^2 W_{\mu\nu}^a (\tilde{W}^a)^{\mu\nu} - g'^2 B_{\mu\nu} \tilde{B}^{\mu\nu} \right)$$

A=3/2

EFT must be completed at a scale $\lesssim \frac{4\pi m_X}{g_X} / \left(\frac{3g^2}{16\pi^2} \right)$ [Preskill 1991]

UV physics implies the presence of Wezz-Zumino terms in the EFT:

$$\mathcal{L} \supset C_B g_X g'^2 \epsilon^{\mu\nu\rho\sigma} X_\mu B_\nu \partial_\rho B_\sigma + C_W g_X g^2 \epsilon^{\mu\nu\rho\sigma} X_\mu (W_\nu^a \partial_\rho W_\sigma^a + \frac{1}{3} g \epsilon^{abc} W_\nu^a W_\rho^b W_\sigma^c)$$

(Need also to specify regularisation of loop diagrams)

Background:

- D'Hoker, Farhi, 1984
- Preskill 1991
- Feruglio, Masiero, Maiani 1992

New constraints for light vectors:

- Dror, Lasenby, Pospelov 1707.01503
- Dror, Lasenby, Pospelov 1705.06726

Other pheno aspects, example DM:

- Ismail, Katz, Racco 1707.00709

UV renormalizable models:



- Michaels, Yu 2020.00021

Energy / Mass Enhancement



Generically, there are E/m_X enhancements of the longitudinal polarisation of X leading to strong bounds.

$$\Gamma(A \rightarrow BX) \propto \frac{g_X^2}{m_X^2}$$

$$\mathcal{L} \supset g_X d_i d_j X_\mu \bar{d}_j \gamma^\mu \mathcal{P}_L d_i + \text{h.c.} + \dots,$$



example from 1707.01503



Constraints from:

$$B \rightarrow KX$$

$$K \rightarrow \pi X$$

$$Z \rightarrow \gamma X$$

Very strong constraints!

However there is a **specific** point in the EFT that do not exhibit this behaviour

What kind of UV physics is associated with that?

Anomalons

The light vector can be elementary or composite. If elementary, it is associated to a conserved symmetry and the current have to be non anomalous. In the UV we need new states to cancel the mass independent part of the triangular diagram:

$$\mathcal{M}^{\mu\nu\rho} \equiv \sum_{f, f_{\text{SM}}} X_\mu \text{ [diagram] },$$

Extra states are called anomalons in what follows. NP carries quantum number under both the SM gauge symmetry and the new symmetry

Anomalons are **chiral** with respect to the full group SM x U(1). The E/m enhancement is due to longitudinal d.o.f

Chiral fermions cannot get an explicit mass term.

SM and the light vector \mathcal{X} (also keeping the Goldstone mode ξ , see App. A for details)

$$\begin{aligned} \mathcal{L}_{\text{EFT}}^{\text{U}(1)\mathcal{X}} \supset & g_X g'^2 \frac{C_{BB}}{24\pi^2} \epsilon^{\alpha\mu\nu\beta} \mathcal{X}_\alpha B_\mu \partial_\beta B_\nu + g_X g'^2 \frac{C_{ab}}{24\pi^2} \epsilon^{\alpha\mu\nu\beta} \mathcal{X}_\alpha W_\mu^a \partial_\beta W_\nu^b \\ & + g_X g g' \frac{C_{aB}}{24\pi^2} \epsilon^{\alpha\mu\nu\beta} \mathcal{X}_\alpha W_\mu^a \partial_\beta B_\nu + g_X g g' \frac{C_{Ba}}{24\pi^2} \epsilon^{\alpha\mu\nu\beta} \mathcal{X}_\alpha B_\mu \partial_\beta W_\nu^a \\ & + g_X g^2 \frac{D_{ab}}{48\pi^2} \frac{\xi}{m_\mathcal{X}} \epsilon^{\alpha\mu\beta\nu} (\partial_\alpha W_\mu^a) (\partial_\beta W_\nu^b) + g_X g'^2 \frac{D_{BB}}{48\pi^2} \frac{\xi}{m_\mathcal{X}} \epsilon^{\alpha\mu\beta\nu} (\partial_\alpha B_\mu) (\partial_\beta B_\nu) \\ & + g_X g g' \frac{D_{aB}}{24\pi^2} \frac{\xi}{m_\mathcal{X}} \epsilon^{\alpha\mu\beta\nu} (\partial_\alpha W_\mu^a) (\partial_\beta B_\nu), \end{aligned}$$

Anomalons have to pick up a mass from EW and/or U(1) symmetry breaking

$$H \bar{f}_L f_R \quad \text{and/or} \quad S \bar{f}_L f_R \quad S \propto e^{i\xi}$$

Enhancement can be understood with the **equivalence** theorem

Goldstone term disappear if the Anomalons take mass from EW only

A Class of Renormalizable Models

Field	Lorentz	SU(3) _C	SU(2) _L	U(1) _Y	U(1) _X
q_L^i	$(\frac{1}{2}, 0)$	3	2	1/6	$\alpha_B/3$
u_R^i	$(0, \frac{1}{2})$	3	1	2/3	$\alpha_B/3$
d_R^i	$(0, \frac{1}{2})$	3	1	-1/3	$\alpha_B/3$
ℓ_L^i	$(\frac{1}{2}, 0)$	1	2	-1/2	α_i
e_R^i	$(0, \frac{1}{2})$	1	1	-1	α_i
H	$(0, 0)$	1	2	1/2	0
\mathcal{L}_L	$(\frac{1}{2}, 0)$	1	2	$\mathcal{Y} - 1/2$	$X_{\mathcal{L}_L}$
\mathcal{L}_R	$(0, \frac{1}{2})$	1	2	$\mathcal{Y} - 1/2$	$X_{\mathcal{L}_R}$
\mathcal{E}_L	$(\frac{1}{2}, 0)$	1	1	$\mathcal{Y} - 1$	$X_{\mathcal{E}_L}$
\mathcal{E}_R	$(0, \frac{1}{2})$	1	1	$\mathcal{Y} - 1$	$X_{\mathcal{E}_R}$
\mathcal{N}_L	$(\frac{1}{2}, 0)$	1	1	\mathcal{Y}	$X_{\mathcal{N}_L}$
\mathcal{N}_R	$(0, \frac{1}{2})$	1	1	\mathcal{Y}	$X_{\mathcal{N}_R}$
ν_R^α	$(0, \frac{1}{2})$	1	1	0	$X_{\nu_R}^\alpha$
\mathcal{S}	$(0, 0)$	1	1	0	$X_{\mathcal{S}}$

$$-\mathcal{L}_Y = y_1 \bar{\mathcal{L}}_L \mathcal{E}_R H + y_2 \bar{\mathcal{L}}_R \mathcal{E}_L H + y_3 \bar{\mathcal{L}}_L \mathcal{N}_R \tilde{H} + y_4 \bar{\mathcal{L}}_R \mathcal{N}_L \tilde{H} + \text{h.c.}$$

- All gauge anomalies have to cancel
- SM quantum number set by a single parameter Y
- Charges are such that anomalous don't couple to S
- **Mass purely from EW**

Two important phenomenological implications:

1) **non-decoupling New Physics**



$$M_{anom} = y \frac{v}{\sqrt{2}} \quad M_{anom} \lesssim 600 \text{ GeV} \quad (y = \sqrt{4\pi})$$

2) **very large coupling to the Higgs**

Anomalous are **heavy leptons** with exotic charges.
No QCD charge is allowed, because of Higgs overproduction at the LHC

Higgs Physics

Main effects are the radiative two-body decays of the Higgs into gauge bosons: $h \rightarrow Z\gamma$ and $h \rightarrow \gamma\gamma$



$$R_{\gamma\gamma, Z\gamma} = \frac{|A_{\gamma\gamma, Z\gamma}^{\text{SM}} + A_{\gamma\gamma, Z\gamma}^{\text{BSM}}|^2}{|A_{\gamma\gamma, Z\gamma}^{\text{SM}}|^2}$$

$$A_{\gamma\gamma}^{\text{BSM}} \simeq \frac{4}{3} (1 + 4Y^2),$$

$$A_{Z\gamma}^{\text{BSM}} \simeq \frac{2}{3} [1 - (1 + 8Y^2) \text{tg}_w^2]$$


- BSM contributions independent from the mass of new states when $m_h \ll M_{anom}$

- Diphoton contribution can be consistent only if $A_{\gamma\gamma}^{\text{BSM}} \simeq -2A_{\gamma\gamma}^{\text{SM}}$ $|Y| \approx \frac{3}{2}$

- In 2204.05945, clean **prediction** for the Z-photon channel $R_{Z\gamma} \approx 2.1$

Higgs to Z gamma

[ATLAS & CMS 2309.03501]



the signal strength is shown in Figure 3. The observed (expected) signal strength at the 68% confidence level is $\mu = 2.0^{+1.0}_{-0.9}$ (1.0 ± 0.9) for the ATLAS analysis, $\mu = 2.4^{+1.0}_{-0.9}$ ($1.0^{+1.0}_{-0.9}$) for the CMS analysis, and $\mu = 2.2 \pm 0.6$ (stat.) $^{+0.3}_{-0.2}$ (syst.) = 2.2 ± 0.7 (1.0 ± 0.6 (stat.) ± 0.2 (syst.)) = 1.0 ± 0.6 for their combination.

This motivated us to study further the phenomenology of chiral leptons in [2311.10130](#) (JHEP)

Direct Searches of Anomalons

Recap: interesting case when new states have SM quantum number: $\mathcal{L}_{L,R} = \begin{pmatrix} \mathcal{N}_{\mathcal{L}} \\ \mathcal{E}_{\mathcal{L}} \end{pmatrix}_{L,R} \sim (\mathbf{1}, \mathbf{2})_Y$, $\mathcal{E}_{L,R} \sim (\mathbf{1}, \mathbf{1})_{Y-\frac{1}{2}}$, $\mathcal{N}_{L,R} \sim (\mathbf{1}, \mathbf{1})_{Y+\frac{1}{2}}$

After EWSB 4 Dirac fields with electric charges about 2 and 1. At the LHC, phenomenology depends if there are stable anomalons or not.

$$|Y| \approx \frac{3}{2}$$

Stable anomalons give charged tracks at the LHC

$$M_{Q=2e} > 1030 \text{ GeV}$$

Adapting and rescaling the ATLAS analysis [arXiv:2303.13613](#)

$$M_{Q=e} > 600 \text{ GeV}$$

Adapting and rescaling the CMS analysis [arXiv:1609.08382](#)

Unstable anomalons are present for some specific U(1) charge assignments and $Y = -\frac{3}{2}$

There is a mixing with the SM charged leptons: $-\mathcal{L}_{\text{mix}} = \lambda_{i,R} \bar{L}_L^i H N_R + \lambda_{i,L} \bar{\mathcal{L}}_L \tilde{H} e_R^i + h.c.$

Doubly charged states decays into same-sign lepton pair via: $\Psi^{\mathcal{E}_i} \rightarrow W^- \ell^- \rightarrow \ell^- \ell^- \cancel{E}_T$

$$M_{Q=2e} > 600 \text{ GeV}$$

Adapting and rescaling the ATLAS analysis [arXiv:1710.09748](#)

Model is **perturbative excluded** when considering a more refined bound using unitarity for Yukawa coupling $M_{anom} \lesssim 400 \text{ GeV}$

(Models that explains a possible large effects in Z-photon can be constructed starting from this benchmark)

LHC and Non Decoupling New Physics

- In absence of New Physics, **old fashioned solutions of the hierarchy problem cannot be ruled out**
- This is because SUSY and Composite Higgs model can be continuously deformed to the SM

$$\lim_{m_{\text{soft}} \rightarrow \infty} \text{MSSM} = SM$$

$$\lim_{f \rightarrow \infty} \text{Composite Higgs} = SM$$

- What is getting worst is the little hierarchy problem. (The tuning of the heavy NP parameters to get the EW scale)

$$\Delta = \frac{v^2}{m_{\text{soft}}^2}$$

$$\Delta = \frac{v^2}{f^2}$$

- **Non-decoupling** New Physics does not have such smooth limit.

$$M_{anom} = y \frac{v}{\sqrt{2}}$$

$$M_{anom} \lesssim 400 \text{ GeV}$$

$$M_{anom} \gtrsim 600 \text{ GeV}$$

- Perturbativity

- Direct searches
and Higgs Physics

- LHC is closing completely the window for new chiral fermions

(Biased) Conclusions

- BSM physics exists, but experimental data does not favor any specific physics scenario.
- Traditional solutions to the hierarchy problem remain relevant despite requiring finer tuning.
- There is a growing interest in light and weakly coupled new physics, though its justification is arguable. However some models are very interesting and motivated.
- When considering light new physics, it is crucial to bear in mind an ultraviolet (UV) completion of the low-energy effective field theory (EFT).
- For instance, a light vector coupled to anomalous currents may exhibit E/mass enhancement but an explicit UV model can circumvent this feature by predicting that new fermions are chiral and acquire mass from the Higgs VEV.
- New chiral fermions are heavily constrained by LHC searches, if not entirely ruled out.

Backup

Perturbative Unitarity

- Unitarity (an axiom of QFT), focus 2 → 2 scattering


$$\begin{aligned}
 SS^\dagger &= 1 \\
 S &= 1 + iT
 \end{aligned}
 \quad \longrightarrow \quad
 \frac{1}{2i} (a_{fi}^J - a_{if}^{J*}) \geq \sum_{h \in 2\text{-particle}} a_{hf}^{J*} a_{hi}^J
 \quad a_{fi}^J \propto \langle f|T|i\rangle_J$$

- For $f = i$ (optical theorem)

$$\text{Im } a_{ii}^J \geq |a_{ii}^J|^2 \quad \longrightarrow \quad (\text{Re } a_{ii}^J)^2 + \left(\text{Im } a_{ii}^J - \frac{1}{2} \right)^2 \leq \frac{1}{4}$$

- In practical perturbative calculations S-matrix unitarity is always approximate

- signals breakdown of perturbative expansion



$$|\text{Re } (a_{ii}^J)^{\text{Born}}| \leq \frac{1}{2}$$

• Works both in the EFT and in explicit renormalizable models