

Describing inclusive semileptonic $b \rightarrow c\ell\bar{\nu}$ decays to higher precision for inclusive V_{cb} determinations

La Thuile 2024 - Les Rencontres de Physique de la Vallée d'Aoste

Ilija S. Milutin¹

in collaboration with
Thomas Mannel¹ and K. Keri Vos²

Based on: JHEP 02 (2024) 226 [ArXiv: 2311.12002]


¹CPPS, Universität Siegen, Germany

²GWFP, Maastricht University and Nikhef, The Netherlands



The V_{cb} puzzle

- $b \rightarrow c(u)l\bar{\nu}$ decays allow for studies of physics beyond the Standard Model and extraction of CKM parameters
- V_{cb} is an important input for many Standard Model predictions
- Long-standing puzzle between different determinations
 - exclusive: known-final state, e.g. $B \rightarrow D l \bar{\nu}$
 - **inclusive**: summed over all possible final states, e.g. $B \rightarrow X_c l \bar{\nu}$



Bernlochner, Prim, Vos (*Eur. Phys. J. Spec. Top.* (2024))

Setting up the Heavy Quark Expansion

- m_b is large compared to Λ_{QCD} \rightarrow power series in Λ_{QCD}/m_b

Chay, Georgi, Bigi, Shifman, Uraltsev, Vainshtein, Manohar, Wise, Neubert, Mannel,...

- Split the momentum of the heavy quark as $p_b = m_b v + k$ where $v^2 = 1$
- Expand in residual momentum $k \sim iD$

- Perform Operator Product Expansion (OPE)

Chay, Georgi, Bigi, Shifman, Uraltsev, Vainshtein, Manohar, Wise, Neubert, Mannel,...

$$d\Gamma = d\Gamma^{(3)} + \frac{1}{m_b^2} d\Gamma^{(5)} + \frac{1}{m_b^3} d\Gamma^{(6)} + \frac{1}{m_b^4} d\Gamma^{(7)} + \dots, \quad d\Gamma^{(n)} = \sum_i \mathcal{C}_i^{(n)} \langle B | \mathcal{O}_i^{(n)} | B \rangle$$

- $d\Gamma^{(n)}$ are power series in $\mathcal{O}(\alpha_s)$
- $\mathcal{C}_i^{(n)}$ perturbative Wilson coefficients
- $\mathcal{O}^{(n)}$ local operators of dimension n
- $\langle B | \mathcal{O}_i^{(n)} | B \rangle$ non-perturbative matrix elements \rightarrow need to be extracted from data

- The number of matrix elements increases very fast with the dimension:

$$d\Gamma = d\Gamma^{(3)} + \frac{1}{m_b^2}d\Gamma^{(5)} + \frac{1}{m_b^3}d\Gamma^{(6)} + \frac{1}{m_b^4}d\Gamma^{(7)} + \frac{1}{m_b^5}d\Gamma^{(8)} + \dots$$

- $d\Gamma^{(3)}$: Partonic result ($d\Gamma^{(4)} = 0$ due to Heavy Quark Symmetries)
- $d\Gamma^{(5)}$: 2 parameters

$$2m_B\mu_\pi^2 = -\langle B|\bar{b}_v(iD)^2b_v|B\rangle$$

$$2m_B\mu_G^2 = \langle B|\bar{b}_v(-i\sigma^{\mu\nu})(iD_\mu)(iD_\nu)b_v|B\rangle$$

- $d\Gamma^{(6)}$: 2 parameters

$$2m_B\rho_D^3 = \frac{1}{2}\langle B|\bar{b}_v[iD_\mu, [ivD, iD^\mu]]b_v|B\rangle$$

$$2m_B\rho_{LS}^3 = \frac{1}{2}\langle B|\bar{b}_v\{iD_\mu, [ivD, iD_\nu]\}(-i\sigma^{\mu\nu})b_v|B\rangle$$

- $d\Gamma^{(7)}$: 9 parameters (at tree level) Mannel, Turczyk, Uraltsev [1009.4622]; Kobach, Pal [1704.00008]
- $d\Gamma^{(8)}$: 18 parameters (at tree level) Mannel, Turczyk, Uraltsev [1009.4622]; Kobach, Pal [1704.00008]

Reparametrization Invariance

- In HQE, choice of v_μ is not unique Dugan, Golden, Grinstein, Chen, Luke, Manohar, Hill, Solon, Heinonen,...
- Lorentz invariance of QCD \rightarrow **Reparametrization Invariance (RPI)** imposed under infinitesimal change $v_\mu \rightarrow v_\mu + \delta v_\mu$, leading to

$$\delta_{\text{RP}} v_\mu = \delta v_\mu, \quad \delta_{\text{RP}} iD_\mu = -m_b \delta v_\mu, \quad \delta_{\text{RP}} b_v(x) = im_b(x \cdot \delta v) b_v(x)$$

Reparametrization Invariance

- In HQE, choice of v_μ is not unique Dugan, Golden, Grinstein, Chen, Luke, Manohar, Hill, Solon, Heinonen,...
- Lorentz invariance of QCD \rightarrow **Reparametrization Invariance** (RPI) imposed under infinitesimal change $v_\mu \rightarrow v_\mu + \delta v_\mu$, leading to


$$\delta_{\text{RP}} v_\mu = \delta v_\mu, \quad \delta_{\text{RP}} iD_\mu = -m_b \delta v_\mu, \quad \delta_{\text{RP}} b_v(x) = im_b(x \cdot \delta v) b_v(x)$$

- **RPI relates different orders** n in $1/m_b$ expansion Mannel, Vos [1802.09409]

$$\delta_{\text{RP}} \mathcal{C}^{(n)} \propto m_b \mathcal{C}^{(n+1)}$$

- This allows us to find combinations of operators which are RPI
- Lorentz invariant quantities depend on reduced set of RPI parameters

Counting RPI operators



- Up to $1/m_b^4$: total of 8 independent parameters Mannel, Vos [1802.09409]
- At $1/m_b^5$, we find **only 10 RPI operators** and 8 non-RPI operators Mannel, ISM, Vos [2311.12002]

New inclusive V_{cb} determination

- Need an RPI-observable \rightarrow only depend on reduced set of RPI operators

New inclusive V_{cb} determination

- Need an RPI-observable \rightarrow only depend on reduced set of RPI operators
- **Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

New inclusive V_{cb} determination

- Need an RPI-observable \rightarrow only depend on reduced set of RPI operators
- **Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

- Data \rightarrow values for reduced set of RPI parameters up to $1/m_b^4 \rightarrow Br(\bar{B} \rightarrow X_c \ell \bar{\nu}) \rightarrow$
 $|V_{cb}^{\text{incl}}| = (41.69 \pm 0.63) \times 10^{-3}$ Bernlochner, Vos, et al. [2205.10274]

New inclusive V_{cb} determination

- Need an RPI-observable \rightarrow only depend on reduced set of RPI operators
- **Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

- Data \rightarrow values for reduced set of RPI parameters up to $1/m_b^4 \rightarrow Br(\bar{B} \rightarrow X_c \ell \bar{\nu}) \rightarrow$
 $|V_{cb}^{\text{incl}}| = (41.69 \pm 0.63) \times 10^{-3}$ Bernlochner, Vos, et al. [2205.10274]

- **First determination** of V_{cb} up to $\mathcal{O}(1/m_b^4)$ and **first extraction** of $1/m_b^4$ matrix elements from data
- Agreement at **1 – 2 σ level** with previous $\mathcal{O}(1/m_b^3)$ determinations

Finauri, Gambino [2310.20324]; Bordone, Capdevila, Gambino [2107.00604]; Alberti, Gambino, Healey, Nandi [1411.6560]; Gambino, Schwanda [1307.4551]

Where do we currently stand?

Γ	tree	α_s	α_s^2	α_s^3
Partonic	✓	✓	✓	✓
$1/m_b^2$	✓	✓		
$1/m_b^3$	✓	✓		
$1/m_b^4$	✓			
$m_b^{\text{kin}}/\bar{m}_c$		✓	✓	✓

$\langle (q^2)^n \rangle$	tree	α_s	α_s^2	α_s^3
Partonic	✓	✓		
$1/m_b^2$	✓	✓		
$1/m_b^3$	✓	✓		
$1/m_b^4$	✓			

- Green: known perturbative corrections Jezabek, Kuhn (1989); Melnikov (2008); Pak, Czarnecki (2008); Becher, Boos, Lunghi (2007); Alberti, Gambino, Nandi (2014); Mannel, Pivovarov, Rosenthal (2015); Gambino, Healey, Turczyk (2016); Mannel, Pivovarov (2020); Fael, Schonwald, Steinhauser (2020, 2021)
- Next step in HQE is dimension 8: $1/m_b^5$



Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]

“genuine” $\frac{1}{m_b^5}$

&


Intrinsic Charm $\frac{1}{m_b^3 m_c^2}$



Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]

“genuine” $\frac{1}{m_b^5}$ & Intrinsic Charm $\frac{1}{m_b^3 m_c^2}$




- Numerically: $m_c^2 \sim m_b \Lambda_{\text{QCD}}$
- Dimension 8 contains enhanced terms which contribute like $1/m_b^4$ terms

Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]

“genuine” $\frac{1}{m_b^5}$ & **Intrinsic Charm** $\frac{1}{m_b^3 m_c^2}$



- Numerically: $m_c^2 \sim m_b \Lambda_{\text{QCD}}$
- Dimension 8 contains enhanced terms which contribute like $1/m_b^4$ terms
- We need $1/m_b^3 \times 1/m_c^2$ contributions to complete the calculation at $1/m_b^4$
 - We calculate the full dimension 8 contributions
 - We extract the **Intrinsic Charm** contribution
 - We find that only **1 combination of parameters** describes the IC in the q^2 -moments

Mannel, ISM, Vos [2311.12002]


How to get the matrix elements...

- At $1/m_b^5 \rightarrow$ too many HQE parameters to fit to data
- **LLSA**: Lowest-Lying-State Approximation to estimate through known parameters μ_π^2 , μ_G^2 , ρ_D^3 and ρ_{LS}^3


Mannel, Turczyk, Uraltsev [1009.4622]

$$\langle B | \bar{b} A \Gamma b(0) | B \rangle = \frac{1}{2m_B} \sum_n \langle B | \bar{b} A b(0) | n \rangle \cdot \langle n | \bar{b} C \Gamma b(0) | B \rangle$$


- $A = iD_{\mu_1} \dots iD_{\mu_k}$, $C = iD_{\mu_{k+1}} \dots iD_{\mu_n}$




Mannel, ISM, Vos [2311.12002]



Mannel, ISM, Vos [2311.12002]



Mannel, ISM, Vos [2311.12002]



Mannel, ISM, Vos [2311.12002]

- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions

Conclusions and outlook

- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an **unexpectedly small** overall contribution of the dimension-8 operators

Conclusions and outlook

- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an **unexpectedly small** overall contribution of the dimension-8 operators
- q^2 moments to $1/m_b^5$ available in open-source library **kolya**

Conclusions and outlook


- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an **unexpectedly small** overall contribution of the dimension-8 operators
- q^2 moments to $1/m_b^5$ available in open-source library **kolya**
- We provide easy calculation of **any kinematic moment** to $1/m_b^5$ for $B \rightarrow X_c \ell \bar{\nu}$

Mannel, ISM, Vos [2311.12002]

- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an **unexpectedly small** overall contribution of the dimension-8 operators
- q^2 moments to $1/m_b^5$ available in open-source library **kolya**
- We provide easy calculation of **any kinematic moment** to $1/m_b^5$ for $B \rightarrow X_c \ell \bar{\nu}$
Mannel, ISM, Vos [2311.12002]
- Work in progress:
 - Full $|V_{cb}|^{\text{incl}}$ fit to $1/m_b^5$
 - Further understand theory uncertainties through the study of **Quark-Hadron Duality Violation**

Back-up: Cancellation without LLSA

- Assume for all dimension 8 operators that $X_i^5 \sim \Lambda_{\text{QCD}}^5$ and vary signs:



- **Step 1:** expand charm propagator ($Q^\mu = m_b v^\mu$)

$$\begin{aligned} -iS_{\text{BGF}} &= \frac{1}{\not{Q} + i\not{D} - m_c} \\ &= \frac{1}{\not{Q} - m_c} - \frac{1}{\not{Q} - m_c} (i\not{D}) \frac{1}{\not{Q} - m_c} \\ &\quad + \frac{1}{\not{Q} - m_c} (i\not{D}) \frac{1}{\not{Q} - m_c} (i\not{D}) \frac{1}{\not{Q} - m_c} + \dots \end{aligned}$$

Back-up: Calculation of forward matrix element

- **Step 2:** insert in forward matrix element

$$\begin{aligned} T = & \left[\Gamma \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha b_\beta \rangle \\ & - \left[\Gamma \frac{1}{\not{Q} - m_c} \gamma^\mu \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha (iD_\mu) b_\beta \rangle \\ & + \left[\Gamma \frac{1}{\not{Q} - m_c} \gamma^\mu \frac{1}{\not{Q} - m_c} \gamma^\nu \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha (iD_\mu) (iD_\nu) b_\beta \rangle \\ & + \dots \end{aligned}$$

Back-up: Calculation of forward matrix element

- **Step 3:** Determine **Trace formula**

- Start at dimension 8 \rightarrow lengthy, but systemically calculable
- Compute dim-7, including $1/m_b$ correction through e.o.m.

$$(ivD)b_v = -\frac{1}{2m_b}(i\cancel{D})(i\cancel{D})b_v$$

- ...
- Compute dim-3, including corrections up to $1/m_b^5$

$$\langle \bar{b}_\alpha b_\beta \rangle = 2m_B \left(\frac{1 + \psi}{4} + \frac{1}{8m_b^2}(\mu_G^2 - \mu_\pi^2) + \mathcal{O}(1/m_b^6) \right)_{\beta\alpha}$$

- Need full (non-RPI) set of **basic parameters** up to $1/m_b^5$
- **Step 4:** Compute the trace with the geometric series