


Accelerazione di particelle tramite laser intensi ed applicazioni ai beni culturali, (biomediche, materiali, ambiente, IA)

Patrizio Antici
antici@emt.inrs.ca

The advent of High-Power lasers thanks to the CPA

Focus on: CPA

CPA: 25 years of intense laser pulses



Basic working principle of CPA. An initially very short but low energy laser pulse (top left) is first stretched, then amplified and then recompressed (bottom right). Numbers are indicative only.



The field of high-power lasers is booming (investments of over >1 B€ in 10 years!) (including INFN – FLAME/Eupraxia!)

BELLA 30 M€

APPOLON 50 M€

CALA 65 M€

ALLS 30 M€

FLAME/INFN-PI 6 M€



2000-2005 Experimental Discovery of laser-accelerated particles

2000

2004

VOLUME 85, NUMBER 14 PHYSICAL REVIEW LETTERS 2 OCTOBER 2000

Intense High-Energy Proton Beams from Petawatt-Laser Irradiation of Solids

R. A. Snavely,^{1,2} M. H. Key,¹ S. P. Hatchett,¹ T. E. Cowan,¹ M. Roth,^{3,4} T. W. Phillips,¹ M. A. Stoyer,¹ E. A. Henry,¹ T. C. Sangster,¹ M. S. Singh,¹ S. C. Wilks,¹ A. MacKinnon,¹ A. Offenberger,^{4,6} D. M. Pennington,¹ K. Yasuike,^{5,*} A. B. Langdon,¹ B. F. Lasinski,¹ J. Johnson,⁶ M. D. Perry,¹ and E. M. Campbell¹

¹Lawrence Livermore National Laboratory, University of California, P.O. Box 808, Livermore, California 94550

²Department of Physics, University of California, Davis, California 95616

³GSI Laboratory, Darmstadt, Germany

⁴University of Alabama, Huntsville, Alabama

⁵George C. Marshall Space Flight Center, Huntsville, Alabama

⁶University of California, San Diego, California 92093

(Received 19 January 2000)

High quality and high energy protons are accelerated by high intensity lasers...

Intense laser beams have been used to accelerate protons to high energies. The maximum energy is transferred to 2×10^{10} protons of energy >10 MeV. The energy spectrum exhibits a sharp high-energy cutoff as high as 58 MeV on the axis of the beam which decreases in energy with increasing off axis angle. Proton induced nuclear processes have been observed and used to characterize the beam.

PACS numbers: 41.75.Jc, 07.77.Ka, 52.50.Jm

The generation of fast protons from laser irradiated solid surfaces is well understood [1,2] and attributable to electrostatic fields produced by hot electrons acting on protons from adsorbed hydrocarbons. An empirical power law relationship between the mean proton energy and intensity x (wavelength) [2] ($I\lambda^2$) was identified and proton energies up to a few MeV were observed for $I\lambda^2$ up to 10^{15} W cm⁻² μm² in nanosecond pulses [1]. Scaling laws for the maximum energy were similarly derived [3].

Chirped pulse amplification (CPA) laser technology [4] has since enabled widespread generation of terawatt power and the first petawatt laser [5]. CPA lasers generate pulses in the range 20 fs to 1 ps. 10 MeV protons were observed with a 1 ps CPA laser at $I\lambda^2 = 10^{19}$ W cm² μm² consistent with the previous scaling [6,7].

New mechanisms of ion acceleration have been studied with CPA lasers. Ponderomotive pressure of the laser radiation causes radial acceleration when laser beams are focused in gas jets and subcritical density plasmas [8] and also axial acceleration into solid targets [9]. Coulomb explosion of molecules [10] and clusters [11] has produced energetic ions. The ion energies from these newly studied processes have been <1 MeV/nucleon.

We report a laser induced proton beam with much higher particle energy (up to 58 MeV). The maximum energy is about $2 \times$ higher than is given by extrapolating the previous scaling law [7] to the $30 \times$ higher intensity of this work. A distinctive feature is emission with good beam collimation perpendicular to the rear-irradiated surface(s) of the target. Similar rear surface beaming of protons of energy <1 MeV was reported in nanosecond pulse CO₂ laser experiments [12]. Our observed high proton energies open up access to nuclear processes.

The experiments used a CPA laser system generating 1 PW pulses of 500 fs duration [5]. With $f/3$ parabolic mirror focusing, the peak intensity was 3×10^{20} W cm⁻²

in a focal spot of 9 μm FWHM with 30% of the energy inside the first minimum. Amplified spontaneous emission in a 4 ns period before the main pulse had 10^{-4} of the main pulse energy and there was a 3×10^{-2} prepulse 2 ns before the main pulse. This precursor radiation generated a plasma which was measured by time-resolved optical interferometry. The main pulse intensity was 3×10^{20} W cm⁻² in a plane 70 μm from a flat CH polymer target with an approximately exponential fall to lower densities having a scale length of 10 μm.

The proton beam was measured by a 10 μm thick RC film which changes through polymerization of a diacetylene active layer, or conversion to dark blue in proportion to the dose. The film is 100 μm thick and weighs 100 ergs/g. A 90° cone of emission of protons, cones, and sheets of RC film was used. The cone is 4 cm deep at its apex, which was 10 cm from the target. The laser axis was the axis of the cone. The RC film response was calibrated absolutely and the image data were analyzed by digital densitometry and transformed geometrically to produce contour plots of dose as a function of angle illustrated in Fig. 1.

The data in Fig. 1 show a collimated intense proton beam emitted perpendicular to the rear target surface of a Au target at 45° to the laser axis. Its angular width narrows to 10° in the image through 600 μm of Ta. The beam is rather uniform in intensity with near circular sharp boundaries. There is a low intensity wide-angle background, which is discussed elsewhere and is due to escaping relativistic electrons [14]. Similar results were seen for normal incidence, S polarization, and target thickness down to 20 μm. The proton beam profiles from CH targets have nonuniform edges and exhibit internal fine structure as illustrated in Fig. 2 for a 100-μm thick CH target at normal incidence. There was typically a 5 times greater proton induced dose recorded from CH relative to Au targets. Data


**Proton beams:
High number
High energy
Very laminar**



1000 x more compact + other properties (short duration)


E-field_{max} ≈ 50-100 MeV /meter (Breakdown)

1 m



RF cavity

Courtesy of W. Mori & L. da Silva




Plasma cavity






100 μm



Opening of new labs for proton acceleration (e.g. FLAME@INFN-LNF or ALLS@EMT-INRS)



Proton parameters of high-power lasers enable many applications !



I. Improvement of a laser-driven proton beamline (high-repetition rate targetry system – implementation of QP)

www.nature.com/scientificreports

SCIENTIFIC REPORTS

OPEN Design and optimization of a compact laser-driven proton beamline

Received: 13 November 2017
Accepted: 19 March 2018
Published online: 19 April 2018

M. Scisciò^{1,2}, M. Migliorati¹, L. Palumbo¹ & P. Antici²

Laser-accelerated protons, generated by irradiating a solid target with a short, energetic laser pulse at high intensity ($I > 10^{18} \text{ W cm}^{-2}$), represent a complementary if not outperforming source compared to conventional accelerators, due to their intrinsic features, such as high beam charge and short

Design and optimization of a laser-PIXE beamline for material science applications

A. Morabito^{1,2}, M. Scisciò^{2,3}, S. Veltri⁴, M. Migliorati² and P. Antici⁵

¹ELI-ALPS, ELI-HU Non profit Ltd., Dugonics ter 13, Szeged, 6720, Hungary; ²INFN and University of Rome, Via Scarpa 14, 00161 Roma, Italy; ³ENEA, Fusion and Nuclear Safety Department, C. R. Frascati, Via E. Fermi 45, Frascati, 00044 Roma, Italy; ⁴Institute of Low Temperature and Structure Research, Polish Academy of Sciences, PL-50422 Wrocław, Poland and ⁵INRS-EMT, 1650 Boul. Lionel Boulet, J3X 1S2, Varennes, Canada

Abstract

Multi-MeV proton beams can be generated by irradiating thin solid foils with ultra-intense ($>10^{18} \text{ W/cm}^2$) short laser pulses. Several of their characteristics, such as high bunch charge and short pulse duration, make them a complementary alternative to conventional radio frequency-based accelerators. A potential material science application is the chemical analysis of small quantities of (OP) materials. The complete characterization of the bulk material composition


Laser and Particle Beams
cambridge.org/lpb

Research Article

Cite this article: Morabito A, Scisciò M, Veltri S, Migliorati M, Antici P (2019). Design and optimization of a laser-PIXE beamline for material science applications. *Laser and Particle Beams* 1–10. <https://doi.org/10.1017/S0263034619000600>

Received: 11 June 2019
Revised: 1 August 2019
Accepted: 6 August 2019


- Possible PhD Thesis:
- Beam dynamic simulations
 - Beam line design






PHYSICAL REVIEW SPECIAL TOPICS - ACCELERATORS AND BEAMS 16, 011302 (2013)

Intrinsic normalized emittance growth in laser-driven electron accelerators



M. Migliorati,^{1,2} A. Bacci,³ C. Benedetti,^{4,*} E. Chiodroni,³ M. Ferrario,³ A. Mostacci,^{1,2} L. Palumbo,^{1,2} A. R. Rossi,³ L. Serafini,⁵ and P. Antici^{1,2,3,†}




2. Laser-PIXE



Example of PIXE to analyze the pigment's composition of *The Trivulzio portrait* by Antonello da Messina



2. Laser-PIXE for Cultural Heritage



C. Hargoues / C2RMF / Aglaé / CNRS Photothèque

High
Inte
>10

Une statuette en bronze du forum de Bavay analysée avec Aglaé.


SCIENTIFIC REPORTS

OPEN Laser-Accelerated Proton Beams as Diagnostics for Cultural Heritage

M. Barberio^{1,2}, S. Veltri³, M. Scisciò^{3,4} & P. Antici^{1,3}

Received: 21 October 2015
Accepted: 07 December 2016
Published: 01 March 2017

This paper introduces the first use of laser-generated proton beams as diagnostic for materials of interest in the domain of Cultural Heritage. Using laser-accelerated protons, as generated by interaction of a high-power short-pulse laser with a solid target, we can produce proton-induced X-ray emission spectroscopies (PIXE). By correctly tuning the proton flux on the sample, we are able to perform the PIXE in a single shot without provoking more damage to the sample than conventional methodologies. We verify this by experimentally irradiating materials of interest in the Cultural Heritage with laser-accelerated protons and measuring the PIXE emission. The morphological and chemical analysis of the sample before and after irradiation are compared in order to assess the damage provoked to the artifact. Monte Carlo simulations confirm that the temperature in the sample stays safely below the melting point. Compared to conventional diagnostic methodologies, laser-driven PIXE has the advantage of being potentially quicker and more efficient.



Explore different materials /
Limits of this and combined
technique / Damage ?

Possible PhD Thesis:

- Laser-PIXE for In-Air analysis of artifacts
- Laser-PIXE for Archeometry


2. A new field: Laser-diagnostics for Cultural Heritage

We are not alone !



ELI Beamlines and its new project: Non-Destructive Laser-driven Heritage Testing

Institute of Physics of the Academy of Sciences of the Czech Republic / ELI Beamlines Research Center is launching a new project called Non-Destructive Methods of Heritage Testing, financed by the Prague Operational Program (MOP) and the State Research Development and Innovation Operational Program (MOP).




Intense Laser Irradiation Laboratory

National Institute of Optics – National Council of Research




CONSIGLIO NAZIONALE DELLE RICERCHE
ISTITUTO NAZIONALE DI OTTICA



Laser-PIXE

Laser driven Particle Induced X-ray Emission: source development and X-ray spectral/spatial analysis


The PIXE (Particle Induced X-ray Emission) is a multi-elemental, quantitative, highly sensitive (trace elements with concentrations in parts-per-million), rapid, non-invasive and non-destructive analysis technique to determine the composition of surface layers of a sample. The technique is based on spectroscopy of the characteristic X radiation emitted by each atom following irradiation with proton/ion beams. PIXE requires protons with MeV energies (2-3 MeV typically) that are currently produced with particle accelerators whose dimensions (and costs) represent a limit for the application of the same technique outside of large research in-



Current topic:

- Analysis of violins with University of Pavia (M. Malagodi)
- Analysis of specific metallic items with Uniroma3 (L.Tortora)

3. Possible thesis: Laser-diagnostics for Environment (Exhaust detection)



Nuclear Instruments and Methods in Physics Research B 363 (2015) 86–91



Contents lists available at ScienceDirect

Nuclear Instruments and Methods in Physics Research B

journal homepage: www.elsevier.com/locate/nimb



Present role of PIXE in atmospheric aerosol research




Willy Maenhaut*

Ghent University, Department of Analytical Chemistry, Krijgslaan 281, S12, BE-9000 Gent, Belgium
Department of Pharmaceutical Sciences, University of Antwerp (Campus Drie Eiken), Universiteitsplein 1, BE-2610 Antwerp, Belgium

Is laser-PIXE applicable for Exhaust-detection ?

4. Possible thesis: Laser-PIXE for early cancer and Alzheimer detection



There is a correlation between the alteration of selected trace elements concentration in brain tissue and certain neurological diseases (Alzheimer's disease, Parkinson's disease, and brain tumors). Most attention has been given to Fe, Cu and Zn and their concentration in different parts of the brain.

In addition:

- **Laser-driven protons for Gene-manipulation (Modification of DNA)**
- **Laser-driven protons for radiotherapy (zebrafish irradiation)**

5. Artificial intelligence for improving beamlines and data mining (together with Uniroma I/INFN ?)



PHYSICAL REVIEW LETTERS **126**, 104801 (2021)

ARTICLE




<https://doi.org/10.1038/s41467-020-20245-6> OPEN

Automation and control of laser wakefield accelerators using Bayesian optimization

R. J. Shalloo^{1,6*}, S. J. D. Dann², J.-N. Gruse¹, C. I. D. Underwood³, A. F. Antoine⁴, C. Arran³, M. Backhouse¹, C. D. Baird^{2,3}, M. D. Balcazar⁴, N. Bourgeois², J. A. Cardarelli⁴, P. Hatfield⁵, J. Kang⁶, K. Krushelnick⁴, S. P. D. Mangles¹, C. D. Murphy³, N. Lu⁷, J. Osterhoff⁸, K. Pöder⁸, P. P. Rajeev², C. P. Ridgers³, S. Rozario¹, M. P. Selwood³, A. J. Shahani⁷, D. R. Symes², A. G. R. Thomas⁴, C. Thornton², Z. Najmudin¹ & M. J. V. Streeter¹

Bayesian Optimization of a Laser-Plasma Accelerator

Sören J alas^{1,4*}, Manuel Kirchen¹, Philipp Messner^{2,1,3}, Paul Winkler^{3,1}, Lars Hübner^{3,1}, Julian Dirkwinkel³, Matthias Schnepf¹, Remi Lehe⁴, and Andreas R. Maier^{3,1}
¹Center for Free-Electron Laser Science and Department of Physics Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany
²International Max Planck Research School for Ultrafast Imaging & Structural Dynamics, Luruper Chaussee 149, 22761 Hamburg, Germany
³Deutsches Elektronen Synchrotron (DESY), Notkestraße 85, 22607 Hamburg, Germany
⁴Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA




Collaboration with:



Mila

6. Laser-accelerated proton for material science application



ARTICLE

DOI: 10.1038/s41467-017-02675-x




OPEN

Laser-accelerated particle beams for stress testing of materials

M. Barberio¹, M. Scisciò^{1,2}, S. Vallières¹, F. Cardelli^{1,2}, S.N. Chen^{3,4}, G. Famularo⁵, T. Gangolf⁶, G. Revet^{3,4}, A. Schiav², M. Senzacqua² & P. Antici¹

Possible Thesis:

- 1) Stress test on different materials
- 2) Doping of semiconductor materials using laser-driven protons



PhD thesis in...

... any other topic that YOU would like to pursue and is feasible (and compatible with the labs)...




International working mode



LLNL @ San Francisco


Current 5 Students:
2 with Uniroma1
1 with Uniroma3
1 with U. Bordeaux
1 only INRS



LULI @ Ecole Polytechnique Paris



LLC @ Lund



FORTH @ Crete

ALLS @ Montréal

We publish... have fun... and they talk about us...

M. Barberio, S. Giusepponi, S. Vallières, M. Scisciò, M. Celino, P. Antici
Ultra-Fast Metallic Nanoparticle Synthesis using Laser-Accelerated Protons
Scientific Reports, in press

C. Bienvenue, S. Vallières, S. Payeur, and P. Antici
Fast Submicrometric Absolute Positioning of Flat Reflective Surface by Monochromatic Interferometry
Review of Scientific Instruments, in press

S. Vallières, M. Barberio, M. Scisciò, E. d'Humieres, and P. Antici
Enhanced laser-driven proton acceleration using ultrasmall nanoparticles
Physical Review Accelerators and Beams 22, 091303 (2019)

A. Morabito, M. Scisciò, S. Veltri, M. Migliorati, P. Antici
Design and optimization of a dedicated laser driven hybrid proton beam-line for cultural heritage applications
Laser and Particle Beams, <https://doi.org/10.1017/S0263034619000600>

S. Vallières, C. Bienvenue, P. Puyuelo-Valdes, M. Salvadori, E. d'Humières, F. Schiettekatte, and P. Antici
Low-energy proton calibration and energy-dependence linearization of E
Review of Scientific Instruments 90, 083301 (2019)

M. Barberio, E. Skantzakis, P. Antici
Material analysis using laser-plasma driven luminescence spectroscopy
Journal of Luminescence 214, 116603 (2019)

M. Barberio, E. Skantzakis, S. Sorieul, P. Antici
Pigment darkening as case study of In-Air Plasma Induced Luminescence
Science Advances 5, 6, eaar6228 DOI: 10.1126/sciadv.aar6228 (2019)

M. Barberio, P. Antici
Laser-PIXE using laser-accelerated proton beams
Scientific Reports 9, 6855 (2019)

Home Berita **Jurnal IPTEK** Tour & Travel Penerbit

Teknik Laser Untuk Diagnosa & Analisa K Seni Rapuh

(STAGING-POINT.COM, 2019/06/14) In **Jurnal IPTEK**



ULTIME NOTIZIE

27/10/2019 - 19:45 : DIFESA: DELEGAZIONE BRASILIANA VISITA I REPARTI DELLA BRIGATA INFORMAZIONI TATTICHE

Home Italiani nel Mondo Esteri Italia Cultura Economia italiana nel mondo Regioni Migrazioni Vaticano Chi Siamo

Sei in: Home / Italiani nel Mondo / Comunità

ITALIA-QUÉBEC: LA COOPERAZIONE VA SU SCIENCE

16:44

- Nell'ambito del **Programma esecutivo di cooperazione culturale, scientifica e tecnologica tra Italia e Québec** per il periodo 2017-2019, un'innovativa ricerca nel campo delle tecnologie laser per spiegare i pigmenti di alcune opere pittoriche, tra cui i celebri Girasoli di Van Gogh, hanno ottenuto la pubblicazione sulla prestigiosa rivista *Science Advances*.
L'In-Air Plasma Induced Luminescence (PIXE) può essere usata non solo per analizzare lo stato di conservazione delle opere d'arte, ma anche per appurare l'autenticità, la composizione e la provenienza. L'In-Air Plasma Induced Luminescence (PIXE) (Particle-Induced X-ray Emission), pertanto si configura come una tecnica di analisi non distruttiva, con potenzialità che non riguardano unicamente la preservazione del patrimonio artistico, ma coinvolgono anche altri settori del patrimonio culturale.



Email Stampa PDF

SUNDAY, OCTOBER 27 2019 | HOME About Terms DMCA Sitemap

Home / australia / Powerful lasers for fragile works of art

POWERFUL LASERS FOR FRAGILE WORKS OF ART

australia navva June 10, 2019 australia

Facebook Twitter Google+ LinkedIn Pinterest



Chi siamo Eventi Lingua e Cultura Avvisi Contatti

In evidenza >> Consolato Generale d'Italia a Montréal Ambasciata d'Italia Ottawa Biblioteca Opportunità

Home > Gli Eventi > Calendario

Gli Eventi

Calendario

Archivio

Eventi in Italia

Italia-Québec, la cooperazione va su "Science Advances"

Data: 11/10/2017

Nell'ambito del **Programma esecutivo di cooperazione culturale, scientifica e tecnologica tra Italia e Québec** per il periodo 2017-2019, un'innovativa ricerca nel campo delle tecnologie laser per spiegare i pigmenti di alcune opere pittoriche, tra cui i celebri Girasoli di Van Gogh, hanno ottenuto la pubblicazione sulla prestigiosa rivista *Science Advances*.

Antici, assieme ai suoi collaboratori e a un gruppo di ricerca internazionale, che vede coinvolta anche l'Università della Calabria, l'acceleratore di particelle francese AIFIRA ed il laboratorio Greco FORTH, ha sviluppato un nuovo strumento diagnostico per esaminare lo stato di conservazione delle opere pittoriche: il metodo di ricerca impiega la tecnologia laser per spiegare come mai alcuni pigmenti di alcune opere pittoriche, tra cui i celebri Girasoli di Van Gogh, abbiano perso la loro lucentezza originaria nel corso del tempo, diventando via via più scuri.

Antici - laureato in ingegneria presso la Sapienza Università di Roma, con dottorato in fisica presso l'École Polytechnique (Francia) e dottorato in ingegneria presso la Sapienza spiega: "La nuova



Advantages / Disadvantages of a joint PhD (but there are no disadvantages...)

- International exposure (network)
- Multidisciplinary
- Multitasking (theory/ experiment)
- Salary is almost doubled (!)
- High productivity and flexibility
- Flexibility in travels and working time
- Good language skills
- Result oriented
- Able to work independently
- Fast decision making, fast acting

-> **Good Job insertion**

What is INRS – Center Energy, Materials, Telecommunications

~40 multidisciplinary professors (7 in photonics), about 190 Students/PhDs (small center, dynamic, low overheads)



RESEARCH Infosource Inc.
Canada's Source of R&D Intelligence

About | Canada's Innovation Leaders | **Top 50 Research Universities** | Top 100 Corporate R&D Spenders | Top 50 Research Colleges | Top 40 Research Hospitals | Explore | Other Resources

Excellence center, only graduate students



SAVE THE STUDENT! FINANCE BANKING MA

Here's the complete top 20 list of t

1. London, United Kingdom
2. Munich, Germany
2. Seoul, South Korea
4. Zurich, Switzerland
5. Melbourne, Australia
6. Berlin, Germany
7. Tokyo, Japan
8. Paris, France
9. Sydney, Australia
10. Edinburgh, United Kingdom
11. Boston, USA
12. Hong Kong SAR, Hong Kong
13. Singapore, Singapore
14. Montreal, Canada



#1 Montreal



Top 50 Research Universities / 2021 / Top 50 List



Canada's Top 50 Research Universities 2021

Rank	2020	2019	University	Sponsored Research		Change	FY2020 \$000	FY2019 \$000	Region	Province
				2020	2019					
25	25		Institut national de la recherche scientifique	\$66,280	\$71,889	-7.8	\$419.5	\$98.6	S	QC
9	8		McMaster University	\$353,530	\$371,599	-4.9	\$369.4	\$68.9	M	ON
5	6		University of Calgary	\$457,296	\$487,805	-6.3	\$298.5	\$67.8	M	AB
12	11		University of Saskatchewan	\$223,328	\$243,531	-8.3	\$207.9	\$65.0	M	SK
3	3		McGill University	\$628,642	\$606,489	3.7	\$344.1	\$63.5		
1	1		University of Toronto	\$1,234,278	\$1,089,287	13.3	\$446.6	\$61.0		
2	2		University of British Columbia	\$652,637	\$624,465	4.5	\$267.7	\$57.5		
44	44		University of Winnipeg	\$13,169	\$12,344	6.7	\$43.9	\$56.6		

Highly "italianized"



Carriero Adriatico
Dir. Resp. Giancarlo Laurenzi
Tiratura: 19.206 Diffusione: 13.369 Lettori: 288.000

Prestigioso premio in Quebec allo scienziato Rosci

Insegna in Canada all'Inrs
Si occupa di nanotecnologie e dei semiconduttori


IL RICONOSCIMENTO
ASCOLI Il professor Federico Rosci è stato premiato con il prestigioso premio "Marie-Victorin" la più alta onorificenza conferita dal governo del Quebec a una persona che ha condotto una carriera di elevato livello nella ricerca delle scienze naturali e dell'ingegneria. Rosci è il nipote del direttore dello storico stabilimento Elettrocaraman di Ascoli e figlio di Enzo, docente per anni all'università di Trieste. C'è, dunque, un pizzico di Ascoli in questo riconoscimento. Il prestigio è doppio perché si tratta di uno "straniero" premiato in terra canadese. Rosci è da tempo impegnato nello studio della nanotecnologia, delle nanostrutture e dei semiconduttori e autore di numerose pubblicazioni sull'argomento. «È uno dei ricercatori più importanti della mia carriera e sono fortunato che un comitato di pari mi abbia scelto», ha dichiarato Rosci. «Sono onorato che il mio nome sia accanto ai vincitori del passato, considerandolo il loro successore». Rosci ha ottenuto un dottorato di ricerca all'Università Sapienza di Roma ed ha fatto esperienze in Danimarca prima di operare in Canada. Ha ricevuto numerosi riconoscimenti, tra cui la Medaglia Rumford della Royal Society of Canada. È stato anche il primo

giganti e progressione
zione e di tra i
Cristian
Rosci
Recherche étudiant-es ou stagiaires
Fabio Boschini
Il professor Federico Rosci
Emanuele Orgiu






designed by freepik.com



Interested ?



**... collaborate or join us !
Patrizio.antici@inrs.ca**

Thank you for your attention !



www.ipat-lab.net



