



Contribution ID: 34

Type: not specified

### Supporting ${}^3P_0$ Quark-Pair Creation using Landau Gauge Green's Functions

Phenomenological evidence suggests that strong decays of low-excitation hadrons often involve the creation of a light quark-antiquark pair with zero angular momentum, known as the  ${}^3P_0$  mechanism, derived from a scalar bilinear. Despite Quantum Chromodynamics being mediated perturbatively by spin-one gluons and exhibiting chiral symmetry in its Lagrangian, a scalar decay term appears spontaneously upon chiral symmetry breaking. We explore this by employing the quark-gluon vertex in the Landau gauge and the nonperturbative effects recently clarified, alongside a constant chromoelectric field similar to the Schwinger pair production in Quantum Electrodynamics. We compare this to a two-field insertion diagram in QED and argue that the relevant quantum numbers for discussing production are  ${}^3\Sigma_0$ ,  ${}^3\Sigma_1$ , and  ${}^3\Pi_0$ , analogous to those in diatomic molecules. Our results indicate significant contributions from the third decay mechanism, supporting the  ${}^3P_0$  phenomenology at momenta at or below the fermion mass scale. However, ultrarelativistic fermions predominantly exhibit  ${}^3\Sigma_1$  quantum numbers. In QED,  ${}^3\Sigma_0$  is dominant, whereas in QCD,  ${}^3\Pi_0$  prevails at sub-GeV momenta due to the requirement to form a color singlet.

**Primary authors:** SALAS-BERNÁRDEZ, Alexandre (Universidad Complutense de Madrid); LLANES-ESTRADA, Felipe J. (Universidad Complutense de Madrid); Prof. ALKOFER, Reinhard (Graz University)

**Presenter:** SALAS-BERNÁRDEZ, Alexandre (Universidad Complutense de Madrid)