

# Breakdown of collinear factorisation in exclusive $\pi^0\gamma$ photoproduction due to Glauber pinch

QCD Evolution 2024  
Pavia, Italy

Saad Nabeebaccus  
IJCLab



université  
PARIS-SACLAY



Gluedynamics

May 28, 2024

Based on 2311.09146 with Jakob Schönleber, Lech Szymanowski and Samuel Wallon  
This project is supported by the European Union's Horizon 2020 research and innovation programme under Grant agreement no. 824093

# Introduction

## Exclusive photon-meson photoproduction

Original motivation: Extraction of **chiral-odd** GPDs at *leading* twist.

►  $\gamma N \rightarrow \rho_T^0 \pi^+ N'$ :

M. El Beiyad, B. Pire, M. Segond, L. Szymanowski, S. Wallon: [1001.4491]

# Introduction

## Exclusive photon-meson photoproduction

Original motivation: Extraction of **chiral-odd** GPDs at *leading* twist.

▶  $\gamma N \rightarrow \rho_T^0 \pi^+ N'$ :

M. El Beiyad, B. Pire, M. Segond, L. Szymanowski, S. Wallon: [1001.4491]

▶  $\gamma N \rightarrow \gamma MN'$ :

–  $M = \rho^0$ : R. Boussarie, B. Pire, L. Szymanowski, S. Wallon: [1609.03830]

–  $M = \pi^\pm$ : G. Duplančić, K. Passek-Kumerički, B. Pire, L. Szymanowski, S. Wallon: [1809.08104]

–  $M = \pi^\pm, \rho^{0,\pm}$ , wider kinematical coverage, various observables:  
G. Duplančić, **S.N.**, K. Passek-Kumerički, B. Pire, L. Szymanowski, S. Wallon:  
[2212.00655, 2302.12026]

# Introduction

## Exclusive photon-meson photoproduction

Original motivation: Extraction of **chiral-odd** GPDs at *leading* twist.

▶  $\gamma N \rightarrow \rho_T^0 \pi^+ N'$ :

M. El Beiyad, B. Pire, M. Segond, L. Szymanowski, S. Wallon: [1001.4491]

▶  $\gamma N \rightarrow \gamma MN'$ :

–  $M = \rho^0$ : R. Boussarie, B. Pire, L. Szymanowski, S. Wallon: [1609.03830]

–  $M = \pi^\pm$ : G. Duplančić, K. Passek-Kumerički, B. Pire, L. Szymanowski, S. Wallon: [1809.08104]

–  $M = \pi^\pm, \rho^{0,\pm}$ , wider kinematical coverage, various observables:  
G. Duplančić, **S.N.**, K. Passek-Kumerički, B. Pire, L. Szymanowski, S. Wallon:  
[2212.00655, 2302.12026]

Richer kinematics of 3-body final state processes allows the sensitivity of GPDs wrt  $x$  to be probed (beyond moment-type dependence, e.g. in DVCS)

J. Qiu, Z. Yu: [2305.15397]

# Introduction

Exclusive photon-meson photoproduction

$$\gamma(q) + N(p_1) \rightarrow \gamma(k) + M(p_M) + N'(p_2)$$

$$\mathcal{A} = \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_M(z)$$


# Introduction

## Exclusive photon-meson photoproduction

$$\gamma(q) + N(p_1) \rightarrow \gamma(k) + M(p_M) + N'(p_2)$$


$$\mathcal{A} = \int_{-1}^1 dx \int_0^1 dz T(x, \xi, z) H(x, \xi, t) \Phi_M(z)$$

- ▶ **Fully differential** cross-section differential covering  $S_{\gamma N}$  from  $\sim 4 \text{ GeV}^2$  to  $20000 \text{ GeV}^2$ .
- ▶ **Good statistics** at various experiments, particularly at *JLab*.
- ▶ Polarisation asymmetries also sizeable.
- ▶ **Small  $\xi$**  limit of quark GPDs can be studied at collider experiments.




# Introduction

Is collinear factorisation justified?



large angle factorisation  
à la Brodsky Lepage



We thus argue *collinear factorisation* of the amplitude at **large**  
 $M_{\gamma M}^2$ ,  $t'$ ,  $u'$ , and **small**  $t$ .

$$\begin{aligned} t &= (p_2 - p_1)^2, & u' &= (p_M - q)^2, \\ t' &= (k - q)^2, & S_{\gamma N} &= (q + p_1)^2. \end{aligned}$$

# Introduction

Is Collinear factorisation justified?

- ▶ Recently, factorisation has been proved for the process  $\pi N \rightarrow \gamma\gamma N'$  by J. Qiu, Z. Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by J. Qiu, Z. Yu [2210.07995]



# Introduction

Is Collinear factorisation justified?

- ▶ Recently, factorisation has been proved for the process  $\pi N \rightarrow \gamma\gamma N'$  by J. Qiu, Z. Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by J. Qiu, Z. Yu [2210.07995]
- ▶ The proof relies on having **large  $p_T$** , rather than large invariant mass (e.g. photon-meson pair).

# Introduction

## Is Collinear factorisation justified?

- ▶ Recently, factorisation has been proved for the process  $\pi N \rightarrow \gamma\gamma N'$  by J. Qiu, Z. Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by J. Qiu, Z. Yu [2210.07995]
- ▶ The proof relies on having **large  $p_T$** , rather than large invariant mass (e.g. photon-meson pair).
- ▶ In fact, NLO computation has been performed for  $\gamma N \rightarrow \gamma\gamma N'$  by O. Grocholski, B. Pire, P. Sznajder, L. Szymanowski, J. Wagner [2110.00048, 2204.00396]
- ▶ Also, NLO computation for  $\gamma\gamma \rightarrow \pi^+\pi^-$  by crossing symmetry G. Duplancic, B. Nizic: [hep-ph/0607069].

# Introduction

## Is Collinear factorisation justified?

- ▶ Recently, factorisation has been proved for the process  $\pi N \rightarrow \gamma\gamma N'$  by J. Qiu, Z. Yu [2205.07846].
- ▶ This was extended to a wide range of  $2 \rightarrow 3$  exclusive processes by J. Qiu, Z. Yu [2210.07995]
- ▶ The proof relies on having **large  $p_T$** , rather than large invariant mass (e.g. photon-meson pair).
- ▶ In fact, NLO computation has been performed for  $\gamma N \rightarrow \gamma\gamma N'$  by O. Grocholski, B. Pire, P. Sznajder, L. Szymanowski, J. Wagner [2110.00048, 2204.00396]
- ▶ Also, NLO computation for  $\gamma\gamma \rightarrow \pi^+\pi^-$  by crossing symmetry G. Duplancic, B. Nizic: [hep-ph/0607069].

*Issues with exclusive  $\pi^0\gamma$  photoproduction...*

# Introduction

## Gluon GPD contributions to exclusive $\pi^0\gamma$ photoproduction

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to *gluon GPD contributions*.

# Introduction


## Gluon GPD contributions to exclusive $\pi^0\gamma$ photoproduction

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to *gluon GPD contributions*.
- ▶ A total of 24 diagrams contribute in this case (compared to 20 diagrams from quark GPD contributions), with 6 groups of 4 related by symmetries ( $x \rightarrow -x$  and  $z \rightarrow 1 - z$  separately).

# Introduction


## Gluon GPD contributions to exclusive $\pi^0\gamma$ photoproduction

- ▶ Because of the quantum numbers of  $\pi^0$  ( $J^{PC} = 0^{-+}$ ), the exclusive photoproduction of  $\pi^0\gamma$  is also sensitive to *gluon GPD contributions*.
- ▶ A total of 24 diagrams contribute in this case (compared to 20 diagrams from quark GPD contributions), with 6 groups of 4 related by symmetries ( $x \rightarrow -x$  and  $z \rightarrow 1 - z$  separately).
- ▶ Diagrams amount to connecting photons to the following two topologies.



# Result assuming collinear factorisation

## Specific diagram



$$CF \sim \frac{\text{Tr} \left[ \not{p}_M \gamma^5 \not{\epsilon}_k \left( \not{k} + z \not{p}_M \right) \gamma^j \left( \not{q} - (x - \xi) \not{p} - \bar{z} \not{p}_M \right) \not{\epsilon}_q \left( -(x - \xi) \not{p} - \bar{z} \not{p}_M \right) \gamma^i \right]}{[2z k p_M] [-2(x - \xi) q p - 2\bar{z} q p_M + 2\bar{z}(x - \xi) p p_M + i\epsilon] [2\bar{z}(x - \xi) p p_M + i\epsilon]}$$

$$\xrightarrow{x \rightarrow \xi, \bar{z} \rightarrow 0} \propto \frac{x - \xi}{[(x - \xi) + A\bar{z} - i\epsilon][\bar{z}(x - \xi) + i\epsilon]}, \quad A \equiv \frac{q p_M}{q p} > 0.$$

(Assuming  $p_M$  is along minus direction)

# Result assuming collinear factorisation

## Specific diagram

Need to dress coefficient function CF with gluon GPD  $\left(\frac{H_g(x)}{(x-\xi+i\epsilon)(x+\xi-i\epsilon)}\right)$ , and DA  $(z\bar{z})$ . This gives

$$\mathcal{A} \sim \frac{\bar{z}(x-\xi)H_g(x)}{(x-\xi+i\epsilon)[(x-\xi)+A\bar{z}-i\epsilon][\bar{z}(x-\xi)+i\epsilon]}$$
$$\longrightarrow \frac{H_g(x)}{[(x-\xi)+A\bar{z}-i\epsilon][x-\xi+i\epsilon]}$$



# Result assuming collinear factorisation

## Specific diagram

Need to dress coefficient function CF with gluon GPD  $\left(\frac{H_g(x)}{(x-\xi+i\epsilon)(x+\xi-i\epsilon)}\right)$ , and DA ( $z\bar{z}$ ). This gives


$$\begin{aligned} \mathcal{A} &\sim \frac{\bar{z}(x-\xi)H_g(x)}{(x-\xi+i\epsilon)[(x-\xi)+A\bar{z}-i\epsilon][\bar{z}(x-\xi)+i\epsilon]} \\ &\longrightarrow \frac{H_g(x)}{[(x-\xi)+A\bar{z}-i\epsilon][x-\xi+i\epsilon]} \end{aligned}$$

The integral over  $z$  and  $x$  diverges if the GPD  $H_g(x)$  is non-vanishing at  $x = \xi$ :

$$\begin{aligned} &\int_{-1}^1 dx \int_0^1 dz \frac{1}{[(x-\xi)+A\bar{z}-i\epsilon][x-\xi+i\epsilon]} \\ &\supset \int_{-1}^1 dx \frac{\ln(x-\xi-i\epsilon)}{[x-\xi+i\epsilon]} \implies \text{divergent imaginary part!} \end{aligned}$$

# Result assuming collinear factorisation

## Specific diagram



$$\int_{-1}^1 dx \int_0^1 dz \frac{1}{[(x - \xi) + A\bar{z} - i\epsilon][x - \xi + i\epsilon]}$$

$\implies$  The “*pinching*” is caused by propagators  $D_a$  and  $D_b$ .

# Result assuming collinear factorisation

## Full Amplitude

What about the sum of diagrams?

$$\begin{aligned} \sum \mathcal{A} &\sim \frac{z\bar{z}(x^2 - \xi^2) \left[ -\alpha \left[ (x^2 - \xi^2)^2 (1 - 2z\bar{z}) + 8x^2\xi^2z\bar{z} \right] - (1 + \alpha^2) z\bar{z}(x^4 - \xi^4) \right] H_g(x)}{z\bar{z}[x - \xi + i\epsilon]^2 [\bar{z}(x + \xi) - \alpha z(x - \xi) - i\epsilon] [z(x - \xi) + \alpha\bar{z}(x + \xi) - i\epsilon]} \\ &\times \frac{1}{[x + \xi - i\epsilon]^2 [\bar{z}(x - \xi) + \alpha z(x + \xi) - i\epsilon] [z(x + \xi) - \alpha\bar{z}(x - \xi) - i\epsilon]} \\ &\xrightarrow{x \rightarrow \xi, \bar{z} \rightarrow 0} \propto \frac{\left[ -\alpha \left[ (x^2 - \xi^2)^2 (1 - 2z\bar{z}) + 8x^2\xi^2z\bar{z} \right] - (1 + \alpha^2) z\bar{z}(x^4 - \xi^4) \right] H_g(x)}{[x - \xi + i\epsilon] [2\xi\bar{z} - \alpha(x - \xi) - i\epsilon] [(x - \xi) + 2\xi\alpha\bar{z} - i\epsilon]} \end{aligned}$$

# Result assuming collinear factorisation

## Full Amplitude

What about the sum of diagrams?

$$\sum \mathcal{A} \sim \frac{z\bar{z}(x^2 - \xi^2) \left[ -\alpha \left[ (x^2 - \xi^2)^2 (1 - 2z\bar{z}) + 8x^2\xi^2 z\bar{z} \right] - (1 + \alpha^2) z\bar{z} (x^4 - \xi^4) \right] H_g(x)}{z\bar{z} [x - \xi + i\epsilon]^2 [\bar{z}(x + \xi) - \alpha z(x - \xi) - i\epsilon] [z(x - \xi) + \alpha\bar{z}(x + \xi) - i\epsilon]}$$
$$\times \frac{1}{[x + \xi - i\epsilon]^2 [\bar{z}(x - \xi) + \alpha z(x + \xi) - i\epsilon] [z(x + \xi) - \alpha\bar{z}(x - \xi) - i\epsilon]}$$
$$\xrightarrow{x \rightarrow \xi, \bar{z} \rightarrow 0} \propto \frac{\left[ -\alpha \left[ (x^2 - \xi^2)^2 (1 - 2z\bar{z}) + 8x^2\xi^2 z\bar{z} \right] - (1 + \alpha^2) z\bar{z} (x^4 - \xi^4) \right] H_g(x)}{[x - \xi + i\epsilon] [2\xi\bar{z} - \alpha(x - \xi) - i\epsilon] [(x - \xi) + 2\xi\alpha\bar{z} - i\epsilon]}$$


Full amplitude (anti)-symmetric in  $x \rightarrow -x$  and  $z \rightarrow \bar{z}$  for (anti)-symmetric GPD. (only symmetric result shown above)

$\implies$  *divergence survives*, and actually adds up.

# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude




$$\xi = 0.5$$

# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude




$$\xi = 0.5$$

- Unfortunately, no cancellations between the 4 corners.

# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude




$$\xi = 0.5$$

- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ In  $\gamma\gamma \rightarrow MM$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.

# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude



$$\xi = 0.5$$


- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ In  $\gamma\gamma \rightarrow MM$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.
- ▶ Indication of problem with naive collinear factorisation?  
At twist-2??



# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude




$$\xi = 0.5$$

- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ In  $\gamma\gamma \rightarrow MM$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.
- ▶ Indication of problem with naive collinear factorisation?  
At twist-2??
- ▶ *Can this divergence be understood from a theoretical point of view?*

# Result assuming collinear factorisation

Singularity structure of the full amplitude

'Phase Space' for amplitude



$$\xi = 0.5$$

- ▶ Unfortunately, no cancellations between the 4 corners.
- ▶ In  $\gamma\gamma \rightarrow MM$ , only ERBL region exists, no poles are crossed, and endpoint contributions are suppressed by DAs.
- ▶ Indication of problem with naive collinear factorisation?  
At twist-2??
- ▶ *Can this divergence be understood from a theoretical point of view?*  
YES!  $\implies$  [S. N., J. Schönleber, L. Szymanowski, S. Wallon: 2311.09146]

# Reduced diagram analysis

## Libby-Sterman power counting

- ▶ How to obtain the dominant contribution of an amplitude (in QCD) in a certain specific kinematics (e. g. collinear)?
  - ⇒ Libby-Sterman power counting rule [Phys.Rev.D 18 (1978) 3252; Phys.Rev.D 18 (1978) 4737]

# Reduced diagram analysis

## Libby-Sterman power counting

- ▶ How to obtain the dominant contribution of an amplitude (in QCD) in a certain specific kinematics (e. g. collinear)?  
⇒ Libby-Sterman power counting rule [Phys.Rev.D 18 (1978) 3252; Phys.Rev.D 18 (1978) 4737]
- ▶ Extensively used in factorisation proofs [Collins: Foundations of perturbative QCD]
- ▶ Basic idea is to identify regions of loop momenta of partons (also number of partons), which gives the dominant contribution to the full amplitude.

# Reduced diagram analysis


## Libby-Sterman power counting

- ▶ How to obtain the dominant contribution of an amplitude (in QCD) in a certain specific kinematics (e. g. collinear)?  
⇒ Libby-Sterman power counting rule [Phys.Rev.D 18 (1978) 3252; Phys.Rev.D 18 (1978) 4737]
- ▶ Extensively used in factorisation proofs [Collins: Foundations of perturbative QCD]
- ▶ Basic idea is to identify regions of loop momenta of partons (also number of partons), which gives the dominant contribution to the full amplitude.
- ▶ Collect all contributions to the *smallest*  $\alpha$ :

$$\mathcal{A} = Q^\beta \sum_{\alpha} f_{\alpha} \lambda^{\alpha}, \quad \lambda = \frac{\Lambda_{\text{QCD}}, m_{\pi}, m_N}{Q} \ll 1$$

# Reduced diagram analysis

## Classic Collinear pinch




In both of the above cases, the power counting is [S. N., J. Schönleber, L. Szymanowski, S. Wallon: 2311.09146]:

$$\mathcal{A} \sim Q^{-1} \lambda^\alpha, \quad \lambda = \frac{\Lambda_{\text{QCD}}, m_\pi, m_N}{Q} \ll 1, \quad \alpha = 1$$

# Reduced diagram analysis

## Classic Collinear pinch



In both of the above cases, the power counting is [S. N., J. Schönleber, L. Szymanowski, S. Wallon: 2311.09146]:

$$\mathcal{A} \sim Q^{-1} \lambda^\alpha, \quad \lambda = \frac{\Lambda_{\text{QCD}}, m_\pi, m_N}{Q} \ll 1, \quad \alpha = 1$$

Collinear factorisation at *all orders* and *leading power* provided:

- ▶ the above collinear **pinch** diagrams (standard) are the *only ones contributing to the leading power of  $\alpha = 1$*
- ▶ the *soft factor S 'cancels'*

# Pinches

## Landau conditions

Pinches correspond to regions of loop momentum which cannot be avoided through contour deformations.

They can be identified efficiently through **Landau conditions**:



Pinches correspond to regions of loop momentum which cannot be avoided through contour deformations.

They can be identified efficiently through **Landau conditions**:

$$I(z) = \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}^{dL}} d^{dL}\omega \frac{N(\omega, z)}{\prod_{j=0}^n (D_j(\omega, z) + i\epsilon)}.$$

Given  $z, \omega_S \in \mathbb{R}^{dL}$  such that the set

$$\mathcal{D} = \{j \in \{1, \dots, n\} \mid D_j(\omega_S, z) = 0\}$$

is non-empty, we have a pinch at  $\omega_S$  iff there exist real and non-negative numbers  $\alpha_j$  for  $j \in \mathcal{D}$  such that

- ▶  $\forall i \in \{1, \dots, dL\} : \sum_{j \in \mathcal{D}} \alpha_j \frac{\partial D_j}{\partial \omega_i}(\omega_S; z) = 0.$
- ▶ At least one of the  $\alpha_j$  is non-zero

Pinches correspond to regions of loop momentum which cannot be avoided through contour deformations.

They can be identified efficiently through **Landau conditions**:

$$I(z) = \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}^{dL}} d^{dL}\omega \frac{N(\omega, z)}{\prod_{j=0}^n (D_j(\omega, z) + i\epsilon)}.$$

Given  $z, \omega_S \in \mathbb{R}^{dL}$  such that the set

$$\mathcal{D} = \{j \in \{1, \dots, n\} \mid D_j(\omega_S, z) = 0\}$$

is non-empty, we have a pinch at  $\omega_S$  iff there exist real and non-negative numbers  $\alpha_j$  for  $j \in \mathcal{D}$  such that

- ▶  $\forall i \in \{1, \dots, dL\} : \sum_{j \in \mathcal{D}} \alpha_j \frac{\partial D_j}{\partial \omega_i}(\omega_S; z) = 0.$
- ▶ At least one of the  $\alpha_j$  is non-zero

*Note:* Existence of pinch does *not* imply existence of a singularity: Need to also perform *power counting*.

# Pinches

Soft pinch always present

Consider the bubble integral, with **massless** internal lines:

$$I_1(p^2) = \lim_{\epsilon \rightarrow 0^+} \int d^4k \frac{1}{(k^2 + i\epsilon)((p-k)^2 + i\epsilon)}.$$

# Pinches

Soft pinch always present

Consider the bubble integral, with **massless** internal lines:

$$I_1(p^2) = \lim_{\epsilon \rightarrow 0^+} \int d^4k \frac{1}{(k^2 + i\epsilon)((p-k)^2 + i\epsilon)}.$$

According to the Landau conditions, there is **always** a pinch related to soft momentum  $k$ , independent of  $p$ .

This is because when  $k = 0$ , both the propagator  $k^2 + i\epsilon$  and its first derivative are zero.

$\implies$  *Landau conditions for a pinch at  $k = 0$  are satisfied.*

# Pinches

Soft pinch always present

Consider the bubble integral, with **massless** internal lines:

$$I_1(p^2) = \lim_{\epsilon \rightarrow 0^+} \int d^4k \frac{1}{(k^2 + i\epsilon)((p-k)^2 + i\epsilon)}.$$

According to the Landau conditions, there is **always** a pinch related to soft momentum  $k$ , independent of  $p$ .

This is because when  $k = 0$ , both the propagator  $k^2 + i\epsilon$  and its first derivative are zero.


$\implies$  *Landau conditions for a pinch at  $k = 0$  are satisfied.*

However, note that the power counting does not give an IR divergence for  $p^2 \neq 0$ :

$$\implies \frac{[\lambda^4]}{[\lambda^2][1]} \sim \lambda^2$$

# Reduced diagram analysis

Other leading pinch surfaces?




Divergence obtained when  $(x - \xi) p$  and  $(1 - z) p_M$  lines become soft:

$\implies D_a$  becomes soft and  $D_b$  becomes collinear with respect to  $q$ .

# Reduced diagram analysis

Other leading pinch surfaces?



Divergence obtained when  $(x - \xi)p$  and  $(1 - z)p_M$  lines become soft:


$\implies D_a$  becomes soft and  $D_b$  becomes collinear with respect to  $q$ .

Is there a **leading pinch** diagram that corresponds to this region?

**Yes!**

# Reduced diagram analysis

Other leading pinch surfaces?




$$\mathcal{A} \sim Q^{-1} \lambda^\alpha, \quad \alpha = 1$$



# Reduced diagram analysis

Other leading pinch surfaces?




$$\mathcal{A} \sim Q^{-1} \lambda^\alpha, \quad \alpha = 1$$

$\implies$  power counting is the same as the collinear region!

# Reduced diagram analysis

Other leading pinch surfaces?



$$\mathcal{A} \sim Q^{-1} \lambda^\alpha, \quad \alpha = 1$$

$\implies$  power counting is the same as the collinear region!

*Note: Corresponding reduced diagram for quark GPD case is power suppressed.*

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

- ▶ Need to distinguish between *ultrasoft*, *soft* and *Glauber* gluons:

$$\text{Ultrasoft } k \sim Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\text{Soft } k \sim Q(\lambda, \lambda, \lambda)$$

$$\text{Glauber } k \sim Q(\lambda^2, \lambda^2, \lambda) \quad (\text{or similar with } |k_{\perp}^2| \gg k^+ k^-)$$

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

- ▶ Need to distinguish between *ultrasoft*, *soft* and *Glauber* gluons:

$$\text{Ultrasoft } k \sim Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\text{Soft } k \sim Q(\lambda, \lambda, \lambda)$$

$$\text{Glauber } k \sim Q(\lambda^2, \lambda^2, \lambda) \quad (\text{or similar with } |k_{\perp}^2| \gg k^+ k^-)$$

- ▶ Libby-Sterman power counting formula strictly applies for *ultrasoft gluons* only.

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

- ▶ Need to distinguish between *ultrasoft*, *soft* and *Glauber* gluons:

$$\text{Ultrasoft } k \sim Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\text{Soft } k \sim Q(\lambda, \lambda, \lambda)$$

$$\text{Glauber } k \sim Q(\lambda^2, \lambda^2, \lambda) \quad (\text{or similar with } |k_{\perp}^2| \gg k^+ k^-)$$

- ▶ Libby-Sterman power counting formula strictly applies for *ultrasoft gluons* only.
- ▶ However, these are typically eliminated by the use of *Ward identities*.

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

- ▶ Need to distinguish between *ultrasoft*, *soft* and *Glauber* gluons:

$$\text{Ultrasoft } k \sim Q(\lambda^2, \lambda^2, \lambda^2)$$

$$\text{Soft } k \sim Q(\lambda, \lambda, \lambda)$$

$$\text{Glauber } k \sim Q(\lambda^2, \lambda^2, \lambda) \quad (\text{or similar with } |k_{\perp}^2| \gg k^+ k^-)$$

- ▶ Libby-Sterman power counting formula strictly applies for *ultrasoft gluons* only.
- ▶ However, these are typically eliminated by the use of *Ward identities*.
- ▶ Glauber gluons *cannot* be eliminated/suppressed by the use of Ward identities.

# What exactly does the pinch surface correspond to?

- ▶ Use Sudakov basis  $(+, -, \perp)$ :

$$\text{Collinear } k \sim Q(1, \lambda^2, \lambda) \quad (\text{or } k \sim Q(\lambda^2, 1, \lambda))$$

- ▶ Need to distinguish between *ultrasoft*, *soft* and *Glauber* gluons:

$$\text{Ultrasoft } k \sim Q(\lambda^2, \lambda^2, \lambda^2)$$


$$\text{Soft } k \sim Q(\lambda, \lambda, \lambda)$$

$$\text{Glauber } k \sim Q(\lambda^2, \lambda^2, \lambda) \quad (\text{or similar with } |k_{\perp}^2| \gg k^+ k^-)$$

- ▶ Libby-Sterman power counting formula strictly applies for *ultrasoft gluons* only.
- ▶ However, these are typically eliminated by the use of *Ward identities*.
- ▶ Glauber gluons *cannot* be eliminated/suppressed by the use of Ward identities.
- ▶ Key Question: Is there a *Glauber pinch* that contributes at *leading power*?



# Glauber pinch



(Notation:  $(+, -, \perp)$ )

$$p_N, p_{N'}, \Delta \sim Q(1, \lambda^2, \lambda), \quad \Delta^+ < 0.$$


$$p_\pi \sim Q(\lambda^2, 1, \lambda)$$

$$q, k \sim Q(1, 1, 1), \quad q^2, k^2 \sim \lambda^2 Q^2$$

$$[\text{Loop}] k_g \sim Q(\lambda, \lambda, \lambda)$$

$$[\text{Loop}] r \sim Q(\lambda, \lambda, \lambda)$$

# Glauber pinch



(Notation:  $(+, -, \perp)$ )

$$p_N, p_{N'}, \Delta \sim Q(1, \lambda^2, \lambda), \quad \Delta^+ < 0.$$

$$p_\pi \sim Q(\lambda^2, 1, \lambda)$$


$$q, k \sim Q(1, 1, 1), \quad q^2, k^2 \sim \lambda^2 Q^2$$

$$[\text{Loop}] k_g \sim Q(\lambda, \lambda, \lambda)$$

$$[\text{Loop}] r \sim Q(\lambda, \lambda, \lambda)$$

Recall: Soft loop momenta  $r$  and  $k$  *always* need to be considered.

# Glauber pinch



(Notation:  $(+, -, \perp)$ )

$$p_N, p_{N'}, \Delta \sim Q(1, \lambda^2, \lambda), \quad \Delta^+ < 0.$$

$$p_\pi \sim Q(\lambda^2, 1, \lambda)$$

$$q, k \sim Q(1, 1, 1), \quad q^2, k^2 \sim \lambda^2 Q^2$$

$$[\text{Loop}] k_g \sim Q(\lambda, \lambda, \lambda)$$

$$[\text{Loop}] r \sim Q(\lambda, \lambda, \lambda)$$

Recall: Soft loop momenta  $r$  and  $k$  *always* need to be considered.

►  $k_g^-$  pinch:


$$(k_g - \Delta)^2 + i0 = -2\Delta^+ k_g^- + \mathcal{O}(\lambda^2) + i0$$

$$\implies k_g^- = \mathcal{O}(\lambda^2) - i0.$$

$$(p_{N'} - k_g)^2 + i0 = -2p_{N'}^+ k_g^- + \mathcal{O}(\lambda^2) + i0$$

$$\implies k_g^- = \mathcal{O}(\lambda^2) + i0.$$

# Glauber pinch



$k_g^+$  pinch:


$$(q - r - k_g)^2 + i0 = -2q^+ r^- - 2q^- k_g^+ + \mathcal{O}(\lambda) + i0$$

$$\implies k_g^+ = \mathcal{O}(\lambda) + i0.$$

$$(r + k_g)^2 + i0 = 2k_g^+ r^- + \mathcal{O}(\lambda^2) + i0$$

$$\implies k_g^+ = \mathcal{O}(\lambda) - \text{sgn}(r^-)i0.$$

# Glauber pinch



$k_g^+$  pinch:

$$(q - r - k_g)^2 + i0 = -2q^+ r^- - 2q^- k_g^+ + \mathcal{O}(\lambda) + i0$$

$$\implies k_g^+ = \mathcal{O}(\lambda) + i0.$$


$$(r + k_g)^2 + i0 = 2k_g^+ r^- + \mathcal{O}(\lambda^2) + i0$$

$$\implies k_g^+ = \mathcal{O}(\lambda) - \text{sgn}(r^-)i0.$$

**Conclusion:**  $k_g^+$  is pinched to be  $\mathcal{O}(\lambda)$ , and  $k_g^-$  is pinched to be  $\mathcal{O}(\lambda^2)$ .

$\implies$  **Glauber pinch**, since  $k^+ k^- \ll |k_\perp|^2$ .

# Glauber pinch is leading



Explicit 2-loop analysis shows that the Glauber pinch demonstrated previously is **leading**, i.e. it scales as  $\lambda^\alpha$ , with  $\alpha = 1$ .

# Glauber pinch

## Exclusive double diffractive processes

Very similar to the **exclusive double diffractive process**, where the Glauber gluon is pinched between the two pairs of incoming and outgoing collinear hadrons.


$$p(p_{N_1}) + p(p_{N_2}) \longrightarrow p(p_{N'_1}) + p(p_{N'_2}) + \gamma(k_1) + \gamma(k_2)$$

# Glauber pinch

## Exclusive double diffractive processes

Very similar to the **exclusive double diffractive process**, where the Glauber gluon is pinched between the two pairs of incoming and outgoing collinear hadrons.

$$p(p_{N_1}) + p(p_{N_2}) \longrightarrow p(p_{N'_1}) + p(p_{N'_2}) + \gamma(k_1) + \gamma(k_2)$$



Here, the Glauber pinch corresponds to  $k_g \sim (\lambda^2, \lambda^2, \lambda)$




# Glauber pinch

## Exclusive double diffractive processes

Very similar to the **exclusive double diffractive process**, where the Glauber gluon is pinched between the two pairs of incoming and outgoing collinear hadrons.

$$p(p_{N_1}) + p(p_{N_2}) \longrightarrow p(p_{N'_1}) + p(p_{N'_2}) + \gamma(k_1) + \gamma(k_2)$$



Here, the Glauber pinch corresponds to  $k_g \sim (\lambda^2, \lambda^2, \lambda)$

Instead, in our case, the Glauber gluon (which corresponds to one of the active partons) is pinched between **a pair of collinear hadrons**, and **a soft line joining the outgoing pion and the incoming photon**.

# Conclusions

- ▶ Collinear factorisation for the exclusive  $\pi^0\gamma$  photoproduction *fails* due to *Glauber pinch* in the *gluon exchange channel*.

# Conclusions

- ▶ Collinear factorisation for the exclusive  $\pi^0\gamma$  photoproduction *fails* due to *Glauber pinch* in the *gluon exchange channel*.
- ▶ Direct calculation assuming collinear factorisation *diverges* already at leading order and leading twist.
- ▶ The same thing happens for the exclusive process  $\pi^0 N \rightarrow N\gamma\gamma$ .

# Conclusions

- ▶ Collinear factorisation for the exclusive  $\pi^0\gamma$  photoproduction *fails* due to *Glauber pinch* in the *gluon exchange channel*.
- ▶ Direct calculation assuming collinear factorisation *diverges* already at leading order and leading twist.
- ▶ The same thing happens for the exclusive process  $\pi^0 N \rightarrow N\gamma\gamma$ .
- ▶ Important to check factorisation of such 3-body final state exclusive processes, before implementing them in the virtual access of STRONG2020: PARTONS and NLOAccess.
- ▶ Channels where *2-gluon exchanges are forbidden* ( $\pi^\pm$  and  $\rho^{0,\pm}$ ) are *safe from the effects discussed here*.

# Conclusions

- ▶ Collinear factorisation for the exclusive  $\pi^0\gamma$  photoproduction *fails* due to *Glauber pinch* in the *gluon exchange channel*.
- ▶ Direct calculation assuming collinear factorisation *diverges* already at leading order and leading twist.
- ▶ The same thing happens for the exclusive process  $\pi^0 N \rightarrow N\gamma\gamma$ .
- ▶ Important to check factorisation of such 3-body final state exclusive processes, before implementing them in the virtual access of STRONG2020: PARTONS and NLOAccess.
- ▶ Channels where *2-gluon exchanges are forbidden* ( $\pi^\pm$  and  $\rho^{0,\pm}$ ) are *safe from the effects discussed here*.
- ▶ Factorisation breaking effects also expected to occur in specific channels that allow for *2-gluon exchanges* in exclusive *di-meson photoproduction*:  $\gamma N \rightarrow M_1 M_2 N'$ . [ongoing]

# Conclusions

- ▶ Collinear factorisation for the exclusive  $\pi^0\gamma$  photoproduction *fails* due to *Glauber pinch* in the *gluon exchange channel*.
- ▶ Direct calculation assuming collinear factorisation *diverges* already at leading order and leading twist.
- ▶ The same thing happens for the exclusive process  $\pi^0 N \rightarrow N\gamma\gamma$ .
- ▶ Important to check factorisation of such 3-body final state exclusive processes, before implementing them in the virtual access of STRONG2020: PARTONS and NLOAccess.
- ▶ Channels where *2-gluon exchanges are forbidden* ( $\pi^\pm$  and  $\rho^{0,\pm}$ ) are *safe from the effects discussed here*.
- ▶ Factorisation breaking effects also expected to occur in specific channels that allow for *2-gluon exchanges* in exclusive *di-meson photoproduction*:  $\gamma N \rightarrow M_1 M_2 N'$ . [ongoing]
- ▶ Compute  $\gamma N \rightarrow \gamma\pi^0 N$  in high-energy ( $k_T$ ) factorisation. [ongoing]

## BACKUP SLIDES

# More about pinches

Soft pinch always present

Consider the *triangle* integral, with *massless* internal lines:

$$I_2 = \lim_{\epsilon \rightarrow 0^+} \int d^4 k \frac{1}{(k^2 + i\epsilon)((k - p_1)^2 + i\epsilon)((k + p_2)^2 + i\epsilon)}.$$

Again, Landau conditions predict the existence of a pinch at  $k = 0$ .

If  $p_1^2 = m_1^2$  and  $p_2^2 = m_2^2$ , then the **power counting** predicts a *logarithmic divergence*:

$$\Rightarrow \frac{[\lambda^4]}{[\lambda^2][\lambda][\lambda]} \sim \lambda^0$$

This is of course the well-known **soft singularity** of triangle integrals, where the massless particle connects to two on-shell legs.



# More about pinches

## Collinear pinch

Consider the bubble integral, with *massless* internal lines:

$$I_1(p^2) = \lim_{\epsilon \rightarrow 0^+} \int d^4 k \frac{1}{(k^2 + i\epsilon)((p-k)^2 + i\epsilon)}.$$

We apply the Landau conditions:

$$k^2 = 0, \quad p^2 - 2p \cdot k = 0, \quad \alpha_1 k + \alpha_2 (k - p) = 0$$
$$\alpha_1, \alpha_2 \geq 0, \quad \alpha_1 + \alpha_2 > 0$$

This implies

$$k^2 = 0, \quad p^2 - 2p \cdot k = 0, \quad k = \alpha p,$$


where  $1 \geq \alpha \geq 0$ . This only has a solution if  $p^2 = 0$ . This is of course nothing but the well-known **collinear singularity**.

The **power counting** indicates a *logarithmic divergence*:

$$\implies \frac{[\lambda^4]}{[\lambda^2][\lambda^2]} \sim \lambda^0, \text{ as expected}$$

# Glauber pinch

Non-analyticity in  $r^-$



Start with  $r \sim Q(\lambda_s, \lambda_s, \lambda_s)$ , where  $\lambda_s \ll 1$ , but completely general wrt  $\lambda$ . *Study pole in  $r^+$ :*

$$r^2 + i0 = 2r^+r^- - |r_\perp|^2 + i0,$$

$$\implies r^+ = \mathcal{O}(\lambda_s) - \text{sgn}(r^-) i0.$$


$$(p_\pi - r)^2 + i0 = -2p_\pi^- r^+ + \mathcal{O}(\max(\lambda^2, \lambda_s^2)) + i0,$$

$$\implies r^+ = \mathcal{O}(\max(\lambda^2, \lambda_s^2)) + i0.$$

**Non-analyticity** at  $r^- = 0$ , and  $r^+$  pinched to be  $\mathcal{O}(\lambda_s)$  for  $\lambda_s \geq \lambda^2$ , or  $r^+$  pinched to be  $\mathcal{O}(\lambda^2)$  for  $\lambda_s \leq \lambda^2$

# Factorisation breaking effects in $\pi^0\gamma$ photoproduction

Gluon GPD contributions



$$D_a = ((x - \xi)p + \bar{z}p_M)^2 + i\epsilon$$

$$= s\bar{\alpha}\bar{z}[x - \xi + i\epsilon] ,$$

$$D_b = (k + zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s[z(x - \xi - i\epsilon) + \alpha\bar{z}(x + \xi - i\epsilon)] ,$$

$$D_c = (zp_M - (x + \xi)p)^2 + i\epsilon$$

$$= -s\bar{\alpha}z[x + \xi - i\epsilon]$$

$\implies$  pinching of poles in the propagators ( $D_a$  and  $D_b$ ) in the limit of  $z \rightarrow 1$