

# ALP contribution to the Strong CP problem

*Tuesday, 2 July 2024 16:30 (5 minutes)*

We compute the one-loop contribution to the  $\bar{\theta}$ -parameter of an axion-like particle (ALP) with CP-odd derivative couplings. Its contribution to the neutron electric dipole moment is shown to be orders of magnitude larger than that stemming from the one-loop ALP contributions to the up- and down-quark electric and chromoelectric dipole moments. This strongly improves existing bounds on ALP-fermion CP-odd interactions, and also sets limits on previously unconstrained couplings. The case of a general singlet scalar is analyzed as well. In addition, we explore how the bounds are modified in the presence of a Peccei-Quinn symmetry.

## Title of the Poster/Talk

Improved nEDM limits on ALP couplings to fermions

## Related Papers/Preprints

2403.12133

**Primary authors:** GAVELA, Belen (Universidad Autónoma de Madrid); GRINSTEIN, Benjamin (University of California San Diego); Dr QUILEZ LASANTA, Pablo (University of California San Diego); ENGUIA-VILETA, Víctor (Instituto de Física Teórica (IFT UAM-CSIC))

**Presenter:** ENGUIA-VILETA, Víctor (Instituto de Física Teórica (IFT UAM-CSIC))

**Session Classification:** Young Scientist Forum