A closer look at dark photon production modes

High luminosity colliders and fixed target facilities using proton beams are sensitive to new weakly coupled degrees of freedom across a broad mass range. We discuss various production modes for dark vector particles in proton beam experiments. In particular, we will have a closer look at bremsstrahlung which is important for dark vectors with masses between 0.5 GeV and 1.5 GeV, due to resonant mixing with hadronic resonances. We revisit the calculation of dark photons via initial state radiation in non-single diffractive scattering, using an improved approach to the splitting function and the timelike electromagnetic form-factor. The framework is benchmarked by applying an analogous calculation to model inclusive rho-meson production, indicating agreement with data from NA27 in the relevant kinematic range. In addition, we add dark photon couplings to quarks to the parton shower in Herwig to study the effect of Parton-level dark photon radiation on the overall production rate of dark photons in proton-proton collisions.

Title of the Poster/Talk

Related Papers/Preprints

Primary authors: RITZ, Adam (University of Victoria); MASOUMINIA, Aidin (IPPP, Durham University); REIMITZ, Peter (Instituto de Fisica, USP); FOROUGHI ABARI, Saeid (University of Victoria); PLAETZER, Simon (University of Vienna)

Presenter: REIMITZ, Peter (Instituto de Fisica, USP)

Session Classification: Poster Session