



# Dark Particles at the LHC: LHC-Friendly Dark Matter Characterization via Non-Linear EFT


David Cabo-Almeida

2407.XXXXX

In collaboration with: Giorgio Arcadi, Sven Fabian, Florian Goertz



# EFT & Simplified Models



# Non-linear Approach

- To solve the dichotomy, recently the ‘extended Dark Matter Effective Field Theory’ (eDMeft) framework was proposed (see T.Alanne et al. Eur. Phys. J. C, 80(5):446, 2020 & T.Alanne et al. JHEP, 10:172, 2020)
- Extended the formalism to two scalar mediators.
- With non-linear formalism more representation of scalar mediator can be cover

Goldstone Matrix

$SU(2)_L \times U(1)_Y$

$$\Sigma(x) = e^{i\sigma^a \phi^a(x)/v} \longrightarrow \Sigma \rightarrow e^{i\varphi_L^a(x)\sigma^a/2} \Sigma e^{-i\varphi_Y(x)\sigma^3/2}$$

$$D_\mu \Sigma = \partial_\mu \Sigma - ig \frac{\sigma^a}{2} W_\mu^a \Sigma + ig' \Sigma \frac{\sigma^3}{2} B_\mu$$

We work adding under this formalism 3 new fields

$\chi, S_1, S_2$

# Extended Dark Matter EFT

$$\mathcal{O}_d^C(h, \mathcal{S}_1, \mathcal{S}_2) \equiv \sum_{k=0}^d \sum_{j=0}^{d-k} \sum_{i=0}^j C_{i,j-i}^{(k)} h^k \mathcal{S}_1^i \mathcal{S}_2^{j-i}$$

$\mathcal{L} = \mathcal{L}_{\text{gauge-ferm}}^{\text{SM}} + \bar{\chi} i \not{D} \chi - \mathcal{O}_2^y(h, \mathcal{S}_1, \mathcal{S}_2) \bar{\chi}_L \chi_R + \text{h.c.} \xrightarrow{\text{DM coupling to scalars}}$

$+ \mathcal{O}_5^\lambda(h, \mathcal{S}_1, \mathcal{S}_2) \xrightarrow{\text{All Scalar interactions}}$

$+ \frac{v^2}{4} \text{Tr} \left[ (D_\mu \Sigma)^\dagger (D^\mu \Sigma) \right] \mathcal{O}_3^\kappa(h, \mathcal{S}_1, \mathcal{S}_2) \xrightarrow{W^+ W^- \phi, ZZ\phi\dots}$





$- \sum_\phi \frac{\phi}{16\pi^2} \left[ g^2 c_B^\phi B^{\mu\nu} B_{\mu\nu} + g^2 c_W^\phi W^{I\mu\nu} W_I^{\mu\nu} + g_s^2 c_G^\phi G^{a\mu\nu} G_a^{\mu\nu} \right] \xrightarrow{GG\phi, W^+ W^- \phi}$

$+ i \frac{v^2}{4} \text{Tr} \left[ \Sigma^\dagger (D^\mu \Sigma) \sigma^3 \right] \left( \partial_\mu h \mathcal{O}_2^s(h, \mathcal{S}_1, \mathcal{S}_2) + \partial_\mu \mathcal{S}_1 \mathcal{O}_2^{s1}(h, \mathcal{S}_1, \mathcal{S}_2) + \partial_\mu \mathcal{S}_2 \mathcal{O}_2^{s2}(h, \mathcal{S}_1, \mathcal{S}_2) \right) \xrightarrow{Z\phi_1\phi_2\dots\phi_m}$



$- \frac{v}{\sqrt{2}} \left( \begin{pmatrix} \bar{u}_L^{(i)} & \bar{d}_L^{(i)} \end{pmatrix} \Sigma \begin{pmatrix} Y_{ij}^u u_R^{(j)} \\ Y_{ij}^d d_R^{(j)} \end{pmatrix} \mathcal{O}_2^{c_q}(h, \mathcal{S}_1, \mathcal{S}_2) + \bar{\ell}_L^{(i)} \Sigma Y_{ij}^\ell \ell_R^{(j)} \mathcal{O}_2^{c_\ell}(h, \mathcal{S}_1, \mathcal{S}_2) + \text{h.c.} \right) \xrightarrow{\bar{f}f\phi}$

# Possible signatures


## Collider Signatures




## EWPO



## Direct Detection



## Relic Density



# Toy Model

$$-\mathcal{L} \supset \mu_\Phi^2 |\Phi|^2 + \lambda_\Phi |\Phi|^4 + \mu_S^2 |S|^2 + \lambda_S |S|^4 + \lambda_{\Phi S} |\Phi|^2 |S|^2 + \mu_a^2 a^2$$

$$-\mathcal{L} \supset y_\chi s \bar{\chi}_L S \chi_R + \sum_{\mathcal{Q}} y_{\mathcal{Q} S} \bar{\mathcal{Q}}_L S \mathcal{Q}_R + \text{h.c.}$$

**With heavy chiral quarks**  $\mathcal{Q} = \mathcal{B}, \mathcal{T}$

Field	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$	global $U(1)$
$S$	<b>1</b>	<b>1</b>	0	+1
$\mathcal{T}_L$	<b>3</b>	<b>1</b>	4/3	+1/2
$\mathcal{T}_R$	<b>3</b>	<b>1</b>	4/3	-1/2
$\mathcal{B}_L$	<b>3</b>	<b>1</b>	4/3	+1/2
$\mathcal{B}_R$	<b>3</b>	<b>1</b>	4/3	-1/2
$\chi_L$	<b>1</b>	<b>1</b>	0	+1/2
$\chi_R$	<b>1</b>	<b>1</b>	0	-1/2


$$\Phi = \begin{pmatrix} G^+ \\ (\nu_h + h + iG^0)/\sqrt{2} \end{pmatrix}, \quad S = \frac{1}{\sqrt{2}} (\nu_s + s + ia)$$

$$\mathcal{M}_s = \begin{pmatrix} 2\nu_h^2 \lambda_\Phi & \times \\ \lambda_{\Phi S} \nu_h \nu_s & 2\lambda_S \nu_s^2 \end{pmatrix}$$

# Toy Model

$$-\mathcal{L} \supset \mu_\Phi^2 \Phi^2 + \lambda_\Phi \Phi^4 + \mu_S^2 S^2 + \lambda_S S^4 + \lambda_{\Phi S} \Phi^2 S^2 + \mu_a^2 a^2$$

$$-\mathcal{L} \supset y_\chi s \bar{\chi}_L S \chi_R + \sum_Q y_{Q S} \bar{Q}_L S Q_R + \text{h.c.}$$



D.D. completely match


Can mimic mono-h and mono-Z

# Toy Model


$$-\mathcal{L} \supset \mu_\Phi^2 \Phi^2 + \lambda_\Phi \Phi^4 + \mu_S^2 S^2 + \lambda_S S^4 + \lambda_{\Phi S} \Phi^2 S^2 + \mu_a^2 a^2$$

$$-\mathcal{L} \supset y_\chi s \bar{\chi}_L S \chi_R + \sum_Q y_{QS} \bar{Q}_L S Q_R + \text{h.c.}$$

**Gluon fusion in  $D = 5$  operator**





# Relic Density



# EFT vs UV at collider


$$\chi \equiv (\sigma_{pp}^{\text{EFT}} - \sigma_{pp}^{\text{UV}}) / \sigma_{pp}^{\text{UV}}$$



# A simplified model

## 2HDM+Pseudoscalar

- Good compromise between theoretical consistency and predictivity (still limited number of free parameters);
- Benchmark for a large variety of collider studies;
- Interesting Dark Matter phenomenology.



LHC Dark Matter Working Group: Phys. Dark. Univ. 27 (2020) 100351

(see also e.g. M. Bauer et al. JHEP 05 (2017) 138, T. Robens Symmetry 13 (2021) 12, 2341)

# 2HDM+P


$Z_2$  symmetry for 2HDM potential

$$V_{2HDM} = m_1^2 \phi_1^\dagger \phi_1 + m_2^2 m_1^2 \phi_2^\dagger \phi_2 - m_3^2 (\phi_1^\dagger \phi_2 + h.c.) + \frac{1}{2} \lambda_1 (\phi_1^\dagger \phi_1)^2 + \frac{1}{2} \lambda_2 (\phi_2^\dagger \phi_2)^2 + \frac{1}{2} \lambda_5 ((\phi_1^\dagger \phi_2)^2 + h.c.) \\ + \lambda_3 (\phi_1^\dagger \phi_1) (\phi_2^\dagger \phi_2) + \lambda_4 (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_1)$$


$$V(\Phi_1, \Phi_2, a_0) = V_{2HDM}(\phi_1, \phi_2) + V_{self}(a_0) + V_{a_0, 2HDM}(\phi_1, \phi_2, a_0)$$

$$\Phi_j = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} \hat{\phi}_j^+ \\ v_j + \hat{\rho}_j + i \hat{\eta}_j \end{pmatrix}$$

with  $j = 1, 2$



$$V_{self}(a_0) = \frac{1}{2} m_{a_0}^2 a_0^2 + \frac{1}{4} \lambda_a a_0^4$$



$$V_{a_0, 2HDM}(\phi_1, \phi_2, a_0) = \kappa (i a_0 \phi_1^\dagger \phi_2 + h.c.) + \lambda_{1P} a_0^2 \phi_1^\dagger \phi_1 + \lambda_{2P} a_0^2 \phi_2^\dagger \phi_2$$

# Mixing and EW Symmetry Breaking

$$\langle \phi_1 \rangle = v_1$$

$$\frac{v_2}{v_1} = \tan \beta$$

$$\langle \phi_2 \rangle = v_2$$

$$(\phi_1, \phi_2, a_0) \longrightarrow (h, a, H, A, H^\pm)$$

## Mixing and Yukawa sector

$$\begin{pmatrix} A^0 \\ a^0 \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} A \\ a \end{pmatrix} \longrightarrow L_{Yuk} = \sum_f \frac{m_f}{v} [g_{Hff} H \bar{f} f + g_{hff} h \bar{f} f - i g_{aff} a \bar{f} \gamma_5 f - i g_{Aaff} A \bar{f} \gamma_5 f]$$

	Type I	Type II	Type X	Type Y
$g_{htt}$	$\frac{\cos \alpha}{\sin \beta} \rightarrow 1$			
$g_{hbb}$	$\frac{\cos \alpha}{\sin \beta} \rightarrow 1$	$-\frac{\sin \alpha}{\cos \beta} \rightarrow 1$	$\frac{\cos \alpha}{\sin \beta} \rightarrow 1$	$-\frac{\sin \alpha}{\cos \beta} \rightarrow 1$
$g_{h\tau\tau}$	$\frac{\cos \alpha}{\sin \beta} \rightarrow 1$	$-\frac{\sin \alpha}{\cos \beta} \rightarrow 1$	$-\frac{\sin \alpha}{\cos \beta} \rightarrow 1$	$\frac{\cos \alpha}{\sin \beta} \rightarrow 1$
$g_{Htt}$	$\frac{\sin \alpha}{\sin \beta} \rightarrow -\frac{1}{\tan \beta}$			
$g_{Hbb}$	$\frac{\sin \alpha}{\sin \beta} \rightarrow -\frac{1}{\tan \beta}$	$\frac{\cos \alpha}{\cos \beta} \rightarrow \tan \beta$	$\frac{\sin \alpha}{\sin \beta} \rightarrow -\frac{1}{\tan \beta}$	$\frac{\cos \alpha}{\cos \beta} \rightarrow \tan \beta$
$g_{H\tau\tau}$	$\frac{\sin \alpha}{\sin \beta} \rightarrow -\frac{1}{\tan \beta}$	$\frac{\cos \alpha}{\cos \beta} \rightarrow \tan \beta$	$\frac{\cos \alpha}{\cos \beta} \rightarrow \tan \beta$	$\frac{\sin \alpha}{\sin \beta} \rightarrow -\frac{1}{\tan \beta}$
$g_{A^0tt}$	$\frac{1}{\tan \beta}$	$\frac{1}{\tan \beta}$	$\frac{1}{\tan \beta}$	$\frac{1}{\tan \beta}$
$g_{A^0bb}$	$-\frac{1}{\tan \beta}$	$\tan \beta$	$-\frac{1}{\tan \beta}$	$\tan \beta$
$g_{A^0\tau\tau}$	$-\frac{1}{\tan \beta}$	$\tan \beta$	$\tan \beta$	$-\frac{1}{\tan \beta}$

$$g_{Aff} = \cos \theta g_{A^0ff}$$

$$g_{aff} = \sin \theta g_{A^0ff}$$


## Alignment limit

$$\alpha \rightarrow \beta - \frac{\pi}{2}$$


# 2HDM+P Relic Density

Better results for double pseudoscalar setup


$$S_1 = A \quad S_2 = a$$



$$S_1 = H \quad S_2 = a$$





# Direct Detection



$S_1 = A \quad S_2 = a$

$$\chi \equiv (\sigma_{\chi-n}^{\text{EFT}} - \sigma_{\chi-n}^{\text{UV}}) / \sigma_{\chi-n}^{\text{UV}}$$

$S_1 = H \quad S_2 = a$



# Maximum Gap

## EWPO

$$\Delta\rho = \frac{\alpha_{\text{QED}}(m_Z^2)}{16\pi^2 m_W^2 (1 - m_W^2/m_Z^2)} \left[ f(m_{H^\pm}^2, m_H^2) + c_\theta^2 (f(m_{H^\pm}^2, m_A^2) - f(m_A^2, m_H^2)) + s_\theta^2 (f(m_{H^\pm}^2, m_P^2) - f(m_P^2, m_H^2)) \right]$$


## Invisible Higgs

$$\Gamma_h^{\text{BSM}} = \sum_{\phi_1, \phi_2} \Gamma(h \rightarrow \phi_1 \phi_2)$$


## Perturbative unitarity & Vacuum stability

See e.g. G. Arcadi et al. Phys. Rev. D, 108(5):055010, 2023 for more detail

$$S_1 = A \quad S_2 = a$$





$$S_1 = H \quad S_2 = a$$



# 2HDM+P Mono-Higgs

$$\chi \equiv (\sigma_{pp}^{\text{EFT}} - \sigma_{pp}^{\text{UV}}) / \sigma_{pp}^{\text{UV}}$$



# Conclusions

- We extend eDMEFT to capture more LHC signatures.
- We compare eDMEFT with simplified models to show the validity.
- Final Steps:
  - Can we detect if our singlets come from a multiplet?

# Back Up

# 2HDM+P Possible Mass Gap

