

# Search for dark matter decay and annihilation using observation by Tibet AS $_{\gamma}$ and LHAASO

*giovedì 11 luglio 2024 14:40 (20 minuti)*

Recently, Tibet AS $_{\gamma}$  and LHAASO have observed very high energy diffuse gamma rays in the Galactic plane between 10 TeV and 1 PeV energies. In our work, we utilize these observations to search for dark matter decay or annihilation signals to Standard Model particles. In addition to the primary gamma-ray originating from various Standard Model particles, we also include secondary gamma-rays generated in these processes. We also consider the effects of dark matter substructures and tidal disruption. We place constraints on dark matter annihilation cross-section and decay lifetime for a wide range of dark matter masses. Future observation of these high-energy gamma rays can further help us either discover particle dark matter or better constrain its properties.

**Autori principali:** DUBEY, Abhishek (Indian institute of science, Bangalore); KUMAR SAHA, Akash (Centre for High Energy Physics (CHEP), Indian Institute of Science, Bengaluru, India); LAHA, Ranjan; NATH MAITY, Tarak

**Relatore:** DUBEY, Abhishek (Indian institute of science, Bangalore)

**Classifica Sessioni:** Parallel 2

**Classificazione della track:** Parallel session: Indirect detection