

# Nucleon decays at JUNO

**Wanlei Guo (IHEP)**

guowl@ihep.ac.cn

On behalf of the JUNO collaboration


NOW 2024, Otranto, Sep 06, 2024



- 1. Status of nucleon decay searches**
- 2. JUNO sensitivity on  $p \rightarrow \bar{\nu} K^+$**
- 3. Neutron invisible decays in JUNO**
- 4. Summary**



## Experimental side: cosmological matter-antimatter asymmetry




**Sakharov's three ingredients:** JETP Letters 5,24 (1967)

1. *Baryon Number Violation*
2. *C and CP Violation*
3. *Non-Equilibrium Conditions*

**Lightest Baryon:**  
Proton  
*Stability?*

*B conservation from accidental global symmetry in SM*



## Theoretical side: Grand Unified Theories (GUTs) Phys. Rept. 441, 191 (2007)



A natural consequence



**Nucleon decay**



Searching for nucleon decays plays a key role to understand baryon asymmetry and test GUTs



# Search history of nucleon decays




In 1974, Georgi and Glashow propose the minimal SU(5) GUT  $\rightarrow \tau_p \approx 10^{28} - 10^{32}$  yrs

PRL 32,438 (1974)

Snowmass: 2203.08771


$\rightarrow$  Detectors with about 1000 ton mass can test the SU(5) GUT

## The first generation of experiments (80s):



## The second generation (90s):

### Super-Kamiokande



After 2000, some neutrino experiments, Such as KamLAND, SNO and BOREXINO, have also searched for nucleon decays.

All these experiments don't find the evidence for nucleon decay, excluded minimal SU(5) 3



# Current limits on nucleon decays



Mode	Partial mean life ( $10^{30}$ years)	90% CL
<b>Antilepton + meson</b>		
$\tau_1$ $N \rightarrow e^+ \pi$	$B$	$> 5300$ (n), $> 24000$ (p) Super-K
$\tau_2$ $N \rightarrow \mu^+ \pi$	$B$	$> 3500$ (n), $> 16000$ (p)
$\tau_3$ $N \rightarrow \nu \pi$		$> 1100$ (n), $> 390$ (p)
$\tau_4$ $p \rightarrow e^+ \eta$		$> 10000$
$\tau_5$ $p \rightarrow \mu^+ \eta$		$> 4700$
$\tau_6$ $n \rightarrow \nu \eta$		$> 158$
$\tau_7$ $N \rightarrow e^+ \rho$		$> 217$ (n), $> 720$ (p)
$\tau_8$ $N \rightarrow \mu^+ \rho$		$> 228$ (n), $> 570$ (p)
$\tau_9$ $N \rightarrow \nu \rho$		$> 19$ (n), $> 162$ (p)
$\tau_{10}$ $p \rightarrow e^+ \omega$		$> 1600$
$\tau_{11}$ $p \rightarrow \mu^+ \omega$		$> 2800$
$\tau_{12}$ $n \rightarrow \nu \omega$		$> 108$
$\tau_{13}$ $N \rightarrow e^+ K$	$B$	$> 17$ (n), $> 1000$ (p)
$\tau_{14}$ $p \rightarrow e^+ K_S^0$		
$\tau_{15}$ $p \rightarrow e^+ K_L^0$		
$\tau_{16}$ $N \rightarrow \mu^+ K$	$B$	$> 26$ (n), $> 4500$ (p)
$\tau_{17}$ $p \rightarrow \mu^+ K_S^0$		
$\tau_{18}$ $p \rightarrow \mu^+ K_L^0$		
$\tau_{19}$ $N \rightarrow \nu K$	$N$	$> 86$ (n), $> 5900$ (p) Super-K
$\tau_{20}$ $n \rightarrow \nu K_S^0$		$> 260$
$\tau_{21}$ $p \rightarrow e^+ K^*(892)^0$		$> 84$
$\tau_{22}$ $N \rightarrow \nu K^*(892)$		$> 78$ (n), $> 51$ (p)

## Other modes include:

- ✓ Antilepton + mesons
- ✓ Lepton + meson
- ✓ Lepton + mesons
- ✓ Antilepton + photon(s)
- ✓ Antilepton + single massless
- ✓ Three (or more) leptons
- ✓ Inclusive modes
- ✓  $\Delta B = 2$  dinucleon modes

<https://pdglive.lbl.gov/Particle.action?init=0&node=S016&home=BXXX005#decays>

## Final state particles:

**Mesons:**  $\pi^\pm, \pi^0, K^\pm, K^0, \eta, \rho, \omega, K^*$  (892);

**Leptons:**  $e^\pm, \mu^\pm, \nu$ ; **Photon:**  $\gamma$




**Total measured modes: 82**

*Super-K analyzed 38 modes!*






# Future nucleon decay experiments



	Hyper-K	DUNE	JUNO
Mass (kton)	258 (186)	4*17 (4*10)	20
Target Nucleus	H2O	Ar40	12% H, 88% C12
Technology	Water Cerenkov	LAr TPC	Liquid Scintillator
Start Time	2027	2028/29	2025
Advantages	Large mass and cheap Good particle Identification Good direction resolution	Excellent track reconstruction Excellent particle Identification Good energy resolution	Excellent energy resolution 3% Excellent E threshold 0.7MeV
Shortcomings	Cerenkov threshold	Complex FSI for Ar40	Direction information lost



# Future sensitives on two favored decay modes






## (2) JUNO sensitivity on $p \rightarrow \bar{\nu} K^+$



# Plenary talk from  
Andrea Serafini








# $p \rightarrow \bar{\nu} K^+$ in free and bound protons



**20 kton LS:** Free proton:  $1.45 \times 10^{33}$   
Bound proton:  $5.30 \times 10^{33}$


## Kinetic energy of $K^+$

Free proton  $\rightarrow$  105 MeV  
Bound proton:  $\downarrow$



## Nuclear Effects:

1. Binding Effect
2. Fermi Motion
3. NN correlation



5. De-excitation of remaining nuclei could emit  $\gamma, p, n, \dots$

- **Modify GENIE generator**
- **Implement de-excitation with TALYS**

H. Hu, **W.L. Guo** et al, PLB 831, 137183(2022)




# Signal characters of $p \rightarrow \bar{\nu} K^+$ in JUNO




## Triple coincident signals :


Decay mode	Branching ratio (%)	Kinetic energy sum (MeV)
$K^+ \rightarrow \mu^+ \nu_\mu$	$63.55 \pm 0.11$	152
$K^+ \rightarrow \pi^+ \pi^0$	$20.66 \pm 0.08$	354
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	$5.59 \pm 0.04$	75
$K^+ \rightarrow \pi^0 e^+ \nu_e$	$5.07 \pm 0.04$	265-493
$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$	$3.353 \pm 0.034$	200-388
$K^+ \rightarrow \pi^+ \pi^0 \pi^0$	$1.761 \pm 0.022$	354






First pulse:  $K^+$  kinetic energy of  $\sim 105$  MeV, decay at rest




15 cm, 1.2ns




152 MeV ( $\mu^+$ ) or 354 MeV ( $\pi^+, \pi^0$ )



Third pulse: Michel  $e^+$



AN and PD candidates  $E_{\text{vis}}$  Distribution





# Backgrounds



1MeV

10MeV

100MeV

1GeV

IBD

Proton Decay

Atmospheric neutrinos ~30k in 10 years.

Cosmic Muon

Type	Ratio (%)	Ratio with $E_{vis}$ in [100 MeV, 600 MeV](%)	Interaction	Signal characteristics
NCES	20.2	15.8	$\nu + n \rightarrow \nu + n$ $\nu + p \rightarrow \nu + p$	Single Pulse
CCQE	45.2	64.2	$\nu_l + p \rightarrow n + l^+$ $\nu_l + n \rightarrow p + l^-$	Single Pulse
Pion Production	33.5	19.8	$\nu_l + p \rightarrow l^- + p + \pi^+$ $\nu + p \rightarrow \nu + n + \pi^+$	Approximate Single Pulse (Second pulse too low)
Kaon Production	1.1	0.2	$\nu_l + n \rightarrow l^- + \Lambda + K^+$ $\nu_l + p \rightarrow l^- + p + K^+$	Double Pulse

- If energetic neutrons do not lost most of the energy within ~10 ns
- Kaon Production has a negligible contribution!







# Event Selection



## Three chains based on delayed signals:



- ✓ Michel electrons → MTag
- ✓ Captured neutrons → CTag







# Multi-pulse fitting



## Efficiency vs background


Criteria		Survival rate of $p \rightarrow \bar{\nu}K^+$ (%)			Survival count (fraction) of atmospheric $\nu$		
		Sample 1	Sample 2	Sample 3	Sample 1	Sample 2	Sample 3
basic selection	$E_{vis}$	94.6			51299 (32.1%)		
	$R_V$	93.7			47849 (29.9%)		
Delayed signal selection	$N_M$	74.4		4.4	20739 (13.0%)		1143 (0.7%)
	$\Delta L_M$	67.0		4.4	13796 (8.6%)		994 (0.6%)
	$N_n$	48.4	17.9	—	5403 (3.4%)	6857 (4.3%)	—
	$\Delta L_n$	—	16.6	—	—	4472 (2.8%)	—
Time character selection	$R_\chi$	45.9	9.0	3.8	4326 (2.7%)	581 (0.4%)	716 (0.4%)
	$\Delta T$	28.3	7.7	2.4	121 (0.07%)	18 (0.01%)	30 (0.02%)
	$E_1, E_2$	27.4	7.3	2.2	1 (0.0006%)	0	0
Total		36.9			1		

## Efficiency uncertainties:


Source	Uncertainty
Statistic	1.6%
Position reconstruction	1.7%
Nuclear model	6.8%
Energy deposition model	11.1%
Total	13.2%




# Sensitivity to $p \rightarrow \bar{\nu} K^+$



(a)  $p \rightarrow \bar{\nu} K^+$



(b) atmospheric  $\nu$



**Background:** 0.2/10years  
**Efficiency :** 36.9%



$\tau/B(p \rightarrow \bar{\nu} K^+) > 0.96 \times 10^{34}$  yrs

$n \rightarrow \mu^- K^+$ ,  $p \rightarrow e^+ K^*(892)^0$ ,  $n \rightarrow \nu K^*(892)^0$ , and  $p \rightarrow \nu K^*(892)^+$



# (3) Neutron invisible decays in JUNO





## Bound neutrons in $^{12}\text{C}$ :

- $n \rightarrow inv$  ( $^{12}\text{C} \rightarrow ^{11}\text{C}^*$ )
- $nn \rightarrow inv$  ( $^{12}\text{C} \rightarrow ^{10}\text{C}^*$ )

**Invisible particle:**

neutrinos, NP particles

**Detect de-excitation products of  $^{11}\text{C}^*$  and  $^{10}\text{C}^*$**



## Triple coincident signals :

$^{11}\text{C}^* \rightarrow n +$	$^{10}\text{C}$	( $Br_{n1} = 3.0\%$ )
$^{11}\text{C}^* \rightarrow n + \gamma +$	$^{10}\text{C}$	( $Br_{n2} = 2.8\%$ )
$^{10}\text{C}^* \rightarrow n +$	$^9\text{C}$	( $Br_{nn1} = 6.2\%$ )
$^{10}\text{C}^* \rightarrow n + p +$	$^8\text{B}$	( $Br_{nn2} = 6.0\%$ )

## Half-life Q value

[19.3 s, 3.65 MeV]
[19.3 s, 3.65 MeV]
[0.13 s, 16.5 MeV]
[0.77 s, 18.0 MeV]

Y. Kamyshev and E. Kolbe, PRD 67, 076007 (2003)



## Five background sources:

1. Reactor neutrinos; 2. Natural radioactivity; 3. Long-lived isotopes;
4. Fast neutrons; 5. Atmospheric neutrinos

## Background combinations:

### ➤ Single signal



- *Natural radioactivity*
- *Long-lived isotopes*



**Single+Single+Single**

### ➤ Double signal


- *IBD from reactor neutrinos*
- *He8/Li9 from long-lived isotopes*
- *Fast neutrons*
- *Alpha-N from radioactivity*
- *Atmospheric neutrino NC*



**Double+Single**

### ➤ Triple signal

- *Atmospheric neutrino NC*



**Triple**





# Event selection




Selection Criterion	$n \rightarrow inv$		$nn \rightarrow inv$	
	$^{11}\text{C}^* \rightarrow n + ^{10}\text{C}$	$^{11}\text{C}^* \rightarrow n + \gamma + ^{10}\text{C}$	$^{10}\text{C}^* \rightarrow n + ^9\text{C}$	$^{10}\text{C}^* \rightarrow n + p + ^8\text{B}$
All triple signals	100	100	100	100
Muon Veto	$65.7 \pm 0.2$	$65.5 \pm 0.2$	$80.8 \pm 0.2$	$78.3 \pm 0.2$
Fiducial Volume	$83.5 \pm 0.4$	$82.7 \pm 0.4$	$82.9 \pm 0.4$	$83.1 \pm 0.4$
Event Selection	$75.4 \pm 0.9$	$89.7 \pm 0.3$	$89.2 \pm 0.3$	$83.5 \pm 0.3$
Multiplicity Cut	$93.8 \pm 0.1$	$93.8 \pm 0.1$	$99.9 \pm \mathcal{O}(10^{-4})$	$99.9 \pm \mathcal{O}(10^{-4})$
<b>Combined Selection</b>	$38.8 \pm 0.5$	$45.6 \pm 0.3$	$59.7 \pm 0.4$	$54.3 \pm 0.4$

## FV cut and selection criteria:

Quantity	$n \rightarrow inv$	$nn \rightarrow inv$
$R_{1,2,3}$ [m]	$< 16.7$	$< 16.7$
$E_1$ [MeV]	0.7-12	0.7-30
$E_2$ [MeV]	1.9-2.5	1.9-2.5
$E_3$ [MeV]	1.5-3.5	3.0-16.0
$\Delta T_{12}$ [ms]	$< 1$	$< 1$
$\Delta T_{23}$ [s]	0.002-100	0.002-3.0
$\Delta R_{12}$ [m]	$< 1.5$	$< 1.5$
$\Delta R_{23}$ [m]	$< 1.5$	$< 1.5$
$\Delta R_{13}$ [m]	$< 1.0$	$< 1.0$

## Muon veto strategy:






# $n \rightarrow inv$ signal vs backgrounds



**Dominant BKGs of  $n \rightarrow inv$ : IBD + Single (1235), Atm- $\nu$  NC (3.0) per 10 years**







# $nn \rightarrow inv$ signal vs backgrounds





Dominant BKGs of  $nn \rightarrow inv$ : IBD + Single (3.0), Atm- $\nu$  NC (4.3) per 10 years





# Summary of Backgrounds and Signal efficiency




Backgrounds (10 years)	$n \rightarrow inv$		$nn \rightarrow inv$	
	Basic selection	PSD + MVA	Basic selection	PSD + MVA
IBD + Single	$1235 \pm 50$	$2.72 \pm 0.10$	$3.01 \pm 0.09$	$0.0110 \pm 0.0003$
Atm- $\nu$ NC	$3.0 \pm 1.1$	$0.93 \pm 0.67$	$4.3 \pm 3.5$	$0.55 \pm 0.63$
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ + Single	$3.4 \pm 1.4$	$0.036 \pm 0.013$	–	–
$^9\text{Li}/^8\text{He}$ + Single	$1.55 \pm 0.39$	$0.29 \pm 0.17$	$0.13 \pm 0.13$	$0.13 \pm 0.13$
Accidental	$1.46 \pm 0.05$	$0.095 \pm 0.004$	–	–
<b>Total</b>	<b><math>1244 \pm 50</math></b>	<b><math>4.07 \pm 0.68</math></b>	<b><math>7.4 \pm 3.5</math></b>	<b><math>0.69 \pm 0.64</math></b>
Signal efficiency (%)	$n \rightarrow inv$		$nn \rightarrow inv$	
	Basic selection	PSD + MVA	Basic selection	PSD + MVA
$\epsilon_{n(nn)1}$	$35.6 \pm 0.2$	$23.5 \pm 0.2$	$54.0 \pm 0.3$	$48.2 \pm 0.3$
$\epsilon_{n(nn)2}$	$43.6 \pm 0.3$	$30.3 \pm 0.3$	$49.2 \pm 0.3$	$36.3 \pm 0.3$





# JUNO sensitivity



An order of magnitude improvement to the current best limits in 2 years data taking 20



## ◆ JUNO is a large multi-purpose LS detector

- ✓ *Large mass (20 kton) → Free p:  $1.43 \times 10^{33}$ ; Bound p/n:  $5.30 \times 10^{33}$*
- ✓ *Excellent energy resolution → 2.95% at 1 MeV*
- ✓ *Low threshold (0.2 MeV) → Identify neutrons and residual nuclei*

## ◆ Competitive sensitivities for some nucleon decay modes

- ✓  $\tau/B(p \rightarrow \bar{\nu} K^+) > 0.96 \times 10^{34}$  yrs
- ✓  $\tau/B(n \rightarrow inv) > 5.0 \times 10^{31}$  yrs **90% CL**
- ✓  $\tau/B(nn \rightarrow inv) > 1.4 \times 10^{32}$  yrs

## ◆ Continually improve physical analyses for advantaged modes

## ◆ Search for other potential decay modes and relevant new physics



Keep digging new physics !

*Thanks for your attention!*





# Backup: Signal Characteristics in LS

