

Beta decay and neutrino mass: **KATRIN** and beyond




Neutrino Oscillation Workshop
NOW 2024, Otranto, Italia

*Magnus Schlösser
Karlsruhe Institute of Technology
Institute for Astroparticle Physics
Tritium Laboratory Karlsruhe*



The role of massive neutrinos and motivations to measure it

Neutrino masses bring in a **fundamental energy scale** (besides Higgs scale)



Cosmology and the role of neutrinos therein **may be more complex** (what is DE, ...?)



Model for mass generation needs:
mixing matrix **AND mass scale**

$0\nu\beta\beta$ observation **not necessarily** points to an **neutrino mass**

Signal is „in reach“: Minimal mass scales exist!

“ $m(\nu_e)$ “ > 10 meV (normal mass ordering)
“ $m(\nu_e)$ “ > 50 meV (inverted mass ordering)


Ways to access the neutrino mass



	Cosmology	Search for $0\nu\beta\beta$	β -decay & electron capture
Observable	$M_\nu = \sum_i m_i$	$m_{\beta\beta}^2 = \left \sum_i U_{ei}^2 m_i \right ^2$	$m_\beta^2 = \sum_i U_{ei} ^2 m_i^2$
Present upper limit	0.12 eV (0.072 eV)	0.156 eV	0.8 eV
Model dependence	Multi-parameter cosmological model	<ul style="list-style-type: none"> - Majorana ν - contributions other than $m(\nu)$? - nuclear matrix elements, g_A 	Direct, only kinematics; no cancellations in incoherent sum

Complementarity and need for direct mass measurements

Standard neutrino picture: **observations have to be found in colored regions**



Tie-breaker needed to exclude exotic models in neutrino nature or cosmology

KATRIN, Nat. Phys. **18** (2022) 160

KamLAND-Zen, PRL **130**, 051801 (2023)

Planck, Astron. Astrophys. **641** (2020) A6


DESI, 2406.14554

Direct mass experiments

- Direct, model-independent access to neutrino mass


$\beta^{(-)}$ decay

$$m_\beta^2 = \sum_i |U_{ei}|^2 m_i^2$$




Measurement of kinetic energy of electron

Electron capture




Measurement of internal excitation of daughter atom

(Anti-) neutrino mass determined from shape distortion near kinematic endpoint



Challenges for achieving mass low sensitivity



Low kinematic endpoint, high decay rate

	β -decay	Electron capture
Chosen isotope	${}^3\text{H} = \text{T}$	${}^{163}\text{Ho}$
Endpoint	18.6 keV	2.8 keV
Half life	12.3 years	4570 years
Typ. production	n-capture in D_2O	n-irradiation of ${}^{162}\text{Er}$

This talk

High signal (\rightarrow statistics)
Low background (\rightarrow statistics)


High energy resolution (\rightarrow sensitivity)
Low and quantified systematic effects

See Talk
06 Sep, 17:50


Elena Ferri

Tritium beta decay experiments


- Direct, model-independent access to neutrino mass



$$m_\beta^2 = \sum_i |U_{ei}|^2 m_i^2$$




$$\frac{dN}{dE} \propto \sqrt{(E_0 - E)^2 - m_{\bar{\nu}_i}^2 c^4}$$




KATRIN's aim: Measurement of $m_{\bar{\nu}}$ with a sensitivity of $0.2 \text{ eV}/c^2$

Molecular decay


Atomic decay



Decay from a molecule



Final-state distribution



Molecular effects need to be taken into account in neutrino mass analysis

„model-dependence“

Established measurement principles

CRES

- Cyclotron Radiation Emission Spectroscopy
- Measuring energy via frequency



MAC-E filter

- Magnetic Adiabatic Collimation with an Electrostatic Filter
- Measuring energy by applying a **high-pass filter**



PTOLEMY

Calorimeters


- Low-temperature micro calorimeters
- Measuring energy by temperature change




See Talk
06 Sep, 17:50

Elena Ferri

Basic principle of KATRIN-like experiment



Karlsruhe Tritium Neutrino Experiment (KATRIN)



Tritium Laboratory Karlsruhe (TLK)

A facility for high activity tritium experiments

- Two missions:
 - Fuel cycle for fusion reactors
 - **KATRIN Experiment**



We develop safe tritium technology and versatile tritium analytics since 1993

- Licensed for 40 g Tritium
- Closed tritium cycle for recycling and purifying tritium in gram amounts
- > 50 experience scientists, engineers and technicians



We are able to setup and operate a large variety of experiments with tritium

KATRIN timeline

2019-2025

Phase 1 (Integral)
Neutrino mass

See Talk
06 Sep, 18:10

Richard Salomon
Details of the latest
KATRIN data analysis

This talk

2026-2027

Phase 2 (Differential)
keV sterile ν

See Talk
04 Sep, 18:50

Andrea Nava
KATRIN: keV sterile
neutrino search

TRISTAN
detector

2028-2034

R&D Phase for KATRIN++

Atomic Tritium Demonstrator

Differential detection demonstrators

Scientific
goal

Neutrino
mass

This talk

KATRIN data releases and neutrino mass results

2019: $m_\nu < 1.1$ eV (90% CL)

2022: $m_\nu < 0.8$ eV (90% CL)


- ~6 Mio counts

Current data set

- 259 measurement days
- 1757 β -scans
- ~36 Mio counts

Expected sensitivity < 0.5 eV

First 5 campaigns:
taking data while finding optimal operation conditions




PRL 123 (2019) 221802
PRD 104 (2021) 012005

Nature Phys. 18 (2022) 160


Data combination challenges

59 stacked spectra with

$$27 + 28 + 14 \times 28 + 28 + 14 \times 28 + 14 \times 25 + 14 \times 28 = 1609 \text{ data points}$$



Systematic uncertainties



- Statistical uncertainties dominate
- Significant reduction of the background-related systematics
- Better control over source scattering
- Increased conservative uncertainties in this release
 - Reduced uncertainties in current data
 - Reduction of the molecular final-states uncertainties

Forecast: individual systematics in final KATRIN analysis (post 2025)
 expected to be $> 0.01 \text{ eV}^2$ range

New best fit and upper limit

New best fit compatible with vanishing neutrino mass

$$m_\nu^2 = -0.14^{+0.13}_{-0.15} \text{ eV}^2$$




Resulting in a new upper limit

$$m_\nu < 0.45 \text{ eV (90 \% CL)}$$

New KATRIN release improves direct neutrino-mass bound by a factor of 2


<https://arxiv.org/abs/2406.13516>



KATRIN data taking continues

Ongoing data taking through 2025 → $\Sigma 1000$ days

→ target sensitivity below 0.3 eV



Established measurement principles

CRES

- Cyclotron Radiation Emission Spectroscopy
- Measuring energy via frequency



MAC-E filter


- Magnetic Adiabatic Collimation with an Electrostatic Filter
- Measuring energy by applying a **high-pass filter**



PTOLEMY


Calorimeters

- Low-temperature micro calorimeters
- Measuring energy by temperature change




See Elena Ferri

Cyclotron Radiation Emission Spectroscopy



$$f_\gamma = \frac{f_c}{\gamma} = \frac{1}{2\pi m_e + E_{\text{kin}}} \frac{eB}{\gamma}$$



Differential measurement with antenna arrays around a (atomic) tritium source
 → Frequency measurement



PROJECT 8

Project 8 – Results

- **Phase I:** First use of CRES for electron spectroscopy (^{83m}Kr)
- **Phase II:** First use of CRES for tritium beta decay electron spectroscopy
→ Neutrino mass limit ($m_\beta < 155 \text{ eV}$)



A. Ashtari Esfahani et al. Phys. Rev. Lett. 131, 102502 (2023)

Outline

2019-2025

Phase 1 (Integral)
Neutrino mass

2026-2027

Phase 2 (Differential)
keV sterile ν

2028-2034

R&D Phase for KATRIN ++

Atomic Tritium Demonstrator


Differential detection demonstrators

Scientific
goal

Neutrino
mass

- **KATRIN** on way to achieve 1000 d measurement time (**final sensitivity $m_\beta < 0.3$ eV**).
Current m_β result : < 0.45 eV
- We will be ready for **TRISTAN**-Operation at the end of 2025 (**Search for keV sterile neutrinos**)


Going beyond KATRIN



- KATRIN final: $< 0.3 \text{ eV}$ (90% CL)
Distinguish between **degenerate** and **hierarchical** scenario
- New technologies: $< 0.05 \text{ eV}$
Cover **inverted** ordering


KATRIN++ mission

- Next generation m_ν experiment
- Identify and develop scalable technology
- Use KATRIN/TLK infrastructure for R&D phase (~ 7 years)




PROJECT 8

Going beyond KATRIN



Current KATRIN performance (integral, $\Delta E = 2.7 \text{ eV}$, $bg = 0.1 \text{ cps}$)

- Differential measurement ($FWHM < 1 \text{ eV}$)
 - Better use of statistics
 - Lower background
- Atomic tritium
 - Avoid broadening ($\sim 1 \text{ eV}$)
 - Avoid limiting systematics of T_2



KATRIN and TLK as ideal R&D facilities

**Differential detector
technology**



**Atomic source
technology**


KATRIN and TLK as ideal R&D facilities

Further options?

Technologies by community?

Differential detector technology



Option 2
Time-of-flight via electron tagging



- eV resolution for differential detection
- immune to Rydberg-like backgrounds

Option 1
Micro-calorimeters / Quantum sensor

Quantum sensors as high resolution differential detectors




Advantages

- Energy resolution $O(\text{eV})$ compared to conventional detectors $O(100 \text{ eV})$
- Nearly 100% quantum efficiency
- Broad spectrum of possible applications

e.g. Metallic Magnetic Calorimeters (MMC)

- Temperature-dependence in sensor magnetization
- Read-out by SQUID
- Energy resolution:
 - Current: $\Delta E \lesssim 2 \text{ eV}$
 - Midterm: $\Delta E \lesssim 1 \text{ eV}$
 - Future: $\Delta E \sim 100 \text{ meV}$




Not yet tested with external electrons

Next R&D goal: Demonstrate KATRIN with a quantum sensor array

Use existing / well-characterised
beamline




+ $10^6 \times$





with Milano's
cryo expertise

- Challenges of coupling quantum sensor detector array to KATRIN infrastructure
 - Type of quantum sensor
 - Operation in magnetic field (~10 mT)
 - Coupling of mK cryo-platform with RT spectrometer
 - Large area detector and multiplexing of ~1e6 channels
 - Limits to energy resolution




ELECTRON: e^- spectroscopy with quantum sensors

8 channel detector chips
& front-end SQUID chips





KIT-IMS (Kempf group)

Metallic Magnetic Calorimeters (MMC)




Calorim. Kr-83m spectrum @ highest resolution
Next: tritium spectroscopy

FWHM ~ 25 eV



Next R&D goal: Demonstrate single electron tagging for ToF



Single electron tagging is challenging

Cyclotron radiation
emission detection
(CRES)

Coreless cryogenic
current comparator

**Tiny signals vs.
minimal noise floor**

Strategy

- „Single channel“ detector
→ less complex than quantum sensor array (QSA)
- Differential measurement with ToF before QSA ready
- Work on techniques to improve ToF resolution (U Münster)

R&D ongoing at U North Carolina


KATRIN and TLK as ideal R&D facilities

**Differential detector
technology**




**Atomic source
technology**

KATRIN and TLK as ideal R&D facilities



- Molecular effects → spectral broadening



Atomic source technology


PROJECT 8

Atomic Tritium Demonstrator at TLK

2022-2025
TLK

2025-2028
TLK+Mainz

2028...
TLK+P8?/...




Aim for investigation

- Develop atom cooling mechanism
- Trapping times / max. densities
- Interplay of beta-driven plasma (meV–eV) and ultra-cold trapped atoms (neV)


Tritium atom throughput on the order of 10 g/day (c.f. KATRIN: 40 g/day)

Atomic source R&D progress



Mass spectrometer

Source



Decontaminated
glove box

- Non-tritium hydrogen cracker being operated
- Characterization measurements for tritium beamline ongoing

- Installation of first ever atomic **tritium** source at TLK ongoing
- First results expected in 2024

Atomic Tritium Demonstrator at TLK

2022-2025
TLK

2025-2028
TLK+Mainz


2028...
TLK+?/...

Molecular tritium

Atomic tritium
(hot, 2500 K)

Atomic tritium
(cold)

Atomic tritium
(trapped, injected)




TLK is offering room and infrastructure
for extended atomic tritium experiments

Final R&D goal

Atomic tritium with Quantum sensor array

Conceptual idea




Work started

- concept
- sensitivities
- R&D

CRES plans aiming at 40-50 meV sensitivity

PROJECT 8

<https://www.project8.org/>




Concept: CRES readout
in **magneto-gravitational trap** for atomic tritium

Staged approach:

- Proof-of-principle
- Demonstrators
- Final experiments in 2030s

H/D/T atom supersonic beam discharge source (30 K)




<https://www.hep.ucl.ac.uk/qtnm/>




Concept: CRES readout
in **race-track** for atomic tritium

PTOLEMY Project




- Aim: direct detection of big-bang neutrinos; determination of neutrino mass is „by-product“

M.G. Betti et al JCAP07(2019)047




Combine technologies (TES, CRES, novel drift filter) with large scale $O(100\text{g})$ tritiated graphene target

Content from „Nicola Rossi,
EPS-HEP July 2023 Hamburg“




<https://ptolemy.lngs.infn.it/>



Start of technology demonstrator
@LNGS soon

KATRIN++ and other next generation projects

- Currently, **no technology proven** to reach ultimate sensitivity
- Neutrino mass detection must be confirmed by **independent technologies**
- **Atomic tritium trap** is key independently of detection techniques
- CRES, bolometer and ToF complementary



Atomic source research

Atomic tritium trap is key independently of detection techniques

Mission: Realize (global) Atomic Tritium Demonstrator (at TLK)

Form joint working group
(kick-off meeting in September)

Possible partners:


Ubachs (VU Amsterdam), T₂, DT, HT

Schiller (U Düsseldorf) HT⁺


Pohl (U Mainz) T

....

**placeholder name



Beta decay and neutrino mass: KATRIN and beyond



- KATRIN on way to achieve 1000 d measurement time (**final sensitivity $m_\beta < 0.3$ eV**).
Current m_β result : < 0.45 eV
- We will be ready for TRISTAN-Operation at the end of 2025 (**Search for keV sterile neutrinos**)
- Ultimate neutrino mass experiment (Normal Ordering; **sensitivity on $m_\beta < 40$ meV**) requires **differential detector principle** und **an atomic tritium source** → R&D Plan for PoF-V
- KATRIN++ invites research groups for **tackling challenges together**




KATRIN collaboration




<https://www.linkedin.com/company/tritiumlab/>




Atomic vs molecular tritium



Going beyond KATRIN




Current KATRIN performance (integral, $\Delta E = 2.7 \text{ eV}$, $\text{bg} = 0.1 \text{ cps}$)



Improved measurement principle




Integral measurement (high pass filter)

- Energy resolution determined by filter
- Detector „only“ counts
- Reduced statistics




Differential measurement

- Energy resolution determined by
 - A) detector**
 - or
 - B) time of flight**



Next R&D goal: Demonstrate single electron tagging for ToF

Pre-spectrometer:
limit rate < 1 khz



KATRIN Source:
 10^{11} Bq


Tagger: **start** (~1000 Hz)

Fast detector: **stop**

Main spectrometer:
delay line due to retardation pot.

Single electron tagging is challenging


Cyclotron radiation
emission detection
(CRES)



See Project 8

Coreless cryogenic
current comparator

**Tiny signals
vs minimal
noise floor**



T. Tanabe et al., Nucl. Instr. Meth. A 427 (1999) 455

Strategy

- „Single channel“ detector
→ less complex than quantum sensor array (QSA)
- Differential measurement with ToF before QSA is ready
- Work on techniques to improve ToF resolution (U Münster)

KAMATE – Karlsruhe Mainz Atomic Tritium experiment



Scientific / technical goals

- Atomic beam characterization
 - Atomic fraction
 - Maximal flow rates / pressure limits
 - Isotopic effects
- Angular dispersion
- Time-of-flight (upgrade)
- Wire-detector

- Cooling / accommodation (upgrade)
 - Velocity measurement
 - Recombination



Karlsruhe



Mainz

Sophisticated setup based on Mainz setup

Multi chamber / collimation design, tilting mechanism, beam control, source parameter control, beam analytics

KAMATE – Karlsruhe Mainz Atomic Tritium experiment



A. Lindman

2024

KAMATE stages

2028

KAMATE 0.5 (at Mainz)

Identify best source at MATS with H/D

KAMATE 1.0 (at TLK)

Operate KAMATE 0.5 setup with T.

T(Beam) ~ 2500 K

KAMATE 2.0 (at TLK)

Add accommodator as first stage cooling.


T(Beam) ~ 150 K

KAMATE 3.0 (at TLK)

Add nozzle for second stage cooling and beam temperature measurement setup (time of flight).

T(Beam) ~ 4 K

Preparing TLK for Atomic Tritium Demonstrator (ATD)



PETRA box

- In process of disposal/repurposing of former experiment
- Next step: decontamination

Other systems

e.g. former tritium retention system, gas bottles, control cabinets, ...

→ Relocation in progress

ALTEX Box

- Decontamination completed
- Installation ongoing

ATD Glove box infrastructure

e.g. large tritium retention system (~500k€), tritium supplies, ...

→ Acquisition on-going

Working principle of KATRIN

