



Contribution ID: 68

Type: oral

Ultra-high acceleration gradient using structured nanomaterials

Thursday, 12 September 2024 10:10 (20 minutes)

Solid-state plasma wakefield acceleration has recently garnered attention as a viable alternative for achieving unprecedented ultra-high acceleration gradients on the order of 1 TV/m or beyond [1, 2]. In this context, recent advancements in nanofabrication techniques [3] have opened up the possibility of creating structured plasmas with inhomogeneous properties. For instance, the utilization of carbon nanotube (CNT) bundles and multi-layer graphene [4] holds great potential for generating stable plasmas with electron densities reaching as high as 10^{24} cm^{-3} , i.e., orders of magnitude higher than conventional gaseous plasmas. As part of a new collaborative effort called NanoAcc (Application of Nanostructures in Accelerator Physics), we have conducted Particle-In-Cell (PIC) simulations to investigate both laser-driven and beam-driven excitation of pre-ionized targets utilizing CNT arrays. Our results confirm the attainment of wakefields at the TV/m scale. Additionally, we have observed phenomena such as self-injection, sub-femtosecond bunch formation, and the acceleration of electrons within micrometre-scale targets, leading to kinetic energies of approximately 10 MeV. These findings open up promising possibilities for manipulating charged particle beams, thereby shaping the future of compact accelerator design and radiation sources. Furthermore, solid-state plasmas offer a high degree of tunability in extracting relevant bunch parameters through effective control over the target structure. In this article, we present an overview of the studies conducted thus far by the NanoAcc collaboration, and discuss future experimental plans as well as potential applications.

References

1. Y. M. Shin, A. H. Lumpkin and R. M. Thurman-Keup, Nucl. Instr. and Meth. B 355 (2015) 94.
2. A. Bonatto et al., Phys. Plasmas 30 (2023) 033105.
3. F. Sarasini et al., Composites Part B 243 (2022) 110136.
4. C. Bontoiu et al., Scientific Reports 13 (2023) 1330.

Primary author: RESTA LOPEZ, Javier (ICMUV-University of Valencia)

Co-authors: BONATTO, Alexandre (School of Physics and Astronomy, University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom); SYTOV, Alexei (Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara; Korea Institute of Science and Technology Information); WELSCH, Carsten Peter; BONTOIU, Cristian; GATTI, Giancarlo; CAVOTO, Gianluca (Istituto Nazionale di Fisica Nucleare); XIA, Guoxing (Cockcroft Institute and the University of Manchester); DREBOT, Ilya (Istituto Nazionale di Fisica Nucleare); BANDIERA, LAURA (Istituto Nazionale di Fisica Nucleare); MARTÍN-LUNA, Pablo

Presenter: SYTOV, Alexei (Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara; Korea Institute of Science and Technology Information)

Session Classification: New Concepts