

Exploring the Quantum Universe: Unveiling Mysteries with Flavour Physics*

2 Jul 2024 – Bari

Niels Tuning (Nikhef)

*courtesy chatGPT...

ChatGPT ▾

Can you give me an exciting title for a seminar on flavour physics?

Historical record of indirect discoveries

GIM mechanism in $K^0 \rightarrow \mu\mu$

Weak Interactions with Lepton-Hadron Symmetry*


S. L. GLASHOW, J. ILIOPOULOS, AND L. MAIANI†
 Lyman Laboratory of Physics, Harvard University, Cambridge, Massachusetts 02139
 (Received 5 March 1970)

We propose a model of weak interactions in which the currents are constructed out of four basic quark fields and interact with a charged massive vector boson. We show, to all orders in perturbation theory, that the leading divergences do not violate any strong-interaction symmetry and the next to the leading divergences respect all observed weak-interaction selection rules. The model features a remarkable symmetry between leptons and quarks. The extension of our model to a complete Yang-Mills theory is discussed.

splitting, beginning at order $G(GA^2)$, as well as contributions to such unobserved decay modes as $K_2 \rightarrow \mu^+ + \mu^-$, $K^+ \rightarrow \pi^+ + l + \bar{l}$, etc., involving neutral lepton

We wish to propose a simple model in which the divergences are properly ordered. Our model is founded in a quark model, but one involving four, not three, fundamental fermions; the weak interactions are mediated

new quantum number C for charm.



Glashow, Iliopoulos, Maiani,
 Phys.Rev. D2 (1970) 1285

“Discovery” of charm

CP violation, $K_L^0 \rightarrow \pi\pi$

27 JULY 1964

EVIDENCE FOR THE 2π DECAY OF THE K_2^0 MESON*†

J. H. Christenson, J. W. Cronin,† V. L. Fitch,† and R. Turlay§
 Princeton University, Princeton, New Jersey
 (Received 10 July 1964)

This Letter reports the results of experimental studies designed to search for the 2π decay of the K_2^0 meson. Several previous experiments have

Progress of Theoretical Physics, Vol. 49, No. 2, February 1973

CP-Violation in the Renormalizable Theory of Weak Interaction

Makoto KOBAYASHI and Toshihide MASKAWA

Department of Physics, Kyoto University, Kyoto

(Received September 1, 1972)

doublet with the same charge assignment. This is because all phases of elements of a 3×3 unitary matrix cannot be absorbed into the phase convention of six fields. This possibility of CP-violation will be discussed later on.

Christenson, Cronin, Fitch, Turlay,
 Phys.Rev.Lett. 13 (1964) 138
 Kobayashi, Maskawa,
 Prog.Theor. Phys. 49 (1973) 652

“Discovery” of beauty

$B^0 \leftrightarrow \bar{B}^0$ mixing

DESY 87-029
 April 1987

OBSERVATION OF $B^0 - \bar{B}^0$ MIXING

The ARGUS Collaboration

In summary, the combined evidence of the investigation of B^0 meson pairs, lepton pairs and B^0 meson-lepton events on the $\Upsilon(4S)$ leads to the conclusion that $B^0 - \bar{B}^0$ mixing has been observed and is substantial.


Parameters	Comments
$r > 0.09$ 90%CL	This experiment
$x > 0.44$	This experiment
$B^{\frac{1}{2}} t_B \approx t_\pi < 160 \text{ MeV}$	B meson (\approx pion) decay constant
$m_b < 5 \text{ GeV}/c^2$	b-quark mass
$\tau_b < 1.4 \cdot 10^{-12} \text{ s}$	B meson lifetime
$ V_{td} < 0.018$	Kobayashi-Maskawa matrix element
$\eta_{\text{QCD}} < 0.86$	QCD correction factor [17]
$m_t > 50 \text{ GeV}/c^2$	t quark mass

ARGUS Coll.
 Phys.Lett.B192 (1987) 245

“Discovery” of top

Historical record of indirect discoveries

Particle	Indirect			Direct		
ν	β decay	Fermi	1932	Reactor ν -CC	Cowan, Reines	1956
W	β decay	Fermi	1932	$W \rightarrow e\nu$	UA1, UA2	1983
c	$K^0 \rightarrow \mu\mu$	GIM	1970	J/ψ	Richter, Ting	1974
b	CPV $K^0 \rightarrow \pi\pi$	CKM, 3 rd gen	1964/72	Y	Ledermann	1977
Z	ν -NC	Gargamelle	1973	$Z \rightarrow e^+e^-$	UA1	1983
t	B mixing	ARGUS	1987	$t \rightarrow Wb$	D0, CDF	1995
H	e^+e^-	EW fit, LEP	2000	$H \rightarrow 4\mu/\gamma\gamma$	CMS, ATLAS	2012
?	What's next ?					?



Outline

- CKM elements

- $\sin 2\beta$
- γ
- Δm_s
- V_{ub}

- Anomalies

- $b \rightarrow c \tau \nu$
- $b \rightarrow s \ell^+ \ell^-$

- ~~Hadron physics~~

- Heavy ion programme
- Spectroscopy

To the backup slides...

- Prospects

- Upgrade II


CKM at the heart of the SM

$$\begin{aligned}\mathcal{L} = & -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \\ & + i\bar{\psi} \not{D} \psi + \text{h.c.} \\ & + \bar{\psi}_i Y_{ij} \psi_j \phi + \text{h.c.} \\ & + |D_\mu \phi|^2 - V(\phi)\end{aligned}$$


$$\bar{\psi}_i Y_{ij} \psi_j \phi$$

(CKM: a quick reminder...)

1) Matrix to transform weak- and mass-eigenstates:




i.e. diagonalize:



(CKM: a quick reminder...)

1) Matrix to transform weak- and mass-eigenstates:




2) Matrix has complex phases:

$$\begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}|e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}|e^{-i\beta} & -|V_{ts}|e^{i\beta_s} & |V_{tb}| \end{pmatrix}$$

3) Matrix is unitary:

$$V^\dagger V = \begin{pmatrix} V_{ud}^* & V_{us}^* & V_{ub}^* \\ V_{cd}^* & V_{cs}^* & V_{cb}^* \\ V_{td}^* & V_{ts}^* & V_{tb}^* \end{pmatrix} \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$V_{ub}^* V_{ud} = 0 \quad V_{cb}^* V_{cd} = 0 \quad V_{tb}^* V_{td} = 0$$



CKM: (1995) LHCb Letter-of-Intent

- LHC-B Letter-of-Intent 1995








Figure 2.1: Limits on the CKM parameters (1σ) ρ and η for $m_t = 174 \text{ GeV}$. The annular region cen-

CKM: (1995) LHCb Letter-of-Intent

- LHC-B Letter-of-Intent 1995



CKM: recent results



$$\left(\begin{array}{ccc}
 |V_{ud}| & |V_{us}| & |V_{ub}|e^{-i\gamma} \\
 -|V_{cd}| & |V_{cs}| & |V_{cb}| \\
 |V_{td}|e^{-i\beta} & -|V_{ts}|e^{i\beta_s} & |V_{tb}|
 \end{array} \right)$$

Global fits:
 CKMfitter: <http://ckmfitter.in2p3.fr/>
 UTFit: <http://www.utfit.org/Utfit/>

Outline

- CKM elements

- $\sin 2\beta$
- γ
- Δm_s
- V_{ub}

- Anomalies

- $b \rightarrow c \tau \nu$
- $b \rightarrow s \ell^+ \ell^-$

- ~~Hadron physics~~

- Heavy ion programme
- Spectroscopy


To the backup slides...

- Prospects

- Upgrade II

sin2β



- CP violation:
 - Two interfering amplitudes
 - Two relative phases
 - Different amplitude under CP conjugation
- $B^0 \rightarrow J/\psi K_S^0$: The golden mode!
 - Relative phase: $\arg(V_{td}^2) = 2\beta$ (and $\pi/2$)




sin2β

$$\begin{aligned}
 \mathcal{A}_{[c\bar{c}]K_S^0}(t) &\equiv \frac{\Gamma(\bar{B}^0(t) \rightarrow [c\bar{c}]K_S^0) - \Gamma(B^0(t) \rightarrow [c\bar{c}]K_S^0)}{\Gamma(\bar{B}^0(t) \rightarrow [c\bar{c}]K_S^0) + \Gamma(B^0(t) \rightarrow [c\bar{c}]K_S^0)} \\
 &= \frac{S \sin(\Delta m t) - C \cos(\Delta m t)}{\cosh(\Delta\Gamma t/2) + A_{\Delta\Gamma} \sinh(\Delta\Gamma t/2)} \approx S \sin(\Delta m t)
 \end{aligned}$$

- “Flavour tagging” essential
 - Which B⁰ was a \bar{B}^0 ?



The LHCb Detector



23 sep 2010
Run 79646



19:49:24
Event 143858637



sin2β

$$\begin{aligned}
 \mathcal{A}_{[c\bar{c}]K_S^0}(t) &\equiv \frac{\Gamma(\bar{B}^0(t) \rightarrow [c\bar{c}]K_S^0) - \Gamma(B^0(t) \rightarrow [c\bar{c}]K_S^0)}{\Gamma(\bar{B}^0(t) \rightarrow [c\bar{c}]K_S^0) + \Gamma(B^0(t) \rightarrow [c\bar{c}]K_S^0)} \\
 &= \frac{S \sin(\Delta m t) - C \cos(\Delta m t)}{\cosh(\Delta\Gamma t/2) + A_{\Delta\Gamma} \sinh(\Delta\Gamma t/2)} \approx S \sin(\Delta m t) \quad (S = \sin 2\beta)
 \end{aligned}$$

- “Flavour tagging” essential
 - Wrong tag fraction $w \sim 39\%$
 - $D = (1 - 2w) \sim 0.22$



sin2β

NEW:

$$S_{J/\psi(\rightarrow\mu^+\mu^-)K_S^0} = 0.716 \pm 0.015 \text{ (stat)} \pm 0.007 \text{ (syst)}$$

(Run 2) LHCb [arXiv:2309.09728](https://arxiv.org/abs/2309.09728) PRL132 (2024) 021801

OLD:

$$\sin(2\beta) \equiv \sin(2\phi_1)$$


HFLAV
2021

BaBar: $\sin 2\beta = 0.691 \pm 0.031$

Belle: $\sin 2\beta = 0.667 \pm 0.026$

LHCb: $\sin 2\beta = 0.760 \pm 0.034$
(Run 1)

Avg: $\sin 2\beta = 0.699 \pm 0.017$





- Large B production competes with good tagging:

$\sigma_{\text{stat}}(\mathcal{S}(J/\psi K_S^0))$	now	50 ab ⁻¹	
Belle/II	0.029	0.005	
	now	50 fb ⁻¹	300 fb ⁻¹
LHCb	0.035	0.006	0.003

φ_s with $B_s^0 \rightarrow J/\psi \phi$

("the $\sin 2\beta$ of the B_s^0 system")



ϕ_s with $B_s^0 \rightarrow J/\psi \phi$




- Some challenges:

- 1) Rapid B_s^0 oscillations: decay time resolution
- 2) "Same side" kaon-tagging: calibration with hadronic final state
- 3) Mix of CP eigenstates: angular analysis

1) Decay time resolution from prompt J/ψ

2) Tagging calibration from $B_s^0 \rightarrow D_s \pi$



3) Angular analysis to disentangle CP + and CP -



LHCb, [arXiv:2308.01468](https://arxiv.org/abs/2308.01468)
Phys.Rev.Lett. 132 (2024) 5, 051802

- LHCb 2011-2016

2012



$\phi_s = -50 \pm 19 \text{ mrad (HFLAV, 2021)}$

$\phi_s = -44 \pm 20 \text{ mrad (LHCb, 2023)}$ (9 fb⁻¹)



$\phi_s = -37 \pm 1 \text{ mrad (SM, CKMfitter)}$

CKMfitter,
Phys. Rev. D84, 033005 (2011),
updated with Summer 2019 results

LHCb, [arXiv:2308.01468](https://arxiv.org/abs/2308.01468)
Phys.Rev.Lett. 132 (2024) 5, 051802

ϕ_s at CMS

- Clean sample of $B_s^0 \rightarrow J/\psi \phi$ decays (2017-2018, 96 fb^{-1})
 - new pioneering flavor tagging algorithm




➤ CPV at 3.2σ from 0

$$\phi_s = -74 \pm 23 \text{ mrad (CMS)}$$



$$\phi_s = -44 \pm 20 \text{ mrad (LHCb)}$$

Constraints on angle γ

- Different yields for B^+ and B^- decays
 - two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$






Constraints on angle γ - with $B^\pm \rightarrow D^{(*)}K^\pm$ and $D^0 \rightarrow h^\pm h^\pm$



$$\Gamma(B^\pm \rightarrow [CP]_D h^\pm) \propto 1 + (r_B^{Dh})^2 + 2r_B^{Dh} \cos(\delta_B^{Dh} \pm \gamma)$$

Constraints on angle γ - with $B^\pm \rightarrow D^{(*)}K^\pm$ and $D^0 \rightarrow h^\pm h^\pm$



$$\Gamma(B^\pm \rightarrow [CP]_D h^\pm) \propto 1 + (r_B^{Dh})^2 + 2r_B^{Dh} \cos(\delta_B^{Dh} \pm \gamma)$$

- Many final states for D^{*0} or D^0 !




- $B^\pm \rightarrow D^0 K^\pm, B^\pm \rightarrow D^0 \pi^\pm, B^\pm \rightarrow D^{*0} K^\pm, B^\pm \rightarrow D^{*0} \pi^\pm$
- $D^0 \rightarrow K^+ K^-, D^0 \rightarrow K^+ \pi^-, D^0 \rightarrow \pi^+ \pi^-, D^0 \rightarrow K_S^0 K^- K^+, D^0 \rightarrow K_S^0 \pi^+ \pi^-,$

- Very precise input for gamma

Constraints on angle γ - with $B^\pm \rightarrow D^{*0}h^\pm$ and $D^0 \rightarrow K_S^0 h^+ h^-$

- Different yields for B^+ and B^- decays

2 amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$



$B^+ \rightarrow D^*K^+, D^* \rightarrow D\pi^0$	112 ± 7
$B^+ \rightarrow D^*K^+, D^* \rightarrow D\gamma$	358 ± 33
$B^- \rightarrow D^*K^-, D^* \rightarrow D\pi^0$	109 ± 6
$B^- \rightarrow D^*K^-, D^* \rightarrow D\gamma$	419 ± 35

$$\gamma = (69^{+13}_{-14})^\circ$$

Component	Reconstructed $B^\pm \rightarrow DK^\pm$
$B^\pm \rightarrow D^*[D\pi^0]K^\pm$	6244 ± 12
$B^\pm \rightarrow D^*[D\pi^0]\pi^\pm$	340 ± 1
$B^\pm \rightarrow D^*[D\gamma]K^\pm$	3144 ± 6
$B^\pm \rightarrow D^*[D\gamma]\pi^\pm$	166 ± 1

$$\gamma = (92^{+21}_{-17})^\circ$$


LHCb, JHEP12(2023)013, [arXiv:2310.04277](https://arxiv.org/abs/2310.04277)

LHCb, JHEP 02 (2024) 118, [arXiv:2311.10434](https://arxiv.org/abs/2311.10434)

CKM angle γ : Combination

- Different yields for B and anti- B decays
 - two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$
 - many $D^{(*)}_{(s)}$ final states:

B decay	D decay	Ref.	Dataset	Status since Ref. [14]
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-$	[29]	Run 1&2	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[30]	Run 1	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K^\pm\pi^\mp\pi^+\pi^-$	[18]	Run 1&2	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-\pi^0$	[19]	Run 1&2	Updated
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 h^+ h^-$	[31]	Run 1&2	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 K^\pm \pi^\mp$	[32]	Run 1&2	As before
$B^\pm \rightarrow D^+ h^\pm$	$D \rightarrow h^+ h^-$	[29]	Run 1&2	As before
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+ h^-$	[33]	Run 1&2(*)	As before
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[33]	Run 1&2(*)	As before
$B^\pm \rightarrow Dh^\pm\pi^\mp$	$D \rightarrow h^+ h^-$	[34]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^+ h^-$	[35]	Run 1&2(*)	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[35]	Run 1&2(*)	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_S^0 \pi^+ \pi^-$	[36]	Run 1	As before
$B^0 \rightarrow D^+ \pi^\pm$	$D^+ \rightarrow K^- \pi^+ \pi^+$	[37]	Run 1	As before
$B^0 \rightarrow D_s^+ K^\pm$	$D_s^+ \rightarrow h^+ h^- \pi^\pm$	[38]	Run 1	As before
$B_s^0 \rightarrow D_s^+ K^\pm \pi^\mp$	$D_s^+ \rightarrow h^+ h^- \pi^\pm$	[39]	Run 1&2	As before
D decay	Observable(s)	Ref.	Dataset	Status since Ref. [14]
$D^0 \rightarrow h^+ h^-$	ΔA_{CP}	[24, 40, 41]	Run 1&2	As before
$D^0 \rightarrow K^+ K^-$	$A_{CP}(K^+ K^-)$	[16, 24, 25]	Run 2	New
$D^0 \rightarrow h^+ h^-$	$y_{CP} - y_{CP}^{K^+ \pi^+}$	[42]	Run 1	As before
$D^0 \rightarrow h^+ h^-$	$y_{CP} - y_{CP}^{K^+ \pi^+}$	[15]	Run 2	New
$D^0 \rightarrow h^+ h^-$	ΔY	[43-46]	Run 1&2	As before
$D^0 \rightarrow K^+ \pi^-$ (Single Tag)	$R^\pm, (x'^\pm)^2, y'^\pm$	[47]	Run 1	As before
$D^0 \rightarrow K^+ \pi^-$ (Double Tag)	$R^\pm, (x'^\pm)^2, y'^\pm$	[48]	Run 1&2(*)	As before
$D^0 \rightarrow K^\pm \pi^\mp \pi^+ \pi^-$	$(x^2 + y^2)/4$	[49]	Run 1	As before
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	x, y	[50]	Run 1	As before
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[51]	Run 1	As before
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[52]	Run 2	As before
$D^0 \rightarrow K_S^0 \pi^+ \pi^-$ (μ^- tag)	$\bar{x}_{CP}, \bar{y}_{CP}, \Delta \bar{x}, \Delta \bar{y}$	[17]	Run 2	New




CKM angle γ


- Different yields for B and anti- B decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$
- many $D^{(*)}_{(s)}$ final states:

B decay	D decay	Ref.	Dataset	Status since Ref. [14]
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-$	[29]	Run 1&2	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[30]	Run 1	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K^\pm\pi^\mp\pi^+\pi^-$	[18]	Run 1&2	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-\pi^0$	[19]	Run 1&2	Updated
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 h^\pm h^\mp$	[31]	Run 1&2	As before
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 K^\pm\pi^\mp$	[32]	Run 1&2	As before
$B^\pm \rightarrow D^+h^\pm$	$D \rightarrow h^+h^-$	[29]	Run 1&2	As before
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+h^-$	[33]	Run 1&2(*)	As before
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[33]	Run 1&2(*)	As before
$B^\pm \rightarrow Dh^\pm\pi^+\pi^-$	$D \rightarrow h^+h^-$	[34]	Run 1	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^+h^-$	[35]	Run 1&2(*)	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[35]	Run 1&2(*)	As before
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_S^0\pi^+\pi^-$	[36]	Run 1	As before
$B^0 \rightarrow D^+\pi^\pm$	$D^+ \rightarrow K^-\pi^+\pi^+$	[37]	Run 1	As before
$B^0 \rightarrow D_s^+K^\pm$	$D_s^+ \rightarrow h^+h^-\pi^+$	[38]	Run 1	As before
$B_s^0 \rightarrow D_s^+K^\pm\pi^+\pi^-$	$D_s^+ \rightarrow h^+h^-\pi^+$	[39]	Run 1&2	As before
D decay	Observable(s)	Ref.	Dataset	Status since Ref. [14]
$D^0 \rightarrow h^+h^-$	ΔA_{CP}	[24, 40, 41]	Run 1&2	As before
$D^0 \rightarrow K^+K^-$	$A_{CP}(K^+K^-)$	[16, 24, 25]	Run 2	New
$D^0 \rightarrow h^+h^-$	$y_{CP} - y_{CP}^{K^+\pi^+}$	[42]	Run 1	As before
$D^0 \rightarrow h^+h^-$	$y_{CP} - y_{CP}^{K^+\pi^+}$	[15]	Run 2	New
$D^0 \rightarrow h^+h^-$	ΔY	[43-46]	Run 1&2	As before
$D^0 \rightarrow K^+\pi^-$ (Single Tag)	$R^\pm, (x^\pm)^2, y^\pm$	[47]	Run 1	As before
$D^0 \rightarrow K^+\pi^-$ (Double Tag)	$R^\pm, (x^\pm)^2, y^\pm$	[48]	Run 1&2(*)	As before
$D^0 \rightarrow K^\pm\pi^\mp\pi^+\pi^-$	$(x^2 + y^2)/4$	[49]	Run 1	As before
$D^0 \rightarrow K_S^0\pi^+\pi^-$	x, y	[50]	Run 1	As before
$D^0 \rightarrow K_S^0\pi^+\pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[51]	Run 1	As before
$D^0 \rightarrow K_S^0\pi^+\pi^-$	$x_{CP}, y_{CP}, \Delta x, \Delta y$	[52]	Run 2	As before
$D^0 \rightarrow K_S^0\pi^+\pi^-$ (μ^- tag)	$\bar{x}_{CP}, \bar{y}_{CP}, \Delta \bar{x}, \Delta \bar{y}$	[17]	Run 2	New




	γ ($^\circ$)
LHCb	$63.8^{+3.5}_{-3.7}$
CKMfitter	$65.6^{+1.1}_{-2.7}$
UTFit	$65.8^{+2.2}_{-2.2}$



LHCb-CONF-2022-002, Oct 2022

Precision Δm_s with $B_s^0 \rightarrow D_s^+ \pi^-$




- Frequency \sim transition rate!
- “Flavour specific” : final state reveals flavour of the decaying B



Precision Δm_s with $B_s^0 \rightarrow D_s^+ \pi^-$


- Legacy “textbook” measurement
- “Flavour specific” : final state reveals flavour of the decaying B
- 3 trillion oscillations per second, with 3×10^{-4} precision

— $B_s^0 \rightarrow D_s^- \pi^+$ — $\bar{B}_s^0 \rightarrow D_s^- \pi^+$ — Untagged



	Δm_s (ps ⁻¹)	Stat	Sys	Ref.
$B_s^0 \rightarrow D_s^+ \pi^-$	17.7683	0.0051	0.0032	arXiv:2104.04421 acc. Nat.Phys
$B_s^0 \rightarrow D_s^+ \pi^- \pi^- \pi^-$	17.757	0.007	0.008	arXiv:2011.12041 JHEP 03(2021)137
Combination	17.7656	0.0057		arXiv:2104.04421 acc. Nat.Phys


Measurement $|V_{ub}|/|V_{cb}|$ from $B(B_s^0 \rightarrow K^- \mu^+ \nu)$




$$\mathcal{B}(B_s \rightarrow K \mu \nu) / \mathcal{B}(B_s \rightarrow D_s \mu \nu) = |V_{ub}|^2 / |V_{cb}|^2 \times \text{FF}_K / \text{FF}_{D_s}$$

- Interesting input to $|V_{ub}|$! (and form factor calculations)

Measurement $|V_{ub}|/|V_{cb}|$ from $B(B_s^0 \rightarrow K^- \mu^+ \nu)$



LHCb, [arXiv:2012.05143](https://arxiv.org/abs/2012.05143) PRL126(2021)8, 081804



$$R_{BF} = \mathcal{B}(B_s \rightarrow K \mu \nu) / \mathcal{B}(B_s \rightarrow D_s \mu \nu) = \frac{N_K}{N_{D_s}} \frac{\epsilon_{D_s}}{\epsilon_K} \times \mathcal{B}(D_s \rightarrow K K \pi)$$

$$\mathcal{B}(B_s \rightarrow K \mu \nu) = (1.06 \pm 0.05(\text{stat})) \pm 0.04(\text{syst}) \pm 0.06(\text{ext}) \pm 0.04(\text{FF}) \times 10^{-4}$$


- First observation of $B_s^0 \rightarrow K^- \mu^+ \nu$

$$R_{BF} = |V_{ub}|^2 / |V_{cb}|^2 \times \text{FF}_K / \text{FF}_{D_s}$$


$$|V_{ub}|/|V_{cb}|(\text{low}) = 0.0607 \pm 0.0015(\text{stat}) \pm 0.0013(\text{syst}) \pm 0.0008(D_s) \pm 0.0030(FF),$$

$$|V_{ub}|/|V_{cb}|(\text{high}) = 0.0946 \pm 0.0030(\text{stat})^{+0.0024}_{-0.0025}(\text{syst}) \pm 0.0013(D_s) \pm 0.0068(FF). \quad (?)$$


CKM: recent results




CKM: recent results




- So far so good, but stay vigilant...
 - V_{ub} and V_{cb} : incl. and excl. measurements differ...
 - V_{us} : too small for unitarity (Cabibbo angle anomaly)
 - $K\pi$ puzzle: CP asymmetries should be related through isospin symmetry...
 - $BR(B \rightarrow Dh)$: Factorisation?
 - ...




Crivellin et al, arXiv:2212.06862



Fleischer, Jaarsma, Malami, Vos, arXiv:1806.08783




Skidmore, 2 Jun 2022, Siegen Workshop




Outline

- CKM elements
 - $\sin 2\beta$
 - γ
 - Δm_s
 - V_{ub}
- Anomalies
 - $b \rightarrow c \tau \nu$
 - $b \rightarrow s \ell^+ \ell^-$
- Hadron physics
 - Heavy ion programme
 - Spectroscopy
- Prospects
 - Upgrade II

Anomalies



Semileptonic
CC
 $b \rightarrow cl\nu$



"Semileptonic"
FCNC EWP Penguin
 $b \rightarrow sl^+l^-$



R(D*) vs R(D)

- Signal: *distinguish "□" from "□-from-τ" ...*

- $B^0 \rightarrow D^{*+} l^- \nu$ → (D*+μ) sample
- $B^+ \rightarrow D^0 l^- \nu$ → (D⁰μ) sample

- Main backgrounds:


- $B \rightarrow DDX$
- $B \rightarrow D^{**} \mu^- \nu$



PRL 115 (2015) 111803

R(D*) vs R(D)


- Simultaneous 3D-fit to 8 samples (and in 4 q² bins...)



o-PAPER-2022-039 supplementary (in preparation)


R(D*) vs R(D)

- World average 3.3σ to 3.2σ



R(D*) vs R(D)

- World average 3.3σ to 3.2σ




New measurement of $R(D^*)$ (hadronic tau decay)

- World average 3.3σ to 3.2σ to **3.34σ**

LHCb ^b	$0.267 \pm 0.012 \pm 0.019$
-------------------	-----------------------------

LHCb, [arXiv:2305.01463](https://arxiv.org/abs/2305.01463) Phys. Rev. D108 (2023) 012018




New measurement of $R(D^+)$, $R(D^{*+})$ (muonic tau decay)


- World average 3.3σ to 3.2σ to 3.34σ to **3.33σ**

LHCb ^c	$0.402 \pm 0.081 \pm 0.085$	$0.249 \pm 0.043 \pm 0.047$
-------------------	-----------------------------	-----------------------------


LHCb, arXiv:2406.03387



CC and FCNC




Semileptonic
CC
 $b \rightarrow cl\nu$



"Semileptonic"
FCNC EWP Penguin
 $b \rightarrow sl^+l^-$

Decay rates

- Study same process with **different** hadrons:




$b \rightarrow s |^+ |^-$


T.Blake et al. arXiv:1606.00916

Rich laboratory:

- 1) Purely leptonic
- 2) Decay rates
- 3) Angular asymmetries
- 4) Ratio of decay rates









Decay rates: consistently low



Decay rates: consistently low



New LQCD calc: Parrot, Bouchard, Davies, [HPQCD], arXiv:2207.13371



Non-local FF: Gubernari, van Dyk, Reboud, Virto, arXiv:2206.03797


Angular asymmetries

P_5'



Angular asymmetries: eg. P_5'


- Compilation:



LHCb, [arXiv:2003.04831](https://arxiv.org/abs/2003.04831), Phys. Rev. Lett. 125 (2020) 011802
Plot from:
Albrecht, van Dyk, Langenbruch, PPNP120 (2021) 103885, arXiv:2107.04822


Angular asymmetries

- Interesting to compare angular asymmetries for μ and e




$B^0 \rightarrow K^{0*} \mu^+ \mu^-$: more than just P_5'

LHCb Coll, arXiv:2003.04831




Coherent pattern






arXiv:2003.04831: $B^0 \rightarrow K^{*0} \mu^+ \mu^-$



arXiv:2012.13241: $B^+ \rightarrow K^{*+} \mu^+ \mu^-$





arXiv:2107.13428: $B_s^0 \rightarrow \phi \mu^+ \mu^-$



→ New vector coupling?



$B^0 \rightarrow K^{0*} \mu^+ \mu^-$ at CMS

- New results with full Run1+Run2 statistics





CMS

- New results with full Run1+Run2 statistics
 - In agreement with LHCb





Coherent pattern

- Charm loop effects could also cause a shift in C_9



$B^0 \rightarrow K^{*0} \mu^+ \mu^-$: unbinned analysis



- New analysis without q^2 binning
 - Run-1 + 2016
 - Use *all* the information (resonant decays with J/ψ or $\psi(2S)$ are removed)
 - Control long-distance (non-factorisable) QCD effects ($B^0 \rightarrow K^{*0} J/\psi$)
 - **Reduced discrepancy: consistent with SM at 1.8σ (1.4σ global significance)**



$B^0 \rightarrow K^{0*} \mu^+ \mu^-$: unbinned analysis

NEW

- New analysis without q^2 binning
 - Run-1 + 2016-2018
 - Use *all* the information, in full range $0.1 < q^2 < 18 \text{ GeV}^2$
 - Control long-distance (non-factorisable) QCD effects ($B^0 \rightarrow K^{0*} J/\psi$)
- Reduced discrepancy: consistent with SM at 2.1σ (1.5σ global significance)




Ratio of decay rates

$$R_K = \frac{\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(\mu^+ \mu^-))} \bigg/ \frac{\mathcal{B}(B^+ \rightarrow K^+ e^+ e^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(e^+ e^-))}$$

- Theoretically “clean”
- Experimentally
 - Signal yields
 - Backgrounds
 - Electron reconstruction
 - Efficiencies cancel in ratio
 - Belle II: good electron reconstruction
 - LHCb: large B sample

Pre 2022




LHCb, arXiv:2103.11769
Nature Physics 18, (2022) 277-282

Ratio of decay rates

$$R_K = \frac{\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(\mu^+ \mu^-))} \bigg/ \frac{\mathcal{B}(B^+ \rightarrow K^+ e^+ e^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(e^+ e^-))}$$

December 2022

- Theoretically “clean”
- Experimentally
 - Signal yields
 - Backgrounds
 - Electron reconstruction
 - Efficiencies cancel in ratio
 - Belle II: good electron reconstruction
 - LHCb: large B sample



LHCb, arXiv:2212.09152
Phys. Rev. Lett. 131 (2023) 051803

Ratio of decay rates


$$R_K = \frac{\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(\mu^+ \mu^-))} \bigg/ \frac{\mathcal{B}(B^+ \rightarrow K^+ e^+ e^-)}{\mathcal{B}(B^+ \rightarrow K^+ J/\psi(e^+ e^-))}$$

December 2022

- Theoretically “clean”


- Experimentally

- Signal yields
- Backgrounds
- Electron reconstruction
- Efficiencies cancel in ratio
- Belle II: good electron reconstruction
- LHCb: large B sample




LHCb, arXiv:2212.09152
Phys. Rev. Lett. 131 (2023) 051803

Pre 2022: 3.1σ



December 2022



➤ **Tightening electron PID**

- Led to uncovering previously underestimated **peaking backgrounds**
- Estimated from data by inverting mis-id cuts and forming control regions

Outline


- CKM elements
 - $\sin 2\beta$
 - γ
 - Δm_s
 - V_{ub}
- Anomalies
 - $b \rightarrow c \tau \nu$
 - $b \rightarrow s \ell^+ \ell^-$
- Hadron physics
 - Heavy ion programme
 - Spectroscopy
- Prospects
 - Upgrade II

Future Plans

2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035+
	Run III				Run IV								Run V	
LS2					LS3							LS4		
LHCb 40 MHz UPGRADE I	$L = 2 \times 10^{33}$				LHCb Consolidate		$L = 2 \times 10^{33}$ 50 fb^{-1}				LHCb UPGRADE II		$L = 1-2 \times 10^{34}$ 300 fb^{-1}	
ATLAS Phase I Upgr	$L = 2 \times 10^{34}$				ATLAS Phase II UPGRADE		HL-LHC $L = 5 \times 10^{34}$						HL-LHC $L = 5 \times 10^{34}$	
CMS Phase I Upgr	300 fb^{-1}				CMS Phase II UPGRADE								3000 fb^{-1}	
Belle II	$L = 3 \times 10^{35}$				7 ab^{-1}						$L = 6 \times 10^{35}$		50 ab^{-1}	

LHC schedule:


<https://lhc-commissioning.web.cern.ch/schedule/LHC-long-term.htm>




You are here!

LHCb is back


1) More luminosity




2) Better trigger




New detector since 2022 !




VELO (pixel)




Tracker (scintillating fibers with SiPM)



Ring Imaging Cherenkov




First rings in RICH2 during LHC test Oct 2021



Calorimeter & Muon detector (new electronics)

New CALO
frontend and
control boards


MUON Station 2
Hit map



... and beyond!

2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035+
Run III					Run IV							Run V		
LS2						LS3						LS4		
LHCb 40 MHz UPGRADE I	$L = 2 \times 10^{33}$				LHCb Consolidate	$L = 2 \times 10^{33}$ 50 fb^{-1}					LHCb UPGRADE II	$L = 1-2 \times 10^{34}$ 300 fb^{-1}		
ATLAS Phase I Upgr	$L = 2 \times 10^{34}$				ATLAS Phase II UPGRADE			HL-LHC $L = 5 \times 10^{34}$				HL-LHC $L = 5 \times 10^{34}$		
CMS Phase I Upgr	300 fb^{-1}				CMS Phase II UPGRADE							3000 fb^{-1}		
Belle II	$L = 3 \times 10^{35}$				7 ab^{-1}							$L = 6 \times 10^{35}$ 50 ab^{-1}		

<https://lhc-commissioning.web.cern.ch/schedule/LHC-long-term.htm>




Planning for Upgrade II: many analyses stat. limited

Observable	Current LHCb (up to 9 fb^{-1})	Upgrade I	
		(23 fb^{-1})	(50 fb^{-1})
CKM tests			
γ ($B \rightarrow DK$, etc.)	4° [9, 10]	1.5°	1°
ϕ_s ($B_s^0 \rightarrow J/\psi\phi$)	49 mrad [8]	14 mrad	10 mrad
$ V_{ub} / V_{cb} $ ($\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu$)	6% [30]	3%	—
a_{sl}^d ($B^0 \rightarrow D^-\mu^+\nu_\mu$)	36×10^{-4} [34]	8×10^{-4}	5×10^{-4}
a_{sl}^s ($B_s^0 \rightarrow D_s^-\mu^+\nu_\mu$)	33×10^{-4} [35]	10×10^{-4}	7×10^{-4}
Charm			
ΔA_{CP} ($D^0 \rightarrow K^+K^-, \pi^+\pi^-$)	29×10^{-5} [5]	17×10^{-5}	—
A_Γ ($D^0 \rightarrow K^+K^-, \pi^+\pi^-$)	13×10^{-5} [38]	4.3×10^{-5}	—
Δx ($D^0 \rightarrow K_s^0\pi^+\pi^-$)	18×10^{-5} [37]	6.3×10^{-5}	4.1×10^{-5}
Rare Decays			
$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	71% [40, 41]	34%	—
$S_{\mu\mu}$ ($B_s^0 \rightarrow \mu^+\mu^-$)	—	—	—
$A_T^{(2)}$ ($B^0 \rightarrow K^{*0}e^+e^-$)	0.10 [52]	0.060	0.043
A_T^{Im} ($B^0 \rightarrow K^{*0}e^+e^-$)	0.10 [52]	0.060	0.043
$\mathcal{A}_{\phi\gamma}^{\Delta\Gamma}$ ($B_s^0 \rightarrow \phi\gamma$)	$^{+0.41}_{-0.44}$ [51]	0.124	0.083
$S_{\phi\gamma}$ ($B_s^0 \rightarrow \phi\gamma$)	0.32 [51]	0.093	0.062
α_γ ($\Lambda_b^0 \rightarrow \Lambda\gamma$)	$^{+0.17}_{-0.29}$ [53]	0.148	0.097
Lepton Universality Tests			
R_K ($B^+ \rightarrow K^+\ell^+\ell^-$)	0.044 [12]	0.025	0.017
R_{K^*} ($B^0 \rightarrow K^{*0}\ell^+\ell^-$)	0.10 [61]	0.031	0.021
$R(D^*)$ ($B^0 \rightarrow D^{*-}\ell^+\nu_\ell$)	0.026 [62, 64]	0.007	—

Planning for Upgrade II

- Increase instantaneous luminosity to $1.5 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Increase integrated luminosity to 300 fb^{-1}



Planning for Upgrade II: Physics Reach

Observable	Current LHCb (up to 9 fb^{-1})	Upgrade I (23 fb^{-1})	Upgrade I (50 fb^{-1})	Upgrade II (300 fb^{-1})
CKM tests				
γ ($B \rightarrow DK$, etc.)	4° [9, 10]	1.5°	1°	0.35°
ϕ_s ($B_s^0 \rightarrow J/\psi\phi$)	49 mrad [8]	14 mrad	10 mrad	4 mrad
$ V_{ub} / V_{cb} $ ($\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu$)	6% [30]	3%	—	1%
a_{sl}^d ($B^0 \rightarrow D^-\mu^+\nu_\mu$)	36×10^{-4} [34]	8×10^{-4}	5×10^{-4}	2×10^{-4}
a_{sl}^s ($B_s^0 \rightarrow D_s^-\mu^+\nu_\mu$)	33×10^{-4} [35]	10×10^{-4}	7×10^{-4}	3×10^{-4}
Charm				
ΔA_{CP} ($D^0 \rightarrow K^+K^-, \pi^+\pi^-$)	29×10^{-5} [5]	17×10^{-5}	—	3.0×10^{-5}
A_Γ ($D^0 \rightarrow K^+K^-, \pi^+\pi^-$)	13×10^{-5} [38]	4.3×10^{-5}	—	1.0×10^{-5}
Δx ($D^0 \rightarrow K_s^0\pi^+\pi^-$)	18×10^{-5} [37]	6.3×10^{-5}	4.1×10^{-5}	1.6×10^{-5}
Rare Decays				
$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	71% [40, 41]	34%	—	10%
$S_{\mu\mu}$ ($B_s^0 \rightarrow \mu^+\mu^-$)	—	—	—	0.2
$A_T^{(2)}$ ($B^0 \rightarrow K^{*0}e^+e^-$)	0.10 [52]	0.060	0.043	0.016
A_T^{Im} ($B^0 \rightarrow K^{*0}e^+e^-$)	0.10 [52]	0.060	0.043	0.016
$\mathcal{A}_{\phi\gamma}^{\Delta\Gamma}$ ($B_s^0 \rightarrow \phi\gamma$)	$^{+0.41}_{-0.44}$ [51]	0.124	0.083	0.033
$S_{\phi\gamma}$ ($B_s^0 \rightarrow \phi\gamma$)	0.32 [51]	0.093	0.062	0.025
α_γ ($\Lambda_b^0 \rightarrow \Lambda\gamma$)	$^{+0.17}_{-0.29}$ [53]	0.148	0.097	0.038
Lepton Universality Tests				
R_K ($B^+ \rightarrow K^+\ell^+\ell^-$)	0.044 [12]	0.025	0.017	0.007
R_{K^*} ($B^0 \rightarrow K^{*0}\ell^+\ell^-$)	0.10 [61]	0.031	0.021	0.008
$R(D^*)$ ($B^0 \rightarrow D^{*-}\ell^+\nu_\ell$)	0.026 [62, 64]	0.007	—	0.002

Planning for Upgrade II: started in 2017

Expression of Interest

Physics Case

Accelerator Study

Luminosity Scenarios

[LHCC-2017-003](#)

[LHCC-2018-027](#)

[CERN-ACC-2018-038](#)

[LHCb-PUB-2019-001](#)

- **LHCC and CERN Research Board (Sep 2019)**

- "The recommendation to prepare a framework TDR for the LHCb Upgrade-II was endorsed, noting that LHCb is expected to run throughout the HL-LHC era."


- **European Strategy Update (Jun 2020)**

- "The flavour physics programme made possible with the proton collisions delivered by the LHC is very rich, and will be enhanced with the ongoing and proposed future upgrade of the LHCb detector."
- "The full potential of the LHC and the HL-LHC, including the study of flavour physics, should be exploited"

Planning for Upgrade II: Tracking


VELO pixel

- Add Timing
- New RF-foil
- 3D, LGADs, 28nm



UT pixel

- MAPS, radiation tolerant





Magnet Station new



Mighty Tracker

- MAPS pixel and Scintillating fibers



Planning for Upgrade II: PID detectors

RICH1 and RICH 2


- Reduced pixel size
- Add timing information
- SiPM, MCP

TORCH new

- TOF – quartz
- MCP




ECAL



ECAL

- Space & time segmentation
- SPACAL with rad hard crystals
- Timing layer with MCP or Si
- W-Si sampling





Muon

- □ – RWELL for inner regions
- MWPC for outer regions (recycles)





Planning for Upgrade II: Testbeam

- Activities for RICH, VELO, ECAL, MUON
- Lots of opportunities for R&D in coming decade!



Conclusions


- Precision measurements to scrutinize the Standard Model
- Precision measurements reach very high mass scales
- Precision measurements are not yet precise enough
- Lots of opportunities to contribute to R&D




The LHCb Detector



The LHCb Detector



The LHCb Detector



23 sep 2010
Run 79646



19:49:24
Event 143858637





Amplitude interferometry

— Total B ampl

— Total \bar{B} ampl




$\phi = 10^\circ$
 $\Gamma/\Delta m = 1.3$



Constraints on angle γ - with $B^\pm \rightarrow D^0 K^\pm$ and $D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$

- Different yields for B^+ and B^- decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$





$$\Gamma_{B^\pm \rightarrow D[K^\mp \pi^\pm \pi^\pm \pi^\mp]K^\pm} \propto r_{K3\pi}^2 + (r_B^K)^2 + 2r_{K3\pi} r_B^K R_{K3\pi} \cos(\delta_B^K + \delta_{K3\pi} \pm \gamma)$$

Constraints on angle γ - with $B^\pm \rightarrow D^0 K^\pm$ and $D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$

- Different yields for B^+ and B^- decays

- two amplitudes contribute with different relative phase: $V_{ub} = |V_{ub}|e^{-i\gamma}$



$$\Gamma_{B^\pm \rightarrow D[K^\mp \pi^\pm \pi^\pm \pi^\mp]K^\pm} \propto r_{K3\pi}^2 + (r_B^K)^2 + 2r_{K3\pi} r_B^K R_{K3\pi} \cos(\delta_B^K + \delta_{K3\pi} \pm \gamma)$$





$$\gamma = (54.8^{+6.0+0.6+6.7}_{-5.8-0.6-4.3})^\circ$$

(Split in 4 regions of $K^\mp \pi^\pm \pi^\pm \pi^\mp$ Dalitz space:)





$$\begin{aligned} \mathcal{A}_K^1 &= -0.469 \pm 0.088 \pm 0.009 \\ \mathcal{A}_K^2 &= -0.852 \pm 0.077 \pm 0.012 \\ \mathcal{A}_K^3 &= -0.284 \pm 0.080 \pm 0.009 \\ \mathcal{A}_K^4 &= +0.107 \pm 0.083 \pm 0.009 \end{aligned}$$

Precision Δm_s with $B_s^0 \rightarrow D_s^+ \pi^-$


- Legacy “textbook” run-2 measurement
 - “Flavour specific” : final state reveals flavour of the decaying B
 - Precision: 3×10^{-4}
 - “Standard candle” for run-3
-
- 2D mass fit on B_s^0 and D_s^+ mass, followed by decay time fit
 - Detailed study of tagging, decay time resolution and bias




More results: CPV



CC and FCNC



Semileptonic
CC
 $b \rightarrow cl\nu$




"Semileptonic"
FCNC EWP Penguin
 $b \rightarrow sl^+l^-$


R(D*) vs R(D)

- Fit was checked on specific subsamples:


$(\Lambda_b \rightarrow D^0 \mu \mu X)$ enriched




$(\eta \rightarrow \pi^+ \pi^- \pi^0)$ enriched




$(D^* \text{ non-}\mu)$ enriched




$(\phi \rightarrow KK)$ enriched



$(D^* \mu + 3\pi)$ enriched





$(DD \text{ WS-K})$ enriched



$B_s^0 \rightarrow \mu^+ \mu^-$


- Purely leptonic $b \rightarrow s l^+ l^-$



+ $B_s^0 \rightarrow e^+ e^-$ (LHCb, arXiv:[2003.03999](https://arxiv.org/abs/2003.03999))

+ $B_s^0 \rightarrow \tau^+ \tau^-$ (LHCb, arXiv:[1703.02508](https://arxiv.org/abs/1703.02508))

$B_s^0 \rightarrow \mu^+ \mu^-$ (LHCb)



LHCb Coll. [arXiv:2108.09284](https://arxiv.org/abs/2108.09284)

Theory :

$$\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-) = (3.66 \pm 0.14) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-) = (1.03 \pm 0.05) \times 10^{-10}$$

$$\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-) = (3.09^{+0.46+0.15}_{-0.43-0.11}) \times 10^{-9}$$


$$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-) < 2.6 \times 10^{-10}$$

$$\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-\gamma)_{m_{\mu\mu} > 4.9 \text{ GeV}/c^2} < 2.0 \times 10^{-9}$$

Beneke, Bobeth, Szafron, arXiv:1908.07011


$B_{(s)}^0 \rightarrow \mu^+ \mu^-$ (2020)

- Including B^0 :



$B_{(s)}^0 \rightarrow \mu^+ \mu^-$

- Including B^0 :
- NB: new result from CMS at ICHEP not included here



LHCb Coll. [arXiv:2108.09284](https://arxiv.org/abs/2108.09284)

- Relative production of B_s^0 wrt B^0 mesons, f_s/f_d :

f_s/f_d (7 TeV)	$= 0.2390 \pm 0.0076$
f_s/f_d (8 TeV)	$= 0.2385 \pm 0.0075$
f_s/f_d (13 TeV)	$= 0.2539 \pm 0.0079$


f_s/f_d (p_T , 7 TeV)	$= (0.244 \pm 0.008) + ((-10.3 \pm 2.7) \times 10^{-4}) \cdot p_T$
f_s/f_d (p_T , 8 TeV)	$= (0.240 \pm 0.008) + ((-3.4 \pm 2.3) \times 10^{-4}) \cdot p_T$
f_s/f_d (p_T , 13 TeV)	$= (0.263 \pm 0.008) + ((-17.6 \pm 2.1) \times 10^{-4}) \cdot p_T$

(Integrated, p_T [0.5,40] GeV/c, η [2.6,4])

LHCb Coll, arXiv:[2103.06810](https://arxiv.org/abs/2103.06810)



$B_s^0 \rightarrow \mu^+ \mu^-$

Summer 2022



Intermezzo: Effective couplings

- Historical example





$$\frac{G_F}{\sqrt{2}} = \frac{g^2}{8M_W^2}$$



- Both are correct, depending on the energy scale you consider

Intermezzo: Effective couplings

- Historical example



- Analog: Flavour-changing neutral current



Intermezzo: Effective couplings

- Effective coupling can be of various “kinds”

- Vector coupling: C_9
- Axial coupling: C_{10}
- Left-handed coupling (V-A): C_9 - C_{10}
- Right-handed (to quarks): C_9' , C_{10}' , ...
- ...

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{\text{CKM}} \sum_i C_i(\mu) Q_i$$



See e.g. Buras & Fleischer, [hep-ph/9704376](https://arxiv.org/abs/hep-ph/9704376)

Semi-Leptonic Operators (fig. 11f):

$$Q_{9V} = (\bar{s}b)_{V-A}(\bar{\mu}\mu)_V$$

$$Q_{10A} = (\bar{s}b)_{V-A}(\bar{\mu}\mu)_A$$

- Analog: Flavour-changing neutral current



Intermezzo: Effective couplings

- C_7 (photon), C_9 (vector) and C_{10} (axial) couplings hide everywhere:

$$\begin{aligned}
 A_{\perp}^{L,R} &\propto (C_9^{eff} + C_9^{eff'}) \mp (C_{10}^{eff} + C_{10}^{eff'}) \frac{V(q^2)}{m_B + m_{K^*}} + \frac{2m_B}{q^2} (C_7^{eff} + C_7^{eff'}) T_1(q^2) \\
 A_{\parallel}^{L,R} &\propto (C_9^{eff} - C_9^{eff'}) \mp (C_{10}^{eff} - C_{10}^{eff'}) \frac{A_1(q^2)}{m_B + m_{K^*}} + \frac{2m_B}{q^2} (C_7^{eff} - C_7^{eff'}) T_2(q^2) \\
 A_0^{L,R} &\propto (C_9^{eff} - C_9^{eff'}) \mp (C_{10}^{eff} - C_{10}^{eff'}) \times [(m_B^2 - m_{K^*}^2 - q^2)(m_B + m_{K^*} A_1(q^2) - \lambda \frac{A_2(q^2)}{m_B + m_{K^*}})] + \\
 &\quad 2m_B (C_7^{eff} - C_7^{eff'}) [(m_B^2 + 3m_{K^*}^2 - q^2) T_2(q^2) - \frac{\lambda}{m_B^2 - m_{K^*}^2} T_3(q^2)]
 \end{aligned}$$

$$\begin{aligned}
 F_L &= \frac{A_0^2}{A_{\parallel}^2 + A_{\perp}^2 + A_0^2} \\
 S_3 &= \frac{A_{\perp}^{L2} - A_{\parallel}^{L2}}{A_{\perp}^{L2} + A_{\parallel}^{L2} + A_0^{L2}} + L \rightarrow R \\
 S_4 &= \frac{\Re(A_0^{L*} A_{\parallel}^L)}{|A_0^L|^2 |A_{\parallel}^L|^2 + |A_0^L|^2} + L \rightarrow R \\
 S_5 &= \frac{\Re(A_0^{L*} A_{\perp}^L)}{|A_0^L|^2 + |A_{\perp}^L|^2 + |A_0^L|^2} - L \rightarrow R \\
 S_6 &= \frac{\Re(A_{\perp}^{L*} A_{\parallel}^L)}{|A_{\perp}^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} - L \rightarrow R = \frac{4}{3} A_{FB} \\
 S_7 &= \frac{\Im(A_0^{L*} A_{\parallel}^L)}{|A_0^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} + L \rightarrow R \\
 S_8 &= \frac{\Im(A_0^{L*} A_{\perp}^L)}{|A_0^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} + L \rightarrow R \\
 S_9 &= \frac{\Im(A_{\perp}^{L*} A_{\parallel}^L)}{|A_{\perp}^L|^2 + |A_{\parallel}^L|^2 + |A_0^L|^2} - L \rightarrow R
 \end{aligned}$$



$$\begin{aligned}
 \frac{1}{\Gamma} \frac{d^3(\Gamma + \bar{\Gamma})}{d \cos \theta_{\ell} d \cos \theta_K d \phi} &= \frac{9}{32\pi} \left[\frac{3}{4} (1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K + \frac{1}{4} (1 - F_L) \sin^2 \theta_K \cos 2\theta_{\ell} \right. \\
 &\quad \left. - F_L \cos^2 \theta_K \cos 2\theta_{\ell} + \right. \\
 &\quad S_3 \sin^2 \theta_K \sin^2 \theta_{\ell} \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_{\ell} \cos \phi + \\
 &\quad S_5 \sin 2\theta_K \sin \theta_{\ell} \cos \phi + S_6 \sin^2 \theta_K \cos \theta_{\ell} + \\
 &\quad S_7 \sin 2\theta_K \sin \theta_{\ell} \sin \phi + \\
 &\quad \left. S_8 \sin 2\theta_K \sin 2\theta_{\ell} \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_{\ell} \sin 2\phi \right]
 \end{aligned}$$

Coherent pattern

Model independent fits:

- C_9^{NP} deviates from 0 by $>4\sigma$
- Independent fits by many groups favour:
 - $C_9^{\text{NP}} = -1$ or
 - $C_9^{\text{NP}} = -C_{10}^{\text{NP}}$

➤ All measurements (175) agree with a single (simple?) shift...



Wilson coefficient	best fit	pull
$C_9^{bs\mu\mu}$	$-0.82^{+0.14}_{-0.14}$	6.2σ
$C_{10}^{bs\mu\mu}$	$+0.56^{+0.12}_{-0.12}$	4.9σ
$C_9^{rbs\mu\mu}$	$-0.09^{+0.13}_{-0.13}$	0.7σ
$C_{10}^{rbs\mu\mu}$	$+0.01^{+0.10}_{-0.09}$	0.1σ
$C_9^{bs\mu\mu} = C_{10}^{bs\mu\mu}$	$-0.06^{+0.11}_{-0.11}$	0.5σ
$C_9^{bs\mu\mu} = -C_{10}^{bs\mu\mu}$	$-0.43^{+0.07}_{-0.07}$	6.2σ

Similar improvement of fit for both scenario's