





*Physics and AI enhanced
reconstruction at the CEPC*




Manqi Ruan

Yields \sim Xsec * Lumi * Time



- 4 Million Higgs (10 years)
- \sim 1 Giga W (1 year) + 4 Tera Z (2 years)
- Upgradable: Top factory (500 k ttbar)

CEPC Physics study



Chinese Physics C Vol. 43, No. 4 (2019) 043002

Precision Higgs physics at the CEPC*

Fenfen An(安芬芬)^{4,23} Yu Bai(白羽)⁹ Chunhui Chen(陈春晖)²³ Xin Chen(陈新)⁵ Zhenxing Chen(陈振兴)⁸ Joao Guimaraes da Costa¹ Zhenwei Cui(崔振威)³ Yaquan Fang(方亚泉)^{4,6,34,35} Chengdong Fu(付成栋)⁴ Jun Gao(高俊)²³ Yanyan Gao(高艳彦)²² Yunning Gao(高原宁)³ Shaofeng Ge(葛韶峰)^{13,29} Jiayin Gu(顾嘉茵)^{13,29} Fangyi Guo(郭方懿)^{4,4} Jun Guo(郭军)¹⁹ Tao Han(韩涛)^{3,31} Shuang Han(韩爽)⁴ Hongjian He(何红建)^{11,19} Xianke He(何显柯)¹⁶ Xiaogang He(何小刚)^{11,16,20} Jifeng Hu(胡继峰)¹⁶ Shih-Chieh Hsu(徐士杰)³² Shan Jin(金山)⁸ Maoqing Jing(荆茂强)^{3,4} Susmita JyotiShamti³³ Ryuta Kinoshita⁴ Chia-Ming Kuo(郭家铭)³¹ Peizhuo Lai(赖培筑)³¹ Boyang Li(李博扬)³ Congqiao Li(李聪乔)³ Gang Li(李刚)^{4,34,35} Haifeng Li(李海峰)¹² Liang Li(李亮)¹⁹ Shu Li(李淑)^{11,19} Tong Li(李通)¹² Qiang Li(李强)³ Hao Liang(梁浩)^{4,6} Zhijun Liang(梁志均)⁴ Libo Liao(廖立波)⁴ Bo Liu(刘波)^{4,23} Jianbei Liu(刘建北)³ Tao Liu(刘涛)¹⁴ Zhen Liu(刘真)^{28,36,40} Xinchou Lou(娄辛丑)^{4,43,44} Lianliang Ma(马连良)¹² Bruce Mellado^{13,18} Xin Mo(莫欣)⁴ Mila Pandurovic¹⁶ Jianming Qian(钱剑明)^{34,35} Zhaosui Qian(钱卓隼)¹⁹ Nikolaos Rempotis²² Manqi Ruan(阮俊奇)⁴⁰ Alex Schryt³² Liangyou Shan(单连友)³ Jingyuan Shi(史静远)³ Xin Shi(史欣)⁴ Shufang Su(苏淑芳)³² Dayong Wang(王大勇)³ Jun Wang(王隼)⁴ Liantao Wang(王连涛)^{37,39} Yifang Wang(王贻芳)^{4,6} Yuqian Wei(魏或琦)⁴ Yue Xu(许悦)³ Haijun Yang(杨海军)^{30,31} Ying Yang(杨迎)⁴ Weiming Yao(姚伟民)³⁸ Dan Yu(于丹)³ Kaili Zhang(张凯栗)^{4,49} Zhaoru Zhang(张照茹)⁴ Mingrui Zhao(赵明锐)² Xiaohu Zhao(赵祥虎)⁴ Ning Zhou(周宁)¹⁰

¹Department of Modern Physics, University of Science and Technology of China, Anhui 230026, China

²China Institute of Atomic Energy, Beijing 102413, China

³School of Physics, Peking University, Beijing 100871, China

⁴Institute of High Energy Physics, Beijing 100049, China

⁵Department of Engineering Physics, Physics Department, Tsinghua University, Beijing 100084, China

⁶University of Chinese Academy of Science (UCAS), Beijing 100049, China

⁷School of Nuclear Science and Technology, University of South China, Hengyang 421001, China

⁸Department of Physics, Nanjing University, Nanjing 210093, China

⁹Department of Physics, Southeast University, Nanjing 210096, China

¹⁰School of Physics and Astronomy, Shanghai Jiao Tong University, KLPPAC-MoE, SKLJPC, Shanghai 200240, China

¹¹Tung-Dao Lee Institute, Shanghai 200240, China

¹²Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao 266237, China

¹³PRISMA Cluster of Excellence & Mainz Institute of Theoretical Physics, Johannes Gutenberg-Universität Mainz, Mainz 55128, Germany

¹⁴Department of Physics, Hong Kong University of Science and Technology, Hong Kong

¹⁵Kevin IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

¹⁶Vincas Institute of Nuclear Sciences, University of Belgrade, Belgrade 11000, Serbia

¹⁷School of Physics and Institute for Collider Particle Physics, University of the Witwatersrand, Johannesburg 2050, South Africa

Received 9 November 2018, Revised 21 January 2019, Published online 4 March 2019

* Supported by the National Key Program for S&T Research and Development (2016YFA040400); CAS Center for Excellence in Particle Physics; Yifang Wang's Science Studio of the Ten Thousand Talents Project; the CAS-SAFEA International Partnership Program for Creative Research Teams (H21510183); BHEP Innovation Grant (Y454517072); Key Research Program of Frontier Sciences, CAS (SQZDYJ-SW-SL3002); Chinese Academy of Science Special Grant for Large-Scale Scientific Project (11311KYSB0170005); the National Natural Science Foundation of China(11675203); the Hundred Talent Programs of Chinese Academy of Science (Y151540U1); the National 1000 Talents Program of China; Front Research Alliance, LLC (DE-AC02-07CH11359); the NSF(PHY1610074); by the Maryland Center for Fundamental Physics (MCFP); Tsinghua University Initiative Scientific Research Program; and the Beijing Municipal Science and Technology Commission project(Z181100004218003)

- 1) E-mail: fangyi@ihep.ac.cn
- 2) E-mail: jiang@ustc.edu.cn
- 3) E-mail: li.gang@mail.ihep.ac.cn
- 4) E-mail: zhangy@ustc.edu.cn
- 5) E-mail: qiong@ustc.edu.cn
- 6) E-mail: manqi.ruan@ihep.ac.cn
- 7) E-mail: liantao@ustc.edu.cn
- 8) E-mail: zhangk@ihep.ac.cn

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd




Table 2.1: Precision of the main parameters of interests and observables at the CEPC, from Ref. [1] and the references therein, where the results of Higgs are estimated with a data sample of 20 ab^{-1} . The HL-LHC projections of 3000 fb^{-1} data are used for comparison. [2]

Observable	Higgs		W, Z and top		
	HL-LHC projections	CEPC precision	Observable	Current precision	CEPC precision
M_H	20 MeV	3 MeV	M_W	9 MeV	0.5 MeV
Γ_H	20%	1.7%	Γ_W	49 MeV	2 MeV
$\sigma(ZH)$	4.2%	0.26%	Γ_Z	2.3 MeV	0.025 MeV
$B(H \rightarrow bb)$	4.4%	0.14%	M_Z	2.1 MeV	0.1 MeV
$B(H \rightarrow cc)$	-	2.0%	Γ_Z	2.3 MeV	0.025 MeV
$B(H \rightarrow gg)$	-	0.81%	R_b	3×10^{-3}	2×10^{-4}
$B(H \rightarrow WW^*)$	2.8%	0.53%	R_c	1.7×10^{-2}	1×10^{-3}
$B(H \rightarrow ZZ^*)$	2.9%	4.2%	R_μ	2×10^{-3}	1×10^{-4}
$B(H \rightarrow \tau^+\tau^-)$	2.9%	0.42%	R_τ	1.7×10^{-2}	1×10^{-4}
$B(H \rightarrow \gamma\gamma)$	2.6%	3.0%	A_μ	1.5×10^{-2}	3.5×10^{-5}
$B(H \rightarrow \mu^+\mu^-)$	8.2%	6.4%	A_τ	4.3×10^{-3}	7×10^{-5}
$B(H \rightarrow Z\gamma)$	20%	8.5%	A_b	2×10^{-2}	2×10^{-4}
$B(\text{upper}(H \rightarrow \text{inv}))$	2.5%	0.07%	N_ν	2.5×10^{-3}	2×10^{-4}


Scientific Significance quantified by **CEPC physics** studies, via full simulation/phenomenology studies:

- Higgs: Precisions exceed HL-LHC ~ 1 order of magnitude.
- EW: Precision improved from current limit by 1-2 orders.
- Flavor Physics, sensitive to NP of 10 TeV or even higher.
- Sensitive to varies of NP signal.
- ...

White papers +
~300 Journal/AxXiv citables




Higgs & Snowmass White Paper

	240 GeV, 20 ab ⁻¹		360 GeV, 1 ab ⁻¹		
	ZH	vvH	ZH	vvH	eeH
inclusive	0.26%		1.40%	\	\
H→bb	0.14%	1.59%	0.90%	1.10%	4.30%
H→cc	2.02%		8.80%	16%	20%
H→gg	0.81%		3.40%	4.50%	12%
H→WW	0.53%		2.80%	4.40%	6.50%
H→ZZ	4.17%		20%	21%	
H → ττ	0.42%		2.10%	4.20%	7.50%
H → γγ	3.02%		11%	16%	
H → μμ	6.36%		41%	57%	
H → Zγ	8.50%		35%		
Br _{upper} (H → inv.)	0.07%				
Γ _H	1.65%		1.10%		



EW measurements & SMEFT

Observable	current precision	CEPC precision (Stat. Unc.)	CEPC runs	main systematic
Δm_Z	2.1 MeV [37–41]	0.1 MeV (0.005 MeV)	Z threshold	E_{beam}
$\Delta \Gamma_Z$	2.3 MeV [37–41]	0.025 MeV (0.005 MeV)	Z threshold	E_{beam}
Δm_W	9 MeV [42–46]	0.5 MeV (0.35 MeV)	WW threshold	E_{beam}
$\Delta \Gamma_W$	49 MeV [46–49]	2.0 MeV (1.8 MeV)	WW threshold	E_{beam}
Δm_t	0.76 GeV [50]	$\mathcal{O}(10)$ MeV ^a	$t\bar{t}$ threshold	
ΔA_e	4.9×10^{-3} [37, 51–55]	1.5×10^{-5} (1.5×10^{-5})	Z pole ($Z \rightarrow \tau\tau$)	Stat. Unc.
ΔA_μ	0.015 [37, 53]	3.5×10^{-5} (3.0×10^{-5})	Z pole ($Z \rightarrow \mu\mu$)	point-to-point Unc.
ΔA_τ	4.3×10^{-3} [37, 51–55]	7.0×10^{-5} (1.2×10^{-5})	Z pole ($Z \rightarrow \tau\tau$)	tau decay model
ΔA_b	0.02 [37, 56]	20×10^{-5} (3×10^{-5})	Z pole	QCD effects
ΔA_c	0.027 [37, 56]	30×10^{-5} (6×10^{-5})	Z pole	QCD effects
$\Delta \sigma_{had}$	37 pb [37–41]	2 pb (0.05 pb)	Z pole	luminosity
δR_b^0	0.003 [37, 57–61]	0.0002 (5×10^{-6})	Z pole	gluon splitting
δR_c^0	0.017 [37, 57, 62–65]	0.001 (2×10^{-5})	Z pole	gluon splitting
δR_e^0	0.0012 [37–41]	2×10^{-4} (3×10^{-6})	Z pole	E_{beam} and t channel
δR_μ^0	0.002 [37–41]	1×10^{-4} (3×10^{-6})	Z pole	E_{beam}
δR_τ^0	0.017 [37–41]	1×10^{-4} (3×10^{-6})	Z pole	E_{beam}
δN_ν	0.0025 [37, 66]	2×10^{-4} (3×10^{-5})	ZH run ($\nu\nu\gamma$)	Calo energy scale



Flavor Physics White paper

Flavor Physics at CEPC: a General Perspective

Contents

1	Introduction	2
2	Description of the CEPC Facility	6
2.1	Key Collider Features for Flavor Physics	6
2.2	Key Detector Features for Flavor Physics	7
2.3	Simulation Method	16
3	Charged Current Semileptonic and Leptonic b Decays	17
4	Rare/Penguin and Forbidden b Decays	21
4.1	Dilepton Modes	23
4.2	Neutrino Modes	25
4.3	Radiative Modes	27
5	CP Asymmetry in b Decays	27
6	Global Symmetry Tests in Z and b Decays	32
7	Charm and Strange Physics	35
7.1	Null tests with rare charm decays	36
8	τ Physics	36
8.1	LFV τ Decays	37
8.2	LFU Tests in τ Decays	38
8.3	Hadronic τ Decays and Other Opportunities	40
8.4	CPV in hadronic τ decays	41
9	Exclusive Hadronic Z Decays	42
10	Flavor Physics beyond Z Pole	43
10.1	$ V_{cb} $ and W Decays	43
10.2	Top FCNC	45
11	Spectroscopy and Exotics	46
12	Light BSM States from Heavy Flavors	50
12.1	Lepton Sector	51
12.2	Quark Sector	52
13	Summary and Outlook	53




Figure 18: Projected sensitivities of measuring the $b \rightarrow s\tau\tau$ [70], $b \rightarrow s\nu\bar{\nu}$ [34] and $b \rightarrow c\tau\nu$ [35, 62] transitions at the Z pole. The sensitivities at Belle II @ 50 ab^{-1} [6] and LHCb Upgrade II [17, 71] have also been provided as a reference. Note, the LHCb sensitivities are generated by combining the analyses of $\tau^+ \rightarrow \pi^+\pi^-\pi^0\nu$ and $\tau \rightarrow \mu\nu\bar{\nu}$. This plot is adapted from [35].




Figure 21: Illustrative Feynman diagrams for the $B_s \rightarrow \phi\nu\bar{\nu}$ transitions in the SM. **LEFT:** EW penguin diagram. **RIGHT:** EW box diagram.






Figure 22: **LEFT:** Relative precision for measuring the signal strength of $B_s \rightarrow \phi\nu\bar{\nu}$ at Tera-Z, as a function of its BR. **RIGHT:** Constraints on the LEFT coefficients $C_L^{\text{NP}} \equiv C_L - C_L^{\text{SM}}$ and C_R with the measurements of the overall $B_s \rightarrow \phi\nu\bar{\nu}$ decay rate (green band) and the ϕ polarization F_L (orange regions). These plots are taken from [34].

40+ benchmarks + ... Access to NP at 10 TeV or higher



New Physics White paper

	4		5		6
ABSTRACT (TO BE UPDATED)					
The Circular Electron Positron Collider (CEPC) is a large-scale collider facility that can serve as a factory of the Higgs, Z , and W bosons and is upgradable to run at the $t\bar{t}$ threshold. This document describes the latest CEPC nominal operation scenario and particle yields and updates the corresponding physics potential. A new detector concept is also briefly described. This submission is for consideration by the Snowmass process.					
CONTENTS					
Contributors (to be updated)	1				
Abstract (to be updated)	4				
I. Executive Summary (Liantao, Xuai, Manqi, Jia, Zhen, Zhao, Yu...)	8	2. Z exotic decays into Long Lived Particles	38	1. ALPs and new scalar particles	95
II. Introduction(Liantao, Xuai, Manqi, Jia, Zhen, Zhao, Yu ...)	10	H. Top quark exotic decays	41	2. New neutral gauge bosons	97
III. Description of CEPC facility, nominal luminosity and Typical Detector Performance (Manqi)	16	V. Supersymmetry (Tianjun, Lei, Xuai, Da)	42	E. Summary and Discussion	99
A. Key Collider Features	16	A. Light electroweakino searches	43	VIII. Flavor Portal NP(Lingfeng, Xinqiang)	102
B. Key Detector Features	17	B. Light slepton searches	45	A. cLFV processes	104
IV. Exotic Higgs potential and Exotic Higgs/ Z /top decays (Yaquan, Zhao)	26	C. Input from the European Strategy	49	B. Decays of b and c hadrons	105
A. Model-independent Sensitivity to Exotic Higgs decays	26	VI. Dark Matter and Dark Sector (Jia, Xiaoping, Yongchao, Bhupal)	50	C. Light BSM degrees of freedom from flavor transitions	106
B. Exotic Higgs potential	28	A. Lepton portal DM	52	IX. Electroweak phase transition and gravitational waves (Ke-Pan Xie, Sai Wang, Fa Peng Huang, Bruce)	108
C. Higgs exotic decays in supersymmetry	30	B. Interplay of dark particles with neutrinos	53	A. Probing the nature of electroweak phase transition at the colliders	109
D. Exotic Decays via Dark Sector	32	C. Leptophilic DM	54	B. Higgs precision measurements	111
1. Higgs Exotic Decays via Dark Sector	32	D. Dark particles from exotic Z decays	55	C. Higgs exotic decay	114
2. Z Exotic Decays via Dark Sector	34	E. DM with millicharge & the effective theory of DM	57	X. Neutrino Physics (Bhupal, Wei, Yongchao)	117
E. Higgs exotic invisible decays	35	F. Dark particles with electromagnetic form factors	59	A. Motivation	117
F. Decays into Long Lived Particles	36	G. Asymmetric DM	60	B. Connection to Leptogenesis and Dark Matter	118
1. Higgs exotic decays into Long Lived Particles	36	H. Long-lived VLL	62	C. Prospects of heavy neutrinos	121
G. The 95 GeV Higgs boson at the CEPC	39	VII. Long-lived Particle Searches (Liang Li, Ying-nan Mao, Kechen Wang, Zeren Simon Wang)	65	1. heavy neutrinos at the near detector	121
		A. Computation of LLP signal-event rates	66	2. Heavy neutrinos at far detectors	124
		B. Studies with near detectors	70	3. Heavy neutrinos at beam dump experiments	126
		1. Higgs boson decays	71	4. SM Higgs decay $h \rightarrow NN$	129
		2. Z -boson decays	81	5. Prospects of heavy neutrinos in $U(1)$ models	133
		3. Supersymmetry (SUSY)	83	6. Prospects of heavy neutrinos in the LRSM	137
		4. Vector-like leptons with scalar	84	D. Prospects of long-lived doubly-charged scalars in seesaw models	138
		C. Studies with far detectors	85	E. Active-sterile neutrino transition magnetic moments	138
		1. Far detectors at hadron colliders	86	F. Non-standard neutrino interactions	141
		2. Proposed far detectors at lepton colliders	88	XI. More Exotics (Yu, Zuowei)	141
		3. Higgs boson decays	90	A. Axion-like particles	142
		4. Z -boson decays	91	B. Lepton form factors	144
		5. Axion-like particles	92	1. General remarks on μ/e g -2	145
		D. Studies with beam dumps	94	2. μ/e dipole moments in SUSY	148
				3. τ weak-electric dipole moments	149
				C. Emergent Hadron Mass	150
				D. Exotic lepton mass models	152
				XII. Global Fits (Jiayin, Yang, Yong Du)	154
				A. SMEFT global fits	154
				B. 2HDM global fits (Tao Han, Shufang Su, Wei Su, Yongcheng Wu)	157
				C. SUSY global fits	160
				XIII. Conclusion (Liantao, Xuai, Manqi, Jia, Zhen, Zhao, Yu...)	163
				Acknowledgements	164
				References	164

Phase Transition in early Universe, LLP, exotic Higgs decays...



95% C.L. upper limit on selected Higgs Exotic Decay BR



Performance requirements

- To reconstruct all kinds of Physics Object


- Identification & Measurements

- Objects:

- Lepton, Photons, Kaon,
- π^0 , Tau, Lambda, Kshort,
- Heavy flavor hadrons,
- **Jets**
- Missing energy/momentum
- Exotics...

- Massive Four in Standard Model:

- Z & W: ~ 70% goes to a pair of jets
- Higgs: ~90% final state with jets (ZH events)
- Top: $t \rightarrow W + b$



- Requirements:



- Excellent pattern. Reco. & Object id ->

1-1 correspondence reco.

- Larger acceptance...
- Excellent intrinsic resolutions
- Extremely stable...

- Be addressed by state-of-art detector design, technology, and reconstruction algorithm!

CEPC Detector & Reconstruction



Full simulation reconstruction Chain with **Arbor, Jol, etc**

Jet origin id

Hao Liang, Yongfeng Zhu, Yuzhi Che, Yuexin Wang, Huiling Qu, Cen Zhou, etc

PHYSICAL REVIEW LETTERS **132**, 221802 (2024)

Eur. Phys. J. C (2024) 84:152
<https://doi.org/10.1140/epjc/s10052-024-12475-5>

THE EUROPEAN
PHYSICAL JOURNAL C



Regular Article - Experimental Physics

Jet-Origin Identification and Its Application at an Electron-Positron Higgs Factory

Hao Liang^{1,2,*}, Yongfeng Zhu^{3,*}, Yuexin Wang^{1,4}, Yuzhi Che^{1,2}, Manqi Ruan^{1,2,†},
Chen Zhou^{3,‡} and Huilin Qu^{5,§}

¹Institute of High Energy Physics, Chinese Academy of Sciences, 19B Yuquan Road, Shijingshan District, Beijing 100049, China

²University of Chinese Academy of Sciences, 19A Yuquan Road, Shijingshan District, Beijing 100049, China

³State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

⁴China Center of Advanced Science and Technology, Beijing 100190, China

⁵CERN, EP Department, CH-1211 Geneva 23, Switzerland

(Received 16 October 2023; revised 26 April 2024; accepted 1 May 2024; published 31 May 2024)

To enhance the scientific discovery power of high-energy collider experiments, we propose and realize the concept of jet-origin identification that categorizes jets into five quark species (b, c, s, u, d), five antiquarks ($\bar{b}, \bar{c}, \bar{s}, \bar{u}, \bar{d}$), and the gluon. Using state-of-the-art algorithms and simulated $\nu\bar{\nu}H, H \rightarrow jj$ events at 240 GeV center-of-mass energy at the electron-positron Higgs factory, the jet-origin identification simultaneously reaches jet flavor tagging efficiencies ranging from 67% to 92% for bottom, charm, and strange quarks and jet charge flip rates of 7%–24% for all quark species. We apply the jet-origin identification to Higgs rare and exotic decay measurements at the nominal luminosity of the Circular Electron Positron Collider and conclude that the upper limits on the branching ratios of $H \rightarrow s\bar{s}, u\bar{u}, d\bar{d}$ and $H \rightarrow sb, db, uc, ds$ can be determined to 2×10^{-4} to 1×10^{-3} at 95% confidence level. The derived upper limit for $H \rightarrow s\bar{s}$ decay is approximately 3 times the prediction of the standard model.

ParticleNet and its application on CEPC jet flavor tagging

Yongfeng Zhu^{1,a}, Hao Liang^{2,3}, Yuexin Wang^{2,3}, Huilin Qu⁴, Chen Zhou^{1,b}, Manqi Ruan^{2,3,c}

¹State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

²Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

³University of Chinese Academy of Sciences (UCAS), Beijing 100049, China




⁴EP Department, CERN, 1211 Geneva 23, Switzerland

Received: 15 November 2023 / Accepted: 23 January 2024
© The Author(s) 2024

<https://arxiv.org/abs/2310.03440>

<https://arxiv.org/abs/2309.13231>


Geo. & Tools





- **Jet origin identification: 11 categories (5 quarks + 5 anti quarks + gluon)**
- Full Simulated vvH, Higgs to two jets sample at CEPC baseline configuration: CEPC-v4 detector, reconstructed with **Arbor + ParticleNet (Deep Learning Tech.)**
 - Input: measurable information of all reconstructed jet particles (~ 10 float)
 - Output: 10(11)-likelihoods to different categories
- 1 Million samples each, 60/20/20% for training, validation & test

11-dim migration behavior

- Let the jet be identified as the category with highest likelihood:
- Pid: ideal Pid – three categories
 - Lepton identification
 - **Charged Kaon identification**
 - Neutral Kaon identification
- Patterns:
 - ~ Diagonal at quark sector...
 - $P(g \rightarrow q) < P(q \rightarrow g)$...
 - Light jet id...



Performance with different PID scenarios & $H \rightarrow ss$ measurements




Flavor tagging: type that maximize $\{L_q + L_{q\text{-bar}}, L_g\}$

If quark jet: jet charge \sim compare $\{L_q, L_{q\text{-bar}}\}$

Remark: current jet flavor tagging efficiency & jet charge flip rates are projections of the 11-dim arrays produced by Jet origin id

Benchmark analyses: Higgs rare/FCNC



Improved by ~3 times

Improved by 1-2 orders of magnitudes


Presumably... firstly quantified

TABLE I: Summary of background events of $H \rightarrow b\bar{b}/c\bar{c}/gg, Z,$ and W prior to flavor-based event selection, along with the expected upper limits on Higgs decay branching ratios at 95% CL. Expectations are derived based on the background-only hypothesis.

	Bkg. (10^3)			Upper limit (10^{-3})						
	H	Z	W	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$	sb	db	uc	ds
$\nu\bar{\nu}H$	151	20	2.1	0.81	0.95	0.99	0.26	0.27	0.46	0.93
$\mu^+\mu^-H$	50	25	0	2.6	3.0	3.2	0.5	0.6	1.0	3.0
e^+e^-H	26	16	0	4.1	4.6	4.8	0.7	0.9	1.6	4.3
Comb.	-	-	-	0.75	0.91	0.95	0.22	0.23	0.39	0.86

- [28] J. Duarte-Campderros, G. Perez, M. Schlaffer, and A. Soffer. Probing the Higgs–strange-quark coupling at e^+e^- colliders using light-jet flavor tagging. *Phys. Rev. D*, 101(11):115005, 2020.
- [50] Alexander Albert et al. Strange quark as a probe for new physics in the Higgs sector. In *Snowmass 2021*, 3 2022.
- [59] J. de Blas et al. Higgs Boson Studies at Future Particle Colliders. *JHEP*, 01:139, 2020.
- [60] Jorge De Blas, Gauthier Durieux, Christophe Grojean, Jiayin Gu, and Ayan Paul. On the future of Higgs, electroweak and diboson measurements at lepton colliders. *JHEP*, 12:117, 2019.

Recent update at more benchmarks



- From Jet Flavor Tagging to Jet Origin ID (**Preliminary**):

- $\nu\nu H, H \rightarrow c\bar{c}$: 3% \rightarrow 1.7%

19/9/2024 V_{cb} : 0.75% \rightarrow 0.5%

Updated result on $\sin^2 \theta_{eff}^l$ measurement

Table 2. Sensitivity S of different final state particles.


\sqrt{s}/GeV	S of $A_{FB}^{e/\mu}$	S of A_{FB}^d	S of A_{FB}^u	S of A_{FB}^s	S of A_{FB}^c	S of A_{FB}^b
70	0.224	4.396	1.435	4.403	1.445	4.352
75	0.530	5.264	2.598	5.269	2.616	5.237
92	1.644	5.553	4.200	5.553	4.201	5.549
105	0.269	4.597	1.993	4.598	1.994	4.586
115	0.035	3.956	1.091	3.958	1.087	3.942
130	0.027	3.279	0.531	3.280	0.520	3.261

Table 3. Cross section of process $e^+e^- \rightarrow f\bar{f}$ calculated using the ZFITTER package. Values of the fundamental parameters are set as $m_Z = 91.1875 \text{ GeV}$, $m_t = 173.2 \text{ GeV}$, $m_H = 125 \text{ GeV}$, $\alpha_s = 0.118$ and $m_W = 80.38 \text{ GeV}$.

\sqrt{s}/GeV	σ_μ/mb	σ_d/mb	σ_u/mb	σ_s/mb	σ_c/mb	σ_b/mb
70	0.039	0.032	0.066	0.031	0.058	0.028
75	0.039	0.047	0.073	0.046	0.065	0.043
92	1.196	5.366	4.228	5.366	4.222	5.268
105	0.075	0.271	0.231	0.271	0.227	0.265
115	0.042	0.135	0.122	0.135	0.118	0.132
130	0.026	0.071	0.068	0.071	0.066	0.069

Verify the RG behavior... using
~1 month of data taking

Expected statistical uncertainties on $\sin^2 \theta_{eff}^l$ measurement.
(Using one-month data collection, ~ **4e12/24 Z events** at Z pole)







\sqrt{s}	b	c	s
70	1.6×10^{-5}	3.2×10^{-5}	2.2×10^{-5}
75	1.3×10^{-5}	1.8×10^{-5}	1.8×10^{-5}
92	1.6×10^{-6}	2.2×10^{-6}	2.2×10^{-6}
105	1.0×10^{-5}	2.4×10^{-5}	1.4×10^{-5}
115	1.9×10^{-5}	6.8×10^{-5}	2.7×10^{-5}
130	3.9×10^{-5}	2.3×10^{-4}	5.4×10^{-5}


B-charge flip rate: Bs oscillations

Opposite side

- p charged Leptons with impact param.
- p charged Kaons with impact param.
- p charged pions with impact param.
- p protons with impact param. ?




- Using all reco P (exc. Bs decay final state):
- Flip rate ~ 15%, Eff. Tagging power > 40%





Same side

- p charged Kaons with impact param.
- p charged pions with impact param.





V.S. Hadronization models



Fast/Full Simulation

Z → μμ (91.2 GeV)





- Delphes ~ Perfect PFA (1 – 1 correspondence..)

1-1 correspondence between
visible \leftrightarrow reconstructed


=

Confusion Free PFA + Pid

Boson Mass Resolution: Key Per. Para





BMR decomposition @ CDR baseline



- 1st, Ultimate Precision ~ 2.8 with CDR baseline
- 3rd, HCAL
- 2nd, HCAL resolution dominant the uncertainties from intrinsic detector resolution: *need better HCAL*
- 3rd Leading contribution: Confusion from shower Fragments (fake particles), *need better Pattern Reco.*

Preliminary: Identify & veto charged shower fragments using AI



Trained at 12E4 events,

Test & Applied at 4E4 events

score > 0.75
efficiency ~83%
purity ~95%

After frag. Veto

Fake particle originated Confusion reduced by 1 order of magnitude, at nominal vvH, $H \rightarrow gg$ event

>95% of the visible energy preserves 1-1 correspondence







TABLE I. Most probable values of the number/count and total energy of f^\pm before and after removing identified f^\pm and corresponding BMR.



	Counts	Total energy (GeV)	BMR (%)
Before	75	6.34	3.35
After	10	0.68	2.88



... At Boson Mass Resolution ...






Pid



- Next step: to improve the neutral hadron reco & to optimize the detector configuration

Impact of Neutral hadron id: Preliminary



- With 1-1 correspondence:
 - Fast Sim Prediction: BMR: 2.9 \rightarrow 2.6
 - Need excellent CALO + ToF \sim o(10 ps)
 - Assume Low energy neutron can be tamed... still very challenge...
- Strongly Boost the light quark ID.

Summary

- Electron positron Higgs factory: extremely rich physics program requires excellent physics performance
 - Excellent Pattern, reco → high eff/purity & precision reco. of all physics objects
 - Large acceptance, Extremely stable & excellent intrinsic det performance
- AI: the trends & indispensable tool towards these requirements
 - Significantly enhance the physics reach & alters the detector design/optimization
 - Jet Origin ID: 'see' quark & gluon as lepton & photon
 - *...A “game changer” and opens new horizon for precise flavor studies at all future experiments...*
 - PFA: reduces significantly the leading confusion,
 - BMR improved from 3.7% to 2.9%, save ~10% of luminosity for all physics measurements with hadronic events
 - Towards One-to-one correspondence Reco.
- Lots to be explored



One to one correspondence reco. at Higgs factory

A goal should be pursued,
and **we could** achieve it...



Via state-of-art det. Design & technology + AI
enhanced algorithms

Back up

Performance V.S. Jet Kinematics









Performance @ Z and Higgs



- *M10 instead of M11*

BMR: impact on critical measurements



Arbor

Tree topology of particle shower

Ori. Idea from Henri Videau @ ALEPH

Eur. Phys. J. C (2018) 78:426
<https://doi.org/10.1140/epjc/s10052-018-5876-z>

THE EUROPEAN
PHYSICAL JOURNAL C



Special Article - Tools for Experiment and Theory

Reconstruction of physics objects at the Circular Electron Positron Collider with Arbor

Manqi Ruan^{1,a}, Hang Zhao¹, Gang Li¹, Chengdong Fu¹, Zhigang Wang¹, Xinchou Lou^{6,7,8}, Dan Yu^{1,2}, Vincent Boudry², Henri Videau², Vladislav Balagura², Jean-Claude Brient², Peizhu Lai³, Chia-Ming Kuo³, Bo Liu^{1,4}, Fenfen An^{1,4}, Chunhui Chen⁴, Soeren Prell⁴, Bo Li⁵, Imad Laketineh⁵

¹ Institute of High Energy Physics, Beijing, China

² Laboratoire Leprince-Ringuet, Ecole Polytechnique, Palaiseau, France

³ Department of Physics and Center of high energy and high field physics, National Central University, Taoyuan City, Taiwan


⁴ Iowa State University, Ames, USA

⁵ Institut de Physique Nucleaire de Lyon, Lyon, France

⁶ Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China



⁷ Physics Department, University of Texas at Dallas, Richardson, TX, USA

⁸ University of Chinese Academy of Sciences (UCAS), Beijing, China



20 GeV Klong reconstructed @ ILD Calo
Curves indicating expected particle trajectories (from MC-truth)

Validation: Arbor Branch Length Vs MC Truth



Arbor: successfully **tag** sub-shower structure

Samples: Particle gun event at ILD HCAL (readout granularity 1cm^2 & layer thickness 2.65cm)

Length:

Charged MCParticle: spatial distance between generation/end points

Arbor branch: sum of distance between neighboring cells

$Z \rightarrow 2 \text{ muon},$
 $H \rightarrow 2 \text{ b}$
 $\sim 2\%$


$Z \rightarrow 2 \text{ jet},$
 $H \rightarrow 2 \text{ tau}$
 $\sim 5\%$

$ZH \rightarrow 4 \text{ jets}$
 $\sim 50\%$



$Z \rightarrow 2 \text{ muon}$
 $H \rightarrow WW^* \rightarrow eevv$
 $\sim 1\%$



CMS Experiment at LHC, CERN
Data recorded: Thu Jan 1 01:00:00 1970 CEST
Run/Event: 1 / 1201
Lumi section: 13




V.S. Multiplicity




- ...many patterns need further understanding & towards further optimization...*

s-jets: dependency on Leading hadron




Fragmentation comparison


Percentage of b hadrons by Whizard & Herwig



Percentage of c hadrons by Whizard & Herwig



Percentage of s hadrons by Whizard & Herwig



Lots to be studied...