



Contribution ID: 12

Type: Poster

Study of Linearity and Internal Background for LaBr₃(Ce) Gamma-Ray Scintillation Detector

Friday, 25 May 2012 13:31 (0 minutes)

Cerium-doped lanthanum bromide, LaBr₃(Ce), crystal is the latest among the family of scintillation counters and shows some attracting properties for γ spectroscopy that makes it a suitable solution for security, medical, geophysics and high energy physics applications. Among these properties are good density (5.1 g/cm³), brightness (>65000 photon/MeV), time resolution and excellent energy resolution (<3% FWHM at $E_\gamma=662$ keV). LaBr₃(Ce) crystal could become an interesting alternative to HpGe detectors. LaBr₃(Ce) exhibits a proportional light yield response to γ -ray energy. Often this property is hampered at higher energies (>5 MeV) by instantaneous current limitation of PMT: fast and bright can have drawbacks for high energy applications. Very good linearity was seen up to 2 MeV. LaBr₃(Ce) has also relatively high intrinsic radiation background due to naturally occurring ¹³⁸La radioisotope. A good use of LaBr₃(Ce) requires an accurate determination of the self activity, particularly when low background is required or when events are collected at extremely low trigger rates (both situations are the usual standard in the case of underground nuclear astrophysics experiments where we plan to test the LaBr₃(Ce) crystal). Although ¹³⁸La background is entirely below the energy of 1500 keV, additional background was seen in the region between 1500 and 2750 keV. The impact of internal background on energy resolution and linearity is discussed.

Primary author: LAVAGNO, A. (Politecnico di Torino and INFN Torino)

Co-authors: SCARFONE, A. (ISC-CNR and INFN Torino); GERVINO, G. (Università di Torino and INFN Torino)

Presenter: LAVAGNO, A. (Politecnico di Torino and INFN Torino)

Session Classification: Experimental Systems without Accelerators - Poster Session

Track Classification: P7 - Experimental Systems without Accelerators