

Modular Flavour Symmetries

Ivo de Medeiros Varzielas

CFTP, Instituto Superior Técnico, Lisbon

FLASY 2025, 2025/07/04



Summary of data: quark mixing

Wolfenstein parametrisation


$$V_{CKM} \simeq \begin{pmatrix} 1 & \lambda & \lambda^3 \\ -\lambda & 1 & \lambda^2 \\ \lambda^3 & -\lambda^2 & 1 \end{pmatrix}$$

$\lambda \simeq 0.23$ (Sine of the Cabibbo angle)

Summary of data: lepton mixing

Tri-bi-maximal (TBM) mixing

$$V_{PMNS} \simeq \begin{pmatrix} -\sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ \sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \\ \sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \end{pmatrix}$$



TBM history

See talk by Harrison

[https://agenda.infn.it/event/42270/
contributions/263811/](https://agenda.infn.it/event/42270/contributions/263811/)

Non-Abelian Discrete Symmetries (NADS)



NADS

PACK INCLUDES:

x1

NAD'S BRAZILIAN & BIKINI WAX



SPATULAS

x2



PRECISION SHAPING STICK

x1

NAD'S POST WAX
CALMING OIL WIPES

x3



x4

CHEEKY BIKINI SHAPES

Some NADS

$\Delta(6), \Delta(12), \Delta(24), \Delta(27), \Delta(54), \dots$

Serving a Flavour triplet



Multiple Modular?

Multiple Modulus enables


- Multiple stabilizers / fixed points
- Stabilizers outside fundamental domain

Modular Models with residual symmetries!

The framework

IdMV, King, Zhou

<https://arxiv.org/abs/1906.02208>



The framework (paper version)




Figure 1: Illustration of the breaking of $S_4^A \times S_4^B \times S_4^C \rightarrow S_4^D$, identified as the diagonal subgroup, via the VEVs of Φ_{AC} and Φ_{BC} .

Example model with 3 S_4

Field	S_4^A	S_4^B	S_4^C	$2k_A$	$2k_B$	$2k_C$
L	1	1	3	0	0	0
e^c	1	1	1	0	0	-6
μ^c	1	1	1	0	0	-4
τ^c	1	1	1	0	0	-2
N_A^c	1	1	1	-6	0	0
N_B^c	1	1	1	0	-4	0
Φ_{AC}	3	1	3	0	0	0
Φ_{BC}	1	3	3	0	0	0

Example superpotential

$$\begin{aligned}
 w_\ell = & \frac{1}{\Lambda} [L\Phi_{AC} Y_A(\tau_A) N_A^c + L\Phi_{BC} Y_B(\tau_B) N_B^c] H_u \\
 & + [LY_e(\tau_C) e^c + LY_\mu(\tau_C) \mu^c + LY_\tau(\tau_C) \tau^c] H_d \\
 & + \frac{1}{2} M_A(\tau_A) N_A^c N_A^c + \frac{1}{2} M_B(\tau_B) N_B^c N_B^c + M_{AB}(\tau_A, \tau_B) N_A^c N_B^c
 \end{aligned}$$

Example effective superpotential

$$w_{\ell}^{\text{eff}} = \left[\frac{v_{AC}}{\Lambda} LY_A(\tau_A) N_A^c + \frac{v_{BC}}{\Lambda} LY_B(\tau_B) N_B^c \right] H_u + (\dots)$$

Multiple Multiple Modular Models

King, Zhou

<https://arxiv.org/abs/1908.02770> (2 S_4)

King, Zhou

<https://arxiv.org/abs/2103.02633> ($SU(5)$)

IdMV, Lourenço

<https://arxiv.org/abs/2107.04042> (2 A_4)

IdMV, Lourenço

<https://arxiv.org/abs/2206.14869> (2 A_5)

IdMV, King, Levy

<https://arxiv.org/abs/2211.00654> (Littlest Modular)

IdMV, King, Levy

<https://arxiv.org/abs/2309.15901> ($SU(5)$ Littlest Modular)

Littlest Modular Seesaw

Ding, King, Liu, Lu

<https://arxiv.org/abs/1910.03460>

Ding, King, Yao

<https://arxiv.org/abs/2103.16311>

Littlest Modular Seesaw

$$\tau_A = \frac{1}{2} + \frac{i}{2} : \quad Y_{\mathbf{3}'}^{(4)}(\tau_A) = (0, -1, 1), \quad (1)$$


for one of the Dirac mass matrix columns, and

$$\tau_B = \frac{3}{2} + \frac{i}{2} : \quad Y_{\mathbf{3}'}^{(2)}(\tau_B) = (1, 1 - \sqrt{6}, 1 + \sqrt{6}), \quad (2)$$

or

$$\tau_B = -\frac{1}{2} + \frac{i}{2} : \quad Y_{\mathbf{3}'}^{(2)}(\tau_B) = (1, 1 + \sqrt{6}, 1 - \sqrt{6}), \quad (3)$$

LMS Results



Automorphic Forms (multiple moduli)

Ding, Feruglio, Liu

<https://arxiv.org/abs/2010.07952>

Modular Hierarchies

See talk by King

[https://agenda.infn.it/event/42270/
contributions/267075/](https://agenda.infn.it/event/42270/contributions/267075/)

King, King

<https://arxiv.org/abs/2002.00969>

IdMV, King, Levy

<https://arxiv.org/abs/2211.00654>

Hierarchies with weightons

Novichkov, Penedo, Petcov

<https://arxiv.org/abs/2102.07488>

IdMV, Levy, Penedo, Petcov


<https://arxiv.org/abs/2307.14410>

Hierarchies near stabilisers

Grand Unified Theories (GUTs)



GUT × NADS



$SU(5)$ with Littlest Modular

IdMV, King, Levy

<https://arxiv.org/abs/2309.15901> ($SU(5)$ Littlest Modular)

$$Y_u = \begin{pmatrix} y_{uu}\epsilon_T^4 + y'_{uu}\epsilon_F^3\epsilon_T & y_{uc}\epsilon_T^3 + y'_{uc}\epsilon_F^3 & y_{ut}\epsilon_T^2 \\ . & y_{cc}\epsilon_T^2 & y_{ct}\epsilon_T \\ . & . & y_{tt} \end{pmatrix},$$

$$Y_d = \begin{pmatrix} y_{dd}\epsilon_F^3 & y_{ds}\epsilon_F^2\epsilon_T & y_{db}\epsilon_F\epsilon_T^2 \\ 0 & y_{ss}\epsilon_F^2 & y_{sb}\epsilon_F\epsilon_T \\ 0 & 0 & y_{bb}\epsilon_F \end{pmatrix} \quad Y_\ell = \begin{pmatrix} y_{ee}\epsilon_F^3 & 0 & 0 \\ y_{\mu e}\epsilon_F^2\epsilon_T & y_{\mu\mu}\epsilon_F^2 & 0 \\ y_{\tau e}\epsilon_F\epsilon_T^2 & y_{\tau\mu}\epsilon_F\epsilon_T & y_{\tau\tau}\epsilon_F \end{pmatrix}.$$

Pati-Salam GUT with Modular S_3


Cárcamo Hernández, IdMV, King, Vishnudath

<https://arxiv.org/abs/2503.14610>

PS S3 diagrams

$$\begin{array}{c}
 \begin{array}{c} \Phi, \Sigma \quad \sigma_1, \square_1 \\ F \xrightarrow{\gamma_L^{(u)}} \psi_1^c \xrightarrow{m_{\psi_1}} \psi_1 \xrightarrow{\gamma_1^{(u)}} F_1^c \end{array} \quad
 \begin{array}{c} \Phi, \Sigma \quad \sigma_1, \square_1 \\ F \xrightarrow{\gamma_L^{(u)}} \psi_1^c \xrightarrow{m_{\psi_1}} \psi_1 \xrightarrow{\gamma_1^{(u)}} F_2^c \end{array} \quad
 \begin{array}{c} \Phi, \Sigma \quad \sigma_1, \square_1 \\ F \xrightarrow{\gamma_L^{(u)}} \psi_1^c \xrightarrow{m_{\psi_1}} \psi_1 \xrightarrow{\gamma_1^{(u)}} F_3^c \end{array} \\
 \\
 \begin{array}{c} \Phi, \Sigma \quad \sigma_2, \square_2 \\ F \xrightarrow{\gamma_L^{(u)}} \psi_2^c \xrightarrow{m_{\psi_2}} \psi_2 \xrightarrow{\gamma_2^{(u)}} F_1^c \end{array} \quad
 \begin{array}{c} \Phi, \Sigma \quad \sigma_1, \square_1 \\ F \xrightarrow{\gamma_L^{(u)}} \psi_2^c \xrightarrow{m_{\psi_2}} \psi_2 \xrightarrow{\gamma_2^{(u)}} F_2^c \end{array} \\
 \\
 \begin{array}{c} \Phi, \Sigma \\ F_3 \xrightarrow{m_{\psi_3}} F_3^c \end{array}
 \end{array}$$

PS S3 diagrams (paper version)



Modular Flavour Symmetries

- Interesting and ambitious to build models $GUT \times NADS$.
- Modular symmetries can play key roles.
- Viable and predictive models.

For the Portuguese participants



Figure: Tripeiro, Tripperia - Roma, 2025/07