

Measurement of charm suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV via D mesons reconstruction in ALICE

D. Caffarri for the ALICE Collaboration
University of Padova – INFN Sez. di Padova



Outline

- ✧ Parton in-medium energy loss
- ✧ D mesons reconstruction strategy in ALICE
- ✧ D mesons cross section in pp collisions at $\sqrt{s} = 7$ TeV
- ✧ D mesons yields in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV
- ✧ Results on D meson suppression in Pb-Pb collisions

Parton energy loss

- “Hard probes” are produced:
- with hard partonic scatterings
 - in a very short time scale




interaction with the medium




parton energy loss

NUCLEAR MODIFICATION FACTOR

$$R_{AA} = \frac{dN_{AA} / dp_t}{\langle N_{coll} \rangle \times dN_{pp} / dp_t}$$



J. Otwinowski for the ALICE Collaboration,
Quark Matter 2011 proceedings arXiv:1110.2985v1



Energy loss mechanisms

Parton energy loss by:



- medium-induced gluon radiation (inelastic scattering)

$$\Delta E = \Delta E(\varepsilon, L, C_R, m)$$

- collisions with in-medium partons

The total energy loss:

- dominating contribution radiative energy loss
- small fraction due to elastic collisions



P. Arnold, G.D. Moore and L.G. Yaffe, JHEP 0011, 057, (2001)

Heavy quark energy loss

Gluon radiation of heavy quarks is suppressed due to the introduction of a mass term in the heavy quark propagator.

Dead cone effect

Energy loss **colour charge** dependence

$$\langle \Delta E \rangle \propto C_R \quad \begin{array}{l} gg \ C_R = 3 \\ qg \ C_R = 4/3 \end{array}$$




Y.L. Dokshitzer, V.A. Khoze and S.I. Troian, J. Phys. G 17, 1602 (1991);
Y.L. Dokshitzer and D.E. Kharzeev, Phys. Lett. B 519, 199 (2001).

Energy loss **quark mass** dependence

$$\Delta E(\text{light}) > \Delta E(c) > \Delta E(b)$$

$$R_{AA}(\pi) < R_{AA}(D) < R_{AA}(B)$$

Wicks, Gyulassy,
"Last Call for LHC Predictions" workshop, 2007




D mesons reconstruction strategy

D mesons full hadronic reconstruction.



Mass = 1864.80 ± 0.14 MeV

$c\tau = 123 \mu\text{m}$



Mass = 1869.60 ± 0.16 MeV

$c\tau = 311.8 \mu\text{m}$

Invariant mass analysis mainly based on:




Mass = 2010.25 ± 0.14 MeV

- secondary vertex reconstruction
- kaon identification


Secondary vertex reconstruction

Displaced vertex topology:

- tracking and vertexing precision crucial for heavy flavour analysis
- Inner Tracking System with 6 Si layers:
two pixel layers at 3.9 cm and 7 cm



ALICE Collaboration arXiv:1203.2160



Impact parameter resolution $\sim 60\mu\text{m}$ for $p_t = 1$ GeV/c


R_{AA} ingredients....

$$R_{AA} = \frac{dN_{AA} / dp_t}{\langle N_{coll} \rangle \times dN_{pp} / dp_t} = \frac{dN_{AA} / dp_t}{\langle T_{AA} \rangle \times d\sigma_{pp} / dp_t}$$

- Glauber fit: measurement of N_{coll} and T_{AA}
- pp reference at $\sqrt{s} = 2.76$ TeV
- Corrected yields in Pb-Pb collisions

D mesons cross section in pp collisions at 7 TeV

[ALICE Collaboration], JHEP 1201, 128 (2012) [arXiv:1111.1553 [hep-ex]].






D mesons cross section measured in the range $1 < p_t < 24 \text{ GeV}/c$
 pQCD predictions (FONLL and GM-VFNS) compatible with our data




ALICE pp measurement at $\sqrt{s} = 7 \text{ TeV}$ scaled to $\sqrt{s} = 2.76 \text{ TeV}$ using
 FONLL predictions.

R.Averbeck et al., arXiv:1107.3243




Pb-Pb central collisions: signal extraction



0 - 20% CC
 3.1×10^6 events
 $2 < p_t < 16$ GeV/c



40-80% CC
 6.3×10^6 events
 $2 < p_t < 16$ GeV/c




ALICE Collaboration arXiv:1203.2160

Pb-Pb collisions: dN/dp_t

pp scaled reference $\times \langle T_{AA} \rangle$

Pb-Pb yield




ALICE Collaboration arXiv:1203.2160

Pb-Pb collisions: dN/dp_t

pp scaled reference $\times \langle T_{AA} \rangle$


Pb-Pb yield

Indication of suppression



ALICE Collaboration arXiv:1203.2160

D meson R_{AA} vs. p_t




For 0-20% CC suppression is a factor 3-4 for $p_t > 5$ GeV/c.

For 40-80% CC suppression is about a factor 1.5 for $p_t > 5$ GeV/c

D meson R_{AA} vs. collision centrality

5 centrality classes: 0-10%, 10-20%, 20-40%, 40-60%, 60-80%.



For 6-12 GeV/c p_t range the suppression shows a clear trend with centrality. For lower p_t the dependency is less pronounced.

Initial state effects

Nuclear PDFs \neq partons PDFs


- Small $x^{(*)}$ gluons tend to merge together to reach a larger x .
- Initial hard scattering probability reduced at low x and Q^2

Shadowing effect computed with pQCD calculation with CTEQ6M PDFs and EPS09 NLO parametrization.

The strong suppression observed is likely to be a final state effect

(*) momentum fraction of the nucleon carried by the parton

ALICE Collaboration arXiv:1203.2160



R_{AA} of D mesons averaged using statistical errors as weights.




$R_{AA}(\pi) < R_{AA}(D) < R_{AA}(B)$?

ALICE Collaboration arXiv:1203.2160
CMS Collaboration arXiv:1201.5069

ALICE charged hadrons R_{AA} in the centrality class 0-20% (*).

Displaced J/ψ from B decays measured by CMS in 0-20%



$R_{AA}(\pi) < R_{AA}(D)$?

There is an indication of saying yes.
Not conclusive.


$R_{AA}(D) < R_{AA}(B)$?

Different p_t range, not possible to conclude.

(*) ALICE preliminary results showed that charged pion R_{AA} coincides with charged hadron R_{AA} for $p_t > 5$ GeV/c



Comparison with energy loss models



ALICE Collaboration arXiv:1203.2160

Comparison with energy loss models

(I) Radiative energy loss
with in-medium D
meson dissociation


(II) Radiative + collisional
energy loss in WHDG

(VII) CUJET1.0

approaches describe
reasonably well at the
same time charm and
light-flavour suppression.

(III) AdS/CFT approach seems to underestimate the D mesons R_{AA}

ALICE Collaboration arXiv:1203.2160



Conclusions



- ✧ First measurement of the **direct charm suppression** in central heavy-ion collisions: arXiv:1203.2160
- ✧ The D meson Nuclear Modification Factor (R_{AA}) has been studied as a function of p_t and of centrality of the collisions.
The suppression measured is a factor ~ 4 for D with $p_t > 5$ GeV/c
The effect is decreasing from central to peripheral collisions.
- ✧ Comparison with theoretical calculations suggests a final state effect, due to **parton energy loss**.
A p-Pb run will take place this year to study nuclear shadowing at the LHC.
- ✧ The mass hierarchy in the energy loss has been tested and no strong conclusion can be drawn so far.
There are indications that **$R_{AA}(\pi) < R_{AA}(D) < R_{AA}(B)$ could be valid.**

Back up slides

Heavy quark energy loss

Gluon radiation of heavy quarks is suppressed due to the introduction of a mass term in the heavy quark propagator.

Dead cone effect

Energy distribution of the radiated gluons

$$\omega \frac{dI_{rad,Q}}{d\omega} = \omega \frac{dI_{rad}}{d\omega} \cdot \left(1 + \frac{\theta_0^2}{\theta^2}\right)^{-2}, \quad \theta_0 = \frac{M}{E} = \frac{1}{\gamma}$$



Y.L. Dokshitzer, V.A. Khoze and S.I. Troian, J. Phys. G 17, 1602 (1991);
Y.L. Dokshitzer and D.E. Kharzeev, Phys. Lett. B 519, 199 (2001).

Energy loss **colour charge** dependence


$$\langle \Delta E \rangle \propto C_R \quad \begin{array}{l} gg \ C_R = 3 \\ qg \ C_R = 4/3 \end{array}$$

Energy loss **quark mass** dependence

$$\Delta E(\text{light}) > \Delta E(c) > \Delta E(b) \rightarrow R_{AA}(\pi) < R_{AA}(D) < R_{AA}(B)$$


Heavy quarks E. loss: some predictions

- Energy loss based predictions (*):
- factor 3-5 suppression for D mesons
 - smaller suppression for B mesons



Wicks, Gyulassy,
 “Last Call for LHC Predictions” workshop, 2007

Pb-Pb collisions at $\sqrt{s} = 5.5$ TeV



Armesto, et al. PRD71 (2005) 014003


(*) not up to date predictions.
 New predictions at the end...

A Large Ion Collider Experiment - ALICE

Trigger:
Silicon Pixel
Detector (SPD),
V0

Tracking:
Inner Tracking
System (ITS),
Time Projection
Chamber (TPC)

PID:
Time Of Flight
(TOF), TPC



Centrality measurement

Collision geometry \rightarrow number of participating nucleons : $N_{part} = 2A - N_{spec}$


Centrality measurement in ALICE:

- use Zero Degree Calorimeter data
- Glauber Fit



Based on fit on multiplicity distributions of:

- VZERO amplitude
- SPD hits
- TPC tracks





Measurement of N_{coll} and $T_{AA} \rightarrow$ ingredients for the R_{AA}

Particle Identification - PID

Conservative PID strategy used to identify the kaon candidates.

Kaons are identified via:

- the energy loss deposit in the TPC ($0.6 < p < 0.8$ GeV/c 2σ cut)
- the velocity measurement in the TOF ($p < 2$ GeV/c 3σ cut)




Keep the signal loss as small as possible


Background reduction by a factor 3 for central Pb-Pb collisions.

LHC as heavy flavour factory

[ALICE Collaboration], JHEP 1201, 128 (2012) [arXiv:1111.1553 [hep-ex]].



ALICE D mesons measurements in pp collisions at 2.76 and 7 TeV used to compute the total charm production cross section



Good agreement with NLO calculation

Increase of a factor ~ 7 with respect to PHENIX and STAR





pp scaled reference at $\sqrt{s} = 2.76$ TeV

ALICE pp measurement at $\sqrt{s} = 7$ TeV scaled to $\sqrt{s} = 2.76$ TeV using FONLL predictions.

R.Averbeck et al., arXiv:1107.3243

D^0 cross section measurement at $\sqrt{s} = 2.76$ TeV (3 days of data taking)


As test the exercise was done scaling at 1.96 TeV and compared to CDF data.





pp scaled reference at $\sqrt{s} = 2.76$ TeV

ALICE pp measurement at $\sqrt{s} = 7$ TeV scaled to $\sqrt{s} = 2.76$ TeV using FONLL predictions.


R.Averbeck et al., arXiv:1107.3243



D^0 cross section measurement at $\sqrt{s} = 2.76$ TeV (3 days of data taking)




D⁰ Signals in peripheral



40-80% CC
 6.3×10^6 events
 $2 < p_t < 16$ GeV/c
 (hint of signal in 1-2)

Efficiencies

Efficiencies are computed using HIJING PbPb Monte Carlo simulation with embedded PYTHIA cc events



ALICE Collaboration arXiv:1203.2160

Feed down correction and beauty energy loss hypothesis

Beauty feed down:

Monte Carlo method based on FONLL predictions.

Subtraction to the D^0 raw yield the expected secondary raw yields.


Beauty energy loss:

Hypothesis on the energy loss of beauty quarks is adopted.

Central value: $R_{AA}^{\text{feed-down}} = R_{AA}^{\text{prompt}}$

Hypothesis $0.3 < R_{AA}^{\text{feed-down}} / R_{AA}^{\text{prompt}} < 3$

(no correction applied,
a systematic uncertainty added)





Feed down correction

Beauty feed down:

Monte Carlo method based on FONLL predictions.

Subtraction to the D^0 raw yield the expected secondary raw yields.



Systematic uncertainties: 0-20 % CC




| Particle | | D ⁰ | |
|------------------|--|------------------|--------------|
| 0-20% centrality | p_t interval (GeV/c) | 2-3 | 12-16 |
| | | Yield extraction | 8% |
| | Tracking efficiency | 10% | 10% |
| | Cut efficiency | 13% | 10% |
| | PID efficiency | +15% -5% | 5% |
| | MC p_t shape | 4% | 3% |
| | FONLL feed-down corr. | +2% -14% | +6% -8% |
| | $R_{AA}^{\text{feed-down}}/R_{AA}^{\text{prompt}}$ (Eq. (3)) | +4% -10% | +14% -27% |
| | BR | 1.3% | |

| Particle | | D ⁰ | |
|------------------|--|-------------------------|--------------|
| 0-20% centrality | p_t interval (GeV/c) | 2-3 | 12-16 |
| | | Data syst. pp and Pb-Pb | +33% -41% |
| | Data syst. in Pb-Pb | +26% -22% | +22% -22% |
| | Data syst. in pp | 17% | 17% |
| | \sqrt{s} -scaling of the pp ref. | +10% -31% | +5% -6% |
| | Feed-down subtraction | +15% -14% | +16% -29% |
| | FONLL feed-down corr. | +12% -2% | +1% -2% |
| | $R_{AA}^{\text{feed-down}}/R_{AA}^{\text{prompt}}$ (Eq. (3)) | +4% -10% | +14% -27% |

ALICE Collaboration arXiv:1203.2160

Open heavy flavour comparison


ALICE D^0 and HF electrons measurements
 CMS displaced J/ψ from B decays



Energy loss models (I)

Path integral approach scattering probability expansion

- pQCD approaches
- Hard parton traversing the medium interacts with various scattering centres.
- Initial parton = lower energy parton + radiated gluon



BDMPS – ASW ⁽¹⁾ multiple-soft bremsstrahlung gluon radiation

DGLV ⁽²⁾ starts from a single-hard radiation spectrum

DGLV + collisional energy loss and path length fluctuations → WHDG ⁽³⁾

$$\hat{q} = \frac{\langle k_{\perp}^2 \rangle}{\lambda}$$

(1) R. Baier, Y. Dokshitzer, A. Mueller, S. Peigné and D. Shiff, Nucl. Phys. B 483, 291 (1997).

C.A. Salgado and U.A. Weidmann, Phys. Rev. Lett. 89, 092303 (2002).

(2) M. Djordjevic, M. Gyulassy, Nucl. Phys. A 733 265 (2004)

(3) S. Wicks, W. Horowitz, M. Djordjevic and M. Gyulassy, Nucl. Phys. A 784, 426 (2007).

AdS/CFT

Weakly coupled gravity theories → four-dimensional gauge theories

Possible determination of medium viscosity, transport coefficient, heavy quark diffusion coefficient

C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, JHEP 0607, 013(2006)

Energy loss models (II)

Transport models

- Interaction of the charm quark with the plasma: drag and diffusion forces that act on the quark
- Charm quark in the QGP \rightarrow treated with Brownian approximation
- Charm quark density \rightarrow Boltzmann equation

A. Beraudo, A. De Pace, W.M. Alberico, A. Molinari. arXiv:0902.0741v2 [hep-ph]
J. Uphoff, O. Fochler, Z. Xu and C. Greiner, arXiv:1112.1559 [hep-ph].


... and outlook

Strong interaction of charm with the medium...
and what about thermalization with the medium?

ALICE first preliminary measurement on charm flow...

The first point suggests a that charm flow is different than zero (1.8σ)

2011 statistics will allow to improve the measurement and to study charm quark thermalization.



Flow measurements

