

# Gauge theory meets cosmology

Bologna, October, 23 2024

---

Based on:

2408.03243 [hep-th]


with M. Bianchi & J.F. Morales



# Outline

- Introduction & Motivation
- Review of FLRW cosmologies
- Linearized perturbations of FLRW universes
- Cosmological perturbations within  $\Lambda$ CDM
- Seiberg-Witten/Cosmology correspondence
- Concluding Remarks


# The Early Universe...




CMB




COBE  
(1992)



WMAP  
(2003)



Planck  
(2013)



# The Late Universe...

BAO

+



SN Ia




$\Lambda$ CDM model




The «cheapest» description BUT...

# Early/Late Universe puzzles appear...



[image credit: Krzysztof Bolejko]



Douspis et al.

## Early & Late Universe



# Why cosmological perturbations?

A HARD question because...



- Different sorts of perturbations are **intrinsically coupled** and analytic treatments are beyond reach...

A RELEVANT question because...



- The **spectrum** of cosmological perturbations crucially depends on the details of a model!
- It may shed a light on **refined models for the early & late universe** beyond  $\Lambda$ CDM

Novel «semi-analytic» treatments?



Insights from String Theory tools?

**Today:**

Motivate & introduce the **SW/cosmology** dictionary

# FLRW cosmologies revisited

$$ds_4^2 = -dt^2 + a(t)^2 ds_{\mathcal{M}_3}^2,$$

$\mathbb{R}^3, S^3$  or  $\mathbb{H}^3$ .

For perfect  
ISOTROPIC fluids...

$$T_{\mu\nu} = (\rho + p) u_\mu u_\nu + p g_{\mu\nu}$$

diag( $\rho, p, p, p$ )

$$p = w\rho$$

Eqn of state

Einstein equations...

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = M_{\text{Pl}}^{-2}T_{\mu\nu}$$


$$a(t) \sim t^{\frac{2}{3(1+w)}}$$

# A «new» time variable...

$$ds_4^2 = -\frac{da^2}{a^2 b(a)^2} + a^2 ds_{\mathcal{M}_3}^2,$$

Use **time reparametrization invariance!!!**

$$dt = \frac{da}{ab(a)}$$



In this way we get **simpler** equations to solve...



# Perfect fluid dynamics

$$\rho = 3(M_{\text{Pl}} H_0)^2 a^{-n} \quad , \quad p = 3w (M_{\text{Pl}} H_0)^2 a^{-n} \quad , \quad b^2 = H_0^2 a^{-n} \quad , \quad n = 3(1 + w)$$

Fluid Type	Symbol	$w$	$n$	$b(a)$	$\eta(a)$
Vacuum	$\Lambda$	$-1$	$0$	$1$	$-1/a$
Strings, Curvature	$\sigma, \kappa$	$-\frac{1}{3}$	$2$	$a^{-1}$	$\log(a)$
Matter	$m$	$0$	$3$	$a^{-3/2}$	$2a^{1/2}$
Radiation	$\gamma$	$\frac{1}{3}$	$4$	$a^{-2}$	$a$
Stiff	$s$	$1$	$6$	$a^{-3}$	$a^2/2$

For the «full» universe we have...


$$b^2 = H_0^2 (\Omega_\gamma a^{-4} + \Omega_m a^{-3} + \Omega_\Lambda + \Omega_\kappa a^{-2})$$

$$\rho(a) = 3(M_{\text{Pl}} H_0)^2 (\Omega_\Lambda + \Omega_m a^{-3} + \Omega_\gamma a^{-4}),$$

$$p(a) = 3(M_{\text{Pl}} H_0)^2 (-\Omega_\Lambda + \frac{1}{3}\Omega_\gamma a^{-4})$$


$$q(a) \equiv -\frac{\ddot{a}a}{\dot{a}^2} \rightarrow \text{Deceleration parameter}$$

$$= \frac{\Omega_\gamma a^{-4} + \frac{1}{2}\Omega_m a^{-3} - \Omega_\Lambda}{\Omega_\gamma a^{-4} + \Omega_m a^{-3} - \Omega_\kappa a^{-2} + \Omega_\Lambda}$$



# What's the (cosmic) time?

$$b(a) = \frac{\dot{a}}{a} = H(a)$$



$$H_0 t = \int \frac{da}{\sqrt{\Omega_\Lambda a^2 + \Omega_\kappa + \Omega_m a^{-1} + \Omega_\gamma a^{-2}}} = \int \frac{ada}{\sqrt{\Omega_\Lambda a^4 + \Omega_\kappa a^2 + \Omega_m a + \Omega_\gamma}}$$

# Review of cosmological perturbations

[Mukhanov]

**Perturbed metric:**

$$g_{\mu\nu} = \underbrace{g_{\mu\nu}^{(0)}}_{\text{hom. + iso.}} + \delta g_{\mu\nu}$$

**Small perturbations**

**Gauge invariant** scalar perturbations may be parametrized as ...

$$ds_4^2|_S \stackrel{L}{=} - (1 + 2\Phi(a, \mathbf{x})) \frac{da^2}{a^2 b(a)^2} + a^2 (1 - 2\Psi(a, \mathbf{x})) ds_{\mathcal{M}_3}^2$$

**Isotropy!**

$$\Psi = \Phi$$

After **Fourier transformation...**

$$\Phi(a, \mathbf{x}) = e^{i\mathbf{k}\mathbf{x}} \Phi(a) \quad \Rightarrow \quad \Delta \Phi = -k^2 \Phi$$


**ODE for time evolution!**

After taking ODE into its **canonical form**...

$$\Psi''(a) + Q(a)\Psi(a) = 0$$

Rational function of time!

Related to  $\Phi$




$N$	Type	Components
3	Hypergeometric	$\Lambda \kappa, \Lambda m, \kappa m, \kappa \gamma$
4	Heun	$\Lambda \gamma, m\gamma, \Lambda \kappa \gamma$
5	Gen. Heun	$\kappa m \Lambda, \kappa m \gamma$
7	Gen. Heun	$\Lambda m \gamma, \Lambda \kappa m \gamma$



Numerical solution for **radiation & matter** (Heun eqn)

# SW/cosmology correspondence

[Hanany, Witten 1996]



# From classical to quantum geometry...

$$(x, z) \in \mathbb{C}^2 : \quad P_0(x)z^2 - P_1(x)z + qP_2(x) = 0 \quad \text{(Classical SW curve)}$$

«Quantum»

$$x = -z\partial_z$$

Quadratic fct's of  
 $m_i \quad q$

$$\left[ P_0(-z\partial_z + \frac{1}{2}) - P_1(-z\partial_z)z^{-1} + qP_2(-z\partial_z - \frac{1}{2})z^{-2} \right] W(z) = 0$$


with

$$W(z) = z^{1 - \frac{m_3 + m_4}{2}} (1 - z)^{-\frac{m_1 + m_2 + 1}{2}} (z - q)^{\frac{m_3 + m_4 - 1}{2}} \Psi(z)$$

# A complete example: radiation & matter

The associated **Schrodinger-like** equation is specified by

$$Q(a) = \frac{64a^2\zeta k^2 (3a^2 + 7a\zeta + 4\zeta^2) - 3 (189a^4 + 924a^3\zeta + 1820a^2\zeta^2 + 1600a\zeta^3 + 512\zeta^4)}{48a^2(a+\zeta)^2(3a+4\zeta)^2}$$


$$\zeta = \frac{\Omega_\gamma}{\Omega_m}$$

which fits the **SW** induced Heun equation, once the **dictionary** is fixed...

$$z = -\zeta a^{-1}, \quad q = \frac{3}{4}, \quad u = \frac{4\hat{k}^2\zeta^2}{3} + \frac{33}{16}$$
$$m_1 = \frac{7}{4}, \quad m_2 = -\frac{5}{4}, \quad m_{3,4} = 1 \pm \frac{1}{12} \sqrt{225 - 64\hat{k}^2\zeta}$$



# Conclusions & Outlook

- Studying the dynamical evolution of cosmological perturbations is an important **challenge** for cosmology
- We can tackle the problem by using tools borrowed from SUSY QFT's
- This analysis offers the possibility of a «semi-analytic» treatment (important for **parametric** control!)
- In our setup it would be interesting to consider more involved scenarios, like **inflationary** dynamics, **early dark energy**, **quintessence**...



Hopefully more to come, so stay tuned...

Thank you for your attention!

