


Studies on alphas  
Semi-Experimental Toy-MC


(David(e))

# Toy MC


- Let's suppose we can identify  $^{222}\text{Rn}$  and  $^{218}\text{Po}$  (thanks to David Marques, we can)
- I simulated  $^{222}\text{Rn}$  with an average time difference of 300 s and their decays with a lifetime (no half-life) of 250 s, with 100% efficiency in detecting both  $^{222}\text{Rn}$  and  $^{218}\text{Po}$



time difference (in tens of seconds)  
between two consecutive  $^{222}\text{Rn}$



time difference (in tens of seconds)  
between a  $^{222}\text{Rn}$  and its own  $^{218}\text{Po}$




time difference (in tens of seconds)  
between to a  $^{222}\text{Rn}$  and the first  
 $^{218}\text{Po}$  found


The “biased” association shortens the visible lifetime from 250 s to 136 s

# Toy MC


- Let's suppose we have a 30% efficiency in detecting both  $^{222}\text{Rn}$  and  $^{218}\text{Po}$  (e.g. mainly for geometrical reasons);
- According to toyMC the visible lifetime increases



time difference (in tens of seconds)  
between two consecutive  $^{222}\text{Rn}$



time difference (in tens of seconds)  
between a  $^{222}\text{Rn}$  and its own  $^{218}\text{Po}$




time difference (in tens of seconds)  
between to a  $^{222}\text{Rn}$  and the first  
 $^{218}\text{Po}$  found

Assuming a 30% efficiency, the visible lifetime is 200 s

# Toy MC

- Let's suppose we can identify  $^{222}\text{Rn}$  and  $^{218}\text{Po}$  (thanks to David Marques, we can)
- I simulated  $^{222}\text{Rn}$  with an average time difference of 30 s and their decays with a lifetime (no half-life) of 250 s






time difference (in tens of seconds)  
between to a  $^{222}\text{Rn}$  and the first  $^{218}\text{Po}$  found

I found a double exp, where at the beginning I got 50 sec (probably due to the wrong pairing Rn-Po) and in the second part 250 sec probably due to the right association when a second radon doesn't arrive in time

# ToyMC: Test I

- Made some tests to check possible bugs
- With very low radon (1 parent each 1e4 sec)




- Found right tau (250 sec) independently from the efficiency

# ToyMC: Test II

---


- Made some tests to check possible bugs
- With very high radon (1 parent each 10 sec, i.e. the minimum possible for my Toy).




# Radon decrease

- Behavior of number of alphas after filter insertion

No cut on alpha lengths



35mm < alpha lengths < 70mm (2D)



- Filter effect clearly visible, decrease is faster the Rn lifetime (130h)