

# The NA62 experiment at CERN

Francesco Brizioli  
(CERN and INFN Perugia)  
[francesco.brizioli@cern.ch](mailto:francesco.brizioli@cern.ch)




CERN, December 8<sup>th</sup>, 2024

# The NA62 experiment at CERN



~200 collaborators from ~30 institutions.



Photograph: Maximilien Brice © 2008 CERN

- Primary goal: measurement of  $\mathcal{B}(K^+ \rightarrow \pi^+ \bar{\nu}\bar{\nu})$
- New Technique:  $K^+$  decay-in-flight
- Results: [\[PLB 791 \(2019\) 156\]](#) [\[JHEP 11 \(2020\) 042\]](#) [\[JHEP 06 \(2021\) 093\]](#)
- Broader physics programme:
  - Precision measurements of kaon and pion decays
  - HNL and LNV/LFV searches in kaon decays
  - Hidden Sector searches with kaons and in dump mode
- Data taking
  - 2016 Commissioning + Physics run (45 days).
  - 2017 Physics run (160 days).
  - 2018 Physics run (217 days).
  - 2021 Physics run (85 days [10 beam dump]).
  - 2022 Physics run (215 days).
  - 2023 Physics run (150 days [10 beam dump]).
  - 2024 Physics run (204 days [12 beam dump, 7 low int.]).


New result!




# $K \rightarrow \pi \nu \bar{\nu}$ : Precision test of the SM



SM: Z-penguin & box diagrams



- $\mathcal{B}(K \rightarrow \pi \nu \bar{\nu})$  highly suppressed in SM
  - GIM mechanism & maximum CKM suppression  $s \rightarrow d$  transition:  $\sim \frac{m_t^2}{m_W^2} |V_{ts}^* V_{td}|$
  - Theoretically clean  $\Rightarrow$  high precision SM predictions
    - Dominated by short distance contributions.
    - Hadronic matrix element extracted from  $\mathcal{B}(K \rightarrow \pi l \nu)$  decays via isospin rotation.




Decay Mode BR	SM [ <a href="#">Buras et al. EPJC 82 (2022) 7, 615</a> ]	SM [ <a href="#">D'Ambrosio et al. JHEP 09 (2022) 148</a> ]	Experimental Status
$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	$(8.60 \pm 0.42) \times 10^{-11}$	$(7.86 \pm 0.61) \times 10^{-11}$	$(10.6^{+4.1}_{-3.5}) \times 10^{-11}$ (NA62)
$\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})$	$(2.94 \pm 0.15) \times 10^{-11}$	$(2.68 \pm 0.30) \times 10^{-11}$	$< 2 \times 10^{-9}$ (KOTO)

# $K \rightarrow \pi \nu \bar{\nu}$ : Beyond the SM

- Correlations between BSM contributions to BRs of  $K^+$  and  $K_L$  modes [[JHEP 11 \(2015\) 166](#)].
  - Must measure both to discriminate between BSM scenarios.
- Correlations with other observables ( $\varepsilon'/\varepsilon$ ,  $\Delta M_B$ , B-decays)  
[\[JHEP 12 \(2020\) 097\]](#)[\[PLB 809 \(2020\) 135769\]](#).
- Leptoquarks [[EPJC 82 \(2022\) 4, 320](#)], Interplay between CC and FCNC [[JHEP 07 \(2023\) 029](#)], NP in neutrino sector [[EPJC 84 \(2024\) 7, 680](#)] and additional scalar/tensor contributions [[JHEP 12 \(2020\) 186](#)][\[arXiv:2405.06742\]](#) ...
- **Green:** CKM-like flavour structure
  - Models with Minimal Flavour Violation
- **Blue:** new flavour-violating interactions where LH or RH currents dominate
  - $Z'$  models with pure LH/RH couplings
- **Red:** general NP models without above constraints
- **Grossman-Nir Bound:** model-independent relation  
[\[PLB 398 \(1997\) 163-168\]](#)

$$\frac{\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})}{\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})} \frac{\tau_{K^+}}{\tau_{K_L}} \lesssim 1$$

$$\Rightarrow \mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \lesssim 4.3 \cdot \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$$




# $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ at NA62



## NA62 Strategy:

- Tag  $K^+$  and measure momentum.
- Identify  $\pi^+$  and measure momentum.
- Match  $K^+$  and  $\pi^+$  in time & form vertex.
  - Determine  $m_{miss}^2 = (P_K - P_\pi)^2$
  - Reject any additional activity.



## NA62 Performance Keystones:


- $\mathcal{O}(100)\text{ps}$  timing between detectors
- $\mathcal{O}(10^4)$  background suppression from kinematics
- $> 10^7$  muon rejection
- $> 10^7$  rejection of  $\pi^0$  from  $K^+ \rightarrow \pi^+ \pi^0$  decays

Decay mode	Branching Ratio [PDG]
$K^+ \rightarrow \mu^+ \nu_\mu$	$(63.56 \pm 0.11)\%$
$K^+ \rightarrow \pi^+ \pi^0$	$(20.67 \pm 0.08)\%$
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	$(5.583 \pm 0.024)\%$
$K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$	$(4.247 \pm 0.024) \times 10^{-5}$

$$K^+ \rightarrow \pi^+ \nu \bar{\nu} \approx 10^{-10}$$

# NA62 beamline & detector

[JINST 12 (2017) 05, P05025]




- Designed & optimised for study of  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ :
  - Particle tracking: beam particle (GTK) & downstream tracks (STRAW)
  - PID:  $K^+$  - KTAG,  $\pi^+$  - RICH, Calorimeters (LKr, MUV1,2), MUV3 ( $\mu$  detector)
  - Comprehensive veto systems: CHANTI (beam interactions), LAV, LKr, IRC, SAC ( $\gamma$ )

# Kinematic constraints & signal regions




$$m_{miss}^2 = (P_{K^+} - P_{\pi^+})^2$$



$\mathcal{O}(10^4)$  background suppression from kinematics

[JHEP 06 (2021) 093]



$\pi^+$  momentum range: 15—45 GeV/c

# Signal sensitivity: results (2021–22 data)



$$N_K = \frac{N_{\pi\pi} D_0}{\mathcal{B}_{\pi\pi} A_{\pi\pi}}$$

$$\mathcal{B}_{SES} = \frac{1}{N_K \varepsilon_{RV} \varepsilon_{trig} A_{\pi\nu\bar{\nu}}}$$

- Display integrals (15–45 GeV/c, 2021+22) for summary tables.
- \* Acceptances evaluated at 0 intensity.

$N_{\pi\pi}$	Normalisation $K^+ \rightarrow \pi^+ \pi^0$	$2.0 \times 10^8$
$A_{\pi\pi}$	Normalisation acceptance	$(13.410 \pm 0.005)\%$
$N_K$	Effective $K^+$ decays	$2.9 \times 10^{12}$
$A_{\pi\nu\bar{\nu}}$	Signal acceptance	$(7.6 \pm 0.2)\%$
$\varepsilon_{trig}$	Trigger efficiency	$(85.9 \pm 1.4)\%$
$\varepsilon_{RV}$	Random veto efficiency	$(63.6 \pm 0.6)\%$
$\mathcal{B}_{SES}$	Single event sensitivity	$(0.84 \pm 0.03) \times 10^{-11}$

$$N_{\pi\nu\bar{\nu}}^{exp} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}}$$

Assuming  $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$ :

2021–22:  $N_{\pi\nu\bar{\nu}} = 10.00 \pm 0.34$

c.f. 2016–18 :  $N_{\pi\nu\bar{\nu}} = 10.01 \pm 0.42$




Double expected signal by including 2021–22 data.

- Significant improvement in SES uncertainty:
  - old: 6.3% → new: 3.5%. Due to:
    - trigger efficiency cancellations
    - improved procedures for evaluation of acceptances and  $\varepsilon_{RV}$

# Signal sensitivity: random veto

$\varepsilon_{RV}$  is independent of track momentum  
(related to additional activity only)

$$N_{\pi\nu\bar{\nu}}^{exp}(p_i) = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}(p_i)} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{\pi\pi}} \frac{A_{\pi\nu\bar{\nu}}(p_i)}{A_{\pi\pi}(p_i)} D_0 N_{\pi\pi}(p_i) \varepsilon_{trig}(p_i) \varepsilon_{RV}$$




- $\varepsilon_{RV}$  = Random Veto Efficiency:
- $1 - \varepsilon_{RV}$  = Probability of rejecting a signal event due to additional activity.
- Balance:
  - Strict vetos  $\Rightarrow$  lower efficiency
  - Loose vetos  $\Rightarrow$  higher background
- Operational intensity higher but re-tuning vetos means  $\varepsilon_{RV}$  is comparable:

$$\varepsilon_{RV}(\text{new}, \overline{\lambda_{21-22}}) \approx 600 \text{ MHz} = (63.6 \pm 0.6)\%$$

$$\varepsilon_{RV}(\text{old}, \overline{\lambda_{2018}}) \approx 400 \text{ MHz} = (66 \pm 1)\%$$

# Optimum NA62 intensity

## Selected signal yield vs intensity




- Saturation of expected signal yield with intensity. Mainly due to:
  - Paralyzable effects from TDAQ dead time and trigger veto windows.
  - Offline selection, due to veto conditions.
- Main sources of uncertainty for model:
  - Online time-dependent mis-calibrations.
  - Fit uncertainty.
- **From August 2023 operate at optimal intensity (~75% of full) to maximise  $\pi\nu\nu$ sensitivity**
  - Maximise signal yield
  - lower expected background
  - Higher DAQ efficiency

Studies of 2021—22 data at high intensity were crucial to establish optimal intensity

# Upstream background mechanism

- A kaon decays upstream the fiducial decay region
- Only a  $\pi^+$  enters the fiducial decay region
- There is an in-time pileup beam particle (in GTK)
- The upstream  $\pi^+$  is scattered in the first STRAW chamber, and a fake vertex in the fiducial decay region is reconstructed



# Upstream background evaluation



$$N_{bg} = \sum_i N_i f_{cda} P_i^{match}$$

Upstream Reference Sample:  
signal selection but invert CDA cut (CDA>4mm)

Scaling factor : bad cda  $\rightarrow$  good cda

Probability to pass  $K^+ - \pi^+$  matching


$N$   
 $f_{cda}$   
 $P_{match}$

Calculate using bins (i) of  $(\Delta T_+, N_{GTK})$   
[Updated to fully data-driven procedure]

$$N = 51 \quad f_{CDA} = 0.20 \pm 0.03 \quad \langle P_{match} \rangle = 73\%$$

$$N_{bg}(\text{Upstream}) = 7.4^{+2.1}_{-1.8}$$


- Upstream reference sample contains all known upstream mechanisms.
  - $N$  provides normalisation.
- $f_{CDA}$  depends only on geometry.
- $P_{match}$  depends on  $(\Delta T_+, N_{GTK})$ .



# Upstream background validation



- Invert & loosen upstream vetos to enrich with different mechanisms:
  - Interaction-enriched: Val1,2,7,8
  - Accidental-enriched: Val3,4,5,6,9,10.
- All independent.
- Expectations and observations are in good agreement.
- Number of events rejected by VetoCounter:
  - (i.e. events in signal region with associated VC signal)
  - $N_{exp}^{VC rej.} = 6.9 \pm 1.4$ ,  $N_{obs}^{VC rej.} = 9$
- VetoCounter is essential to control upstream background.



# Background expectations (2021-22 data)



## Backgrounds

$K^+ \rightarrow \pi^+ \pi^0 (\gamma)$	$0.83 \pm 0.05$
$K^+ \rightarrow \pi^+ \pi^0$	$0.76 \pm 0.04$
$K^+ \rightarrow \pi^+ \pi^0 \gamma$	$0.07 \pm 0.01$
$K^+ \rightarrow \mu^+ \nu (\gamma)$	$1.70 \pm 0.47$
$K^+ \rightarrow \mu^+ \nu$	$0.87 \pm 0.19$
$K^+ \rightarrow \mu^+ \nu \gamma$	$0.82 \pm 0.43$
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	$0.11 \pm 0.03$
$K^+ \rightarrow \pi^+ \pi^- e^+ \nu$	$0.89^{+0.34}_{-0.28}$
$K^+ \rightarrow \pi^0 \ell^+ \nu$	$< 0.001$
$K^+ \rightarrow \pi^+ \gamma \gamma$	$0.01 \pm 0.01$
Upstream	$7.4^{+2.1}_{-1.8}$
Total	$11.0^{+2.1}_{-1.9}$

## Signal Sensitivity

$$\mathcal{B}_{SES} = (0.84 \pm 0.03) \times 10^{-11}$$

$$N_{\pi\nu\bar{\nu}}^{SM,exp} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}}$$

Assuming  $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$ :

2021–22:  $N_{\pi\nu\bar{\nu}} = 10.00 \pm 0.34$

c.f. 2016–18 :  $N_{\pi\nu\bar{\nu}} = 10.01 \pm 0.42$



→ Expected signal doubled by including 2021–22 data

- $N_{\pi\nu\bar{\nu}}^{SM}$  per SPS burst:  $2.5 \times 10^{-5}$  in 2022
  - c.f.  $1.7 \times 10^{-5}$  in 2018. ⇒ signal yield increased by 50%
- Sensitivity for  $\text{BR} \sim \sqrt{S + B}/S$  similar but improved with respect to 2018 analysis, for same amount of data

# The new NA62 results: 2016–22 data



- Integrating 2016–22 data:  $N_{bg} = 18^{+3}_{-2}$ ,  $N_{obs} = 51$ .
- Background-only hypothesis p-value =  $2 \times 10^{-7} \Rightarrow$  significance  $Z > 5$



$$\mathcal{B}_{16-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (13.0^{+3.3}_{-2.9}) \times 10^{-11} = (13.0^{(+3.0)}_{(-2.7)}{}_{stat}^{(+1.3)}{}_{syst}) \times 10^{-11}$$

# Results in context



BNL E787/E949 experiment  
[Phys. Rev. D 79 (2009) 092004]


$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-18} = (10.6_{-3.5}^{+4.1}) \times 10^{-11}$$

[JHEP 06 (2021) 093]

$$\mathcal{B}_{\pi\nu\bar{\nu}}^{21-22} = (16.0_{-4.5}^{+5.0}) \times 10^{-11}$$

$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-22} = (13.0_{-2.9}^{+3.3}) \times 10^{-11}$$

- NA62 results are consistent
- Central value moved up (now 1.5–1.7 $\sigma$  above SM)
- Fractional uncertainty decreased: 40% to 25%
- Bkg-only hypothesis rejected with significance  $Z>5$




# Results in context



- Fractional uncertainty: 25%
- Bkg-only hypothesis rejected with significance  $Z>5$
- **Observation of the  $K^+ \rightarrow \pi^+ \nu\bar{\nu}$  decay with BR consistent with SM prediction, within  $1.7\sigma$** 
  - Need full NA62 data-set to clarify SM agreement or tension

$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-22} = (13.0^{+3.3}_{-2.9}) \times 10^{-11}$$

$2\sigma$  range :  $[7.4 - 19.7] \times 10^{-11}$




# The NA62 published results






1. First detection of a tagged neutrino in the NA62 experiment, arXiv: 2412.04033, submitted to Phys. Lett. B.
2. First search for  $K^+ \rightarrow \pi^0 \pi^+ \mu^- \nu$  decays, Phys. Lett. B 859 (2024) 139122.
3. Search for leptonic decays of the dark photon at NA62, Phys. Rev. Lett. 133 (2024) 111802.
4. Measurement of the  $K^+ \rightarrow \pi^+ \gamma\gamma$  decay, Phys. Lett. B 850 (2024) 138513.
5. Search for  $K^+$  decays into the  $\pi^+ e^+ e^- e^-$  final state, Phys. Lett. B 846 (2023) 138193.
6. A study of the  $K^+ \rightarrow \pi^0 e^+ \nu\gamma$  decay, JHEP 09 (2023) 040.
7. Search for dark photon decays to  $\mu^+ \mu^-$  at NA62, JHEP 09 (2023) 035.
8. A search for the  $K^+ \rightarrow \mu^- \nu e^+ e^+$  decay, Phys. Lett. B 838 (2023) 137679.
9. A measurement of the  $K^+ \rightarrow \pi^+ \mu^+ \mu^-$  decay, JHEP 11 (2022) 011.
10. Searches for lepton number violating  $K^+ \rightarrow \pi^-(\pi^0) e^+ e^-$  decays, Phys. Lett. B 830 (2022) 137172.
11. Search for Lepton Number and Flavor Violation in  $K^+$  and  $\pi^0$  Decays, Phys. Rev. Lett. 127 (2021) 131802.
12. Measurement of the very rare  $K^+ \rightarrow \pi^+ \bar{\nu} \nu$  decay, JHEP 06 (2021) 093.
13. Search for  $K^+$  decays to a muon and invisible particles, Phys. Lett. B 816 (2021) 136259.
14. Search for a feebly interacting particle X in the decay  $K^+ \rightarrow \pi^+ X$ , JHEP 03, (2021) 058.
15. Search for  $\pi^0$  decays to invisible particles, JHEP 02, (2021) 201.
16. An investigation of the very rare  $K^+ \rightarrow \pi^+ \bar{\nu} \nu$  decay, JHEP 11 (2020) 042.
17. Search for heavy neutral lepton production in  $K^+$  decays to positrons, Phys. Lett. B 807 (2020) 135599.
18. Searches for lepton number violating  $K^+$  decays, Phys. Lett. B 797 (2019) 134794.
19. Search for production of an invisible dark photon in  $\pi^0$  decays, JHEP 1905 (2019) 182.
20. First search of  $K^+ \rightarrow \pi^+ \bar{\nu} \nu$  using the decay-in-flight technique, Phys. Lett. B 791 (2019) 156.
21. Search for heavy neutral lepton production in  $K^+$  decays, Phys. Lett. B 778 (2018) 137.

# Supplemental



# The story so far: $K^+ \rightarrow \pi^+\nu\bar{\nu}$ with 2016–18 data



Data-taking year	[Reference]	$N_{bg}$	$N_{\pi\nu\bar{\nu}}^{SM,exp}$	$N_{obs}$
2016	[PLB 791 (2019) 156]	$0.152^{+0.093}_{-0.035}$	$0.267 \pm 0.020$	1
2017	[JHEP 11 (2020) 042]	$1.46 \pm 0.33$	$2.16 \pm 0.13$	2
2018	[JHEP 06 (2021) 093]	$5.42^{+0.99}_{-0.75}$	$7.58 \pm 0.40$	17
2016–18	[JHEP 06 (2021) 093]	$7.03^{+1.05}_{-0.82}$	$10.01 \pm 0.42$	20

$N_{\pi\nu\bar{\nu}}^{SM,exp}$  assumes:  
 $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$


Statistical combination:  $\mathcal{B}(K^+ \rightarrow \pi^+\nu\bar{\nu}) = (10.6^{+4.0}_{-3.4}|_{stat} \pm 0.9|_{syst}) \times 10^{-11} @ 68\% CL$

Background-only hypothesis:  $p = 3.4 \times 10^{-4} \Rightarrow \text{significance} = 3.4\sigma$

# Backgrounds from kinematic tails




$K^+ \rightarrow \pi^+ \pi^0(\gamma)$   
 control sample of  $K^+ \rightarrow \pi^+ \pi^0$  events with  
 $\pi^0 \rightarrow \gamma\gamma$  and 2 photons detected in LKr:




$$N_{bg}(K^+ \rightarrow \pi^+ \pi^0(\gamma)) = 0.83 \pm 0.05$$

$K^+ \rightarrow \mu^+ \nu$   
 control sample of  $K^+ \rightarrow \mu^+ \nu$  events  
 with RICH PID= $\pi^+$  and Calo PID= $\mu^+$ :



- <1% contribution from  $K^+ \rightarrow \mu^+ \nu$  followed by  $\mu^+ \rightarrow e^+ \nu\nu$

$$N_{bg}(K^+ \rightarrow \mu^+ \nu) = 0.9 \pm 0.2$$





- Use MC to measure  $f_{tail}$ :

$$N_{bg}(K^+ \rightarrow \pi^+ \pi^+ \pi^-) = 0.11 \pm 0.03$$

# Radiative decays: $K^+ \rightarrow \pi^+\pi^0\gamma$ & $K^+ \rightarrow \mu^+\nu\gamma$





- $K^+ \rightarrow \pi^+\pi^0\gamma$  : included with “kinematic tails” estimation.
  - Suppression: photon vetos, rejection with additional  $\gamma$  is 30x stronger.
  - Estimation: MC + measured single photon rejection efficiency :  $N_{bg}(K^+ \rightarrow \pi^+\pi^0\gamma) = 0.07 \pm 0.01$
  - Validation:  $m_{miss}^2$  control regions (CR1,2 - see later)
- $K^+ \rightarrow \mu^+\nu\gamma$  : not included in “kinematic tails” estimation if  $\gamma$  overlaps  $\mu^+$  at LKr (leading to misID as  $\pi^+$ )
  - Suppression: based on  $(P_K - P_\mu - P_\gamma)^2$  and  $E_\gamma$  with  $\gamma$  = LKr cluster (mis)associated to muon.
    - Necessary for 2021–22 data, since Calorimetric PID degraded at higher intensities.
  - Estimation: min. Bias data control sample with signal in MUV3 :  $N_{bg}(K^+ \rightarrow \mu^+\nu\gamma) = 0.8 \pm 0.4$
  - Validation: data sample without  $K^+ \rightarrow \mu^+\nu\gamma$  veto and PID = “less pion-like” (Calo BDT bins below  $\pi^+$  bin).



# Background mechanism: $K^+ \rightarrow \mu^+\nu\gamma$

- $K^+ \rightarrow \mu^+\nu\gamma$  decay with fairly energetic photon ( $E_\gamma > 5\text{ GeV}$ ) and high momentum  $\mu^+$  ( $p \gtrsim 35\text{ GeV}/c$ ).
- $\gamma$  and  $\mu^+$  hit LKr together and are misidentified as a  $\pi^+$ .
- No rejection power from photon vetos (LKr  $\gamma$  cluster associated to track).
- Additional  $\gamma$  naturally shifts  $m_{miss}^2 = (P_K - P_\pi)^2$  towards higher values (i.e. towards signal regions).




# $K^+ \rightarrow \mu^+\nu\gamma$ Background




- Kinematically select  $K^+ \rightarrow \mu^+\nu\gamma$  events:  $m_{miss}^2(K_{\mu 2\gamma}) = (P_K - P_\mu - P_\gamma)^2$ 
  - $P_K$ : 4-momentum of  $K^+$  from GTK (as normal)
  - $P_\mu$ : 4-momentum of track with  $\mu^+$  mass hypothesis.
  - $P_\gamma$ : reconstructed from energy and position of LKr cluster (and position of  $K^+ - \mu^+$  vertex).


Validation: data sample with PID = “less pion-like” (Calo BDT bins below  $\pi^+$  bin).



Evaluate background expectation using  $\nu\nu\gamma$  control sample from MinimumBias trigger, not applying Calorimetric BDT classifier and MUV3 signal:



- Before  $K^+ \rightarrow \mu^+\nu\gamma$  veto: found excess of events at  $p > 35 \text{ GeV}/c$  in Region 2 relative to 2016–18 data.
- Additional background identified and studied in data control samples & MC.
- $K^+ \rightarrow \mu^+\nu\gamma$  veto added to selection criteria for final analysis.



# Upstream background evaluation



$$N_{bg} = \sum_i N_i f_{cda} P_i^{match}$$

Upstream Reference Sample:  
signal selection but invert CDA cut (CDA>4mm)

Scaling factor : bad cda  $\rightarrow$  good cda


Probability to pass  $K^+ - \pi^+$  matching

Calculate using bins (i) of  $(\Delta T_+, N_{GTK})$   
[Updated to fully data-driven procedure]

$$N = 51 \quad f_{CDA} = 0.20 \pm 0.03 \quad \langle P_{match} \rangle = 73\%$$

$$N_{bg}(Upstream) = 7.4^{+2.1}_{-1.8}$$



- Upstream reference sample contains all known upstream mechanisms.
  - $N$  provides normalisation.
- $f_{CDA}$  depends only on geometry.
- $P_{match}$  depends on  $(\Delta T_+, N_{GTK})$ .



# Control regions: 2021–22 Data




2021–22 data




- Good agreement in control regions validates background expectations.

# Signal regions: 2021–22 Data



1D projection with differential background predictions  
& SM signal expectation [not a fit]:



Expected SM signal:  $N_{\pi\nu\bar{\nu}}^{\text{SM}} \approx 10$


Expected background:  $N_{bg} = 11.0^{+2.1}_{-1.9}$

Observed:  $N_{obs} = 31$

# Results: 2021–22 Data




- Measure  $\mathcal{B}_{\pi\nu\bar{\nu}}$  and 68% ( $1\sigma$ ) confidence interval using a profile likelihood ratio test statistic  $q(\theta)$



$$\mathcal{B}_{21-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (16.0^{+5.0}_{-4.5}) \times 10^{-11} = (16.0^{(+4.8)}_{(-4.2)}{}_{stat}^{(+1.4)}{}_{syst}) \times 10^{-11}$$

- Use 6 (momentum bin) categories  
After fit (use measured BR)



# Results in context: the long story of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$



- Experimental measurements:
  - Camerini et al. [[PRL 23 \(1969\) 326-329](#)]
  - Klems et al. [[PRD 4 \(1971\) 66-80](#)]
  - Ljung et al. [[PRD 8 \(1973\) 1307-1330](#)]
  - Cable et al. [[PRD 8 \(1973\) 3807-3812](#)]
  - Asano et al. [[PLB 107 \(1981\) 159](#)]
- E787 :
  - [[PRL 64 \(1990\) 21-24](#)]
  - [[PRL 70 \(1993\) 2521-2524](#)]
  - [[PRL 76 \(1996\) 1421-1424](#)]
  - [[PRL 79 \(1997\) 2204-2207](#)]
  - [[PRL 84 \(2000\) 3768-3770](#)]
  - [[PRL 88 \(2002\) 041803](#)]
- E949 (+E787)
  - [[PRL 93 \(2004\) 031801](#)]
  - [[PRL 101 \(2008\) 191802](#)]
- NA62:
  - 2016 data: [[PLB 791 \(2019\) 156](#)]
  - 2016+17 data: [[JHEP 11 \(2020\) 042](#)]
  - 2016–18 data: [[JHEP 06 \(2021\) 093](#)]
  - 2016–22 data : this result.
- Theory:
  - [[Phys. Rev. 163 \(1967\) 1430-1440](#)]
  - [[PRD 10 \(1974\) 897](#)]
  - [[Prog.Theor.Phys. 65 \(1981\)](#)]
  - [[PLB 133 \(1983\) 443-448](#)]
  - [[PLB 192 \(1987\) 201-206](#)]
  - [[Nucl.Phys.B 304 \(1988\) 205-235](#)]
  - [[PRD 54 \(1996\) 6782-6789](#)]
  - [[PRD 76 \(2007\) 034017](#)]
  - [[PRD 78 \(2008\) 034006](#)]
  - [[PRD 83 \(2011\) 034030](#)]
  - [[JHEP 11 \(2015\) 033](#)]
  - [[JHEP 09 \(2022\) 148](#)]

