

The Cabibbo Angle from Inclusive T Decays



Francesco Sanfilippo – INFN Roma Tre

XXXVIII Convegno Nazionale di Fisica Teorica, 22/05/2025




Based on

[A. evangelista et al.
Phys Rev. D 108, 074513 (2023)]

EDITORS' SUGGESTION

Inclusive hadronic decay rate of the τ lepton from lattice QCD


The authors express the inclusive hadronic decay rate of the tau lepton as an integral over the spectral density of the two-point correlator of the weak $V - A$ hadronic current which they compute fully nonperturbatively in lattice QCD. In a lattice QCD computation with all systematic errors except for isospin breaking effects under control, they then obtain the CKM matrix element V_{ud} with subpercent errors showing that their nonperturbative method can become a viable alternative to superallowed nuclear beta decays for obtaining V_{ud} .



Flavor in the SM: the CKM matrix

In the SM, the charged change of flavors occur through the exchange of a W boson

$$\mathcal{L}_w = -\frac{g_w}{\sqrt{2}} (\bar{u}, \bar{c}, \bar{t}) \gamma^\mu P_L W_\mu^+ V_{\text{CKM}} \begin{pmatrix} d \\ s \\ b \end{pmatrix} + \text{h.c.}, \quad V_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$



Unitary matrix connecting up-like and down-like quarks

Hierarchy of amplitude between different generation

$$\underbrace{\begin{pmatrix} |V_{ud}| & & \\ & |V_{cs}| & \\ & & |V_{tb}| \end{pmatrix}}_{\mathcal{O}(1)} \gg \underbrace{\begin{pmatrix} & |V_{us}| & \\ |V_{cd}| & & \end{pmatrix}}_{\mathcal{O}(10^{-1})} \gg \underbrace{\begin{pmatrix} & & |V_{ub}| \\ & |V_{ts}| & |V_{cb}| \\ |V_{td}| & & \end{pmatrix}}_{\mathcal{O}(10^{-2}-10^{-3})}$$

We focus on the first row of the CKM matrix

First row unitarity

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1 \quad (?)$$

$|V_{ud}|$ from superallowed nuclear beta decays: 0.03%


$|V_{ud}|$ from semileptonic $\pi \rightarrow \pi^0 e \nu_e$ beta decay: 0.3%

$|V_{ub}|$ is very small
and we can drop it

$|V_{us}|$ from semileptonic decay $K \rightarrow \pi \ell \nu$: 0.2%

$\frac{|V_{us}|}{|V_{ud}|}$ from the ratio of semileptonic decay
 $K/\pi \rightarrow \ell \nu(\gamma)$: 0.3%

Tension between the value of $|V_{us}|$ from
leptonic and semileptonic decays vs. $|V_{us}^{uni}|$




Usefulness of τ decay

Three leptons are known experimentally, their measured masses is:



$$m_e = 0.5110 \text{ KeV}, \quad m_\mu = 105.66 \text{ MeV}, \quad m_\tau = 1.776 \text{ GeV}$$

The lightest charged hadron is the **pion**, with mass: $m_\pi^\pm = 139.57 \text{ MeV}$

→ τ is unique among leptons, as **it can decay into hadrons**



Two hadronic decay channels




Studying the ud/us channels one can determine $V_{ud/us}$

Comparison with $K_{\ell 2}, K_{\ell 3}$ determinations shows tension

τ determinations also in tension with unitarity constraints

Tension is more pronounced for inclusive process $\tau \rightarrow X_{us}$



Inclusive process, $\mathcal{B}_{X_{ud/us}} \equiv \Gamma[\tau \rightarrow X_{ud/us} \nu_\tau] / \Gamma_{\text{tot}}$

Sum of exclusive processes for a given channel

$\tau \rightarrow X_{us}$ channel:

Branching fraction	HFLAV 2021 fit (%)
$K^- \nu_\tau$	0.6957 ± 0.0096
$K^- \pi^0 \nu_\tau$	0.4322 ± 0.0148
$K^- 2\pi^0 \nu_\tau$ (ex. K^0)	0.0634 ± 0.0219
$K^- 3\pi^0 \nu_\tau$ (ex. K^0, η)	0.0465 ± 0.0213
$\pi^- \bar{K}^0 \nu_\tau$	0.8375 ± 0.0139
$\pi^- \bar{K}^0 \pi^0 \nu_\tau$	0.3810 ± 0.0129
$\pi^- \bar{K}^0 2\pi^0 \nu_\tau$ (ex. K^0)	0.0234 ± 0.0231
$\bar{K}^0 h^- h^- h^+ \nu_\tau$	0.0222 ± 0.0202
$K^- \eta \nu_\tau$	0.0155 ± 0.0008
$K^- \pi^0 \eta \nu_\tau$	0.0048 ± 0.0012
$\pi^- \bar{K}^0 \eta \nu_\tau$	0.0094 ± 0.0015
$K^- \omega \nu_\tau$	0.0410 ± 0.0092
$K^- \phi (K^+ K^-) \nu_\tau$	0.0022 ± 0.0008
$K^- \phi (K_S^0 K_L^0) \nu_\tau$	0.0015 ± 0.0006
$K^- \pi^- \pi^+ \nu_\tau$ (ex. K^0, ω)	0.2924 ± 0.0068
$K^- \pi^- \pi^+ \pi^0 \nu_\tau$ (ex. K^0, ω, η)	0.0387 ± 0.0142
$K^- 2\pi^- 2\pi^+ \nu_\tau$ (ex. K^0)	0.0001 ± 0.0001
$K^- 2\pi^- 2\pi^+ \pi^0 \nu_\tau$ (ex. K^0)	0.0001 ± 0.0001
$X_s^- \nu_\tau$	2.9076 ± 0.0478

Accuracy: 1.7%

$\tau \rightarrow X_{ud}$ channel:

In the past, obtained by difference

$$\mathcal{B}_{X_{ud}} = 1 - \mathcal{B}_{X_{us}} - \mathcal{B}_e - \mathcal{B}_\mu$$

with $\mathcal{B}_{e/\mu}$ branching to electrons/muons

Nowadays directly measured

$$\mathcal{B}_{X_{ud}} = 0.6183 \pm 0.0010$$

according to HFLAV average

Weak Hamiltonian Effective theory



$$H_{ud}^{\text{eff}}(x) = \frac{G_F}{\sqrt{2}} V_{ud} \sqrt{S_{\text{EW}}} \underbrace{\bar{\nu}_\tau(x) \gamma^\alpha (1 - \gamma^5) \tau(x)}_{J_{\nu_\tau \tau}^\alpha(x)} \times \underbrace{\bar{d}(x) \gamma^\alpha (1 - \gamma^5) u(x)}_{(J_{ud}^\alpha)^\dagger(x)} + \text{h.c.}$$

Factorization of the amplitude in leptonic and **nonperturbative** hadronic parts

$$\mathcal{A}(\tau \rightarrow X_{ud} \nu_\tau) = \frac{G_F}{\sqrt{2}} V_{ud} \sqrt{S_{\text{EW}}} \langle \nu_\tau | J_{\nu_\tau \tau}^\alpha(0) | \tau \rangle \langle X_{ud} | J_{ud}^\alpha(0)^\dagger | 0 \rangle$$

From matrix element to decay width

$$\sum X_{ud} \left[\text{Diagram 1} \right]^2 = \frac{1}{2m_\tau} 2\text{Im} \left[\text{Diagram 2} \right]$$

All that has been done so far is an application of the **optical theorem**

$$\Gamma[\tau \rightarrow X_{ud}\nu_\tau] = \frac{1}{2m_\tau} 2\text{Im}[\Gamma_{\tau\tau}]$$

where $\Gamma_{\tau\tau} = \langle \tau | T | \tau \rangle$ is the forward amplitude $S = 1 + iT$ related to ud states

The hadronic tensor

Squaring $\mathcal{A}(\tau \rightarrow X_{ud}\nu_\tau)$ and summing over all X_{ud} states:

$$|\mathcal{A}|^2 = \sum_{X_{ud}} |\mathcal{A}(\tau \rightarrow X_{ud}\nu_\tau)|^2 = \left(\frac{G_F}{\sqrt{2}}\right)^2 |V_{ud}|^2 S_{EW} L^{\alpha\beta}(p_\tau, p_\nu) \rho_{ud}^{\alpha\beta}(q)$$

with p_τ, p_ν being the τ, ν_τ four-momentum, and $q = p_\tau - p_\nu$.

Leptonic tensor $L^{\alpha\beta}(p_\tau, p_\nu)$ obtained averaging/summing over τ/ν_τ polarizations:

$$L^{\alpha\beta}(p_\tau, p_\nu) = 4(p_\tau^\alpha p_\nu^\beta + p_\tau^\beta p_\nu^\alpha - g^{\alpha\beta} p_\tau \cdot p_\nu - i\varepsilon^{\alpha\beta\rho\sigma} p_{\tau\rho} p_{\nu\sigma})$$

Missing ingredient: the hadronic tensor

$$\rho_{ud}^{\alpha\beta}(q) \equiv \sum_{X_{ud}} \langle 0 | J_{ud}^\alpha(0) | X_{ud}(q) \rangle \langle X_{ud}(q) | J_{ud}^\beta(0)^\dagger | 0 \rangle$$


Determining the Hadronic Tensor

In Minkowsky spacetime, from the spectral representation (Fourier transform):

$$C^{\alpha\beta}(t, q) \stackrel{t \geq 0}{=} \int_0^\infty \frac{dE}{2\pi} e^{-iEt} \rho_{ud}^{\alpha\beta}(E, q)$$

...of the current-current correlation function in time-momentum representation:

$$C^{\alpha\beta}(t, q) = \int d^3x e^{-iq \cdot x} \langle 0 | T \left(J_{ud}^\alpha(-it, x) J_{ud}^\beta(0)^\dagger \right) | 0 \rangle$$




- Nonperturbative quantity
- Severe disease with a bad prognosis
- Call a latticist!

Hadronic amplitudes in Minkowskian

Correlation function, e.g. J_A, J_B currents on state $|P\rangle$

$$C(t) \equiv \langle 0 | T \{ J_A(t) J_B(0) \} | P \rangle \stackrel{t \geq 0}{=} \sum_{n=0}^{\infty} C_n e^{-iE_n t} \quad (\text{Fourier transform})$$



Extracting Hadronic Amplitude $H(E)$
in **Minkowskian** continuum spacetime:

$$\rho(E) = i \lim_{T \rightarrow \infty} \int_0^T dt e^{iEt} C(t)$$

Inverse Fourier transform: a reliable horse

Hadronic amplitudes in Euclidean lattice

Analytic continuation to discrete Euclidean spacetime: $\tau = it$

$$C_e(\tau) \equiv C(-i\tau) = \sum_{n=0}^{\infty} C_n e^{-E_n \tau}$$

Laplace transform



Extracting Hadronic Amplitude

in discrete **Euclidean** spacetime of length T:

$$\rho^T(E) = \int_0^T d\tau e^{E\tau} C_e(\tau)$$

Inverse Laplace transform!??

A wild and unreliable horse!

Analytic continuation issues

In Euclidean spacetime, it holds true that

$$C^{\alpha\beta}(t, q) \stackrel{t \geq 0}{=} \int_0^\infty \frac{dE}{2\pi} e^{-Et} \rho_{ud}^{\alpha\beta}(E, q)$$

i.e the current-current correlator is the Laplace transform of the needed Hadronic tensor

But

One cannot naively invert this relation if the energy E lies above the minimal threshold E_0

This requires an inverse Laplace transform, **ill-posed problem on finite noisy dataset**

CFR Nazario's talk

Spectral representation

$$\rho^T(E) = \int_0^T dt e^{Et} C_E(t) = \int_0^T dt e^{Et} \sum_{n=0}^{\infty} C_n e^{-E_n t} = \sum_{n=0}^{\infty} C_n \frac{1 - e^{-(E_n - E)T}}{E_n - E}$$

Let us break down two energy regimes

$$E < E_0$$

$$\rho^T(E) \xrightarrow{T \rightarrow \infty} \sum_{n=0}^{\infty} \frac{C_n}{E_n - E}$$

GOOD!

$$E > E_0$$

$$\rho^T(E) \xrightarrow{T \rightarrow \infty} \sum_{n=0}^{\infty} \frac{C_n}{E_n - E} + \sum_{n}^{E_n < E} C_n \frac{e^{(E - E_n)T}}{E - E_n}$$

BAD!!!

- How to subtract the divergent part?
- Needs all C_n to recover the imaginary part!

Bypassing the determination of $\rho(E)$

Setting the external spatial momentum to zero we can simplify the needed correlators

$$\rho_{ud}^{00}(q) = q^2 \rho_L(q^2) , \quad \frac{1}{3} \sum_{i=1}^3 \rho_{ud}^{ii}(q) = q^2 \rho_T(q^2)$$

and rewrite the decay rate in terms of a **convolution integral** over energy

$$R_{ud}^{(\tau)} \propto \int_0^\infty dE \left[K_T \left(\frac{E}{m_\tau} \right) E^2 \rho_T(E^2) + K_L \left(\frac{E}{m_\tau} \right) E^2 \rho_L(E^2) \right]$$

having incorporated back the leptonic term into the kernels:

$$K_L(x) \equiv \frac{1}{x} (1 - x^2)^2 \theta(1 - x) , \quad K_T(x) \equiv (1 + 2x^2) K_L(x)$$

The explicit determination of the spectral density is avoided: only its **convolution** with K is needed

Inverse Laplace transform for $K(E)$

We don't need to determine the spectral densities, just its convolution with $K(E)$

$$R_{ud}^{(\tau)} = 12\pi S_{EW} \frac{|V_{ud}|^2}{m_\tau^3} \int_0^\infty dE \left[K_T \left(\frac{E}{m_\tau} \right) E^2 \rho_T(E^2) + K_L \left(\frac{E}{m_\tau} \right) E^2 \rho_L(E^2) \right]$$


Both kernels can be expressed in terms of an approximated inverse Laplace transform

$$R_{ud}^{(\tau)} \propto \sum_{I \in \{L, T\}} \int_0^\infty dE K_I \left(\frac{E}{m_\tau} \right) E^2 \rho_I(E^2) = \sum_{I \in \{L, T\}} \int_0^\infty dE \sum_{n=1}^N g_{n,I} e^{-nE} E^2 \rho_I(E^2)$$

with coefficients $g_{n,I}$ determined imposing minimal L^2 distance

...will this be numerically stable...?

...not quite...



- This is due to the θ function, needed to cut out energy larger than m_τ
- Using this coefficients introduces uncontrolled error in the rate
- We need to smooth out the sharp θ function


Smoothing out the kernels

HLT framework to solve the inverse Laplace transform in the context of Lattice simulations
[M.Hansen, A.Lupo, N.Tantalo, PRD 96 (2017)]


Smoothing the kernel defining the rate has been tested in the inclusive semilptonic decays
[P.Gambino et al., PRL 125 (2020)]

We introduce a smeared kernel in terms of a smoothed θ , and compute the rate

$$\theta(x) \rightarrow \Theta_\sigma(x) \equiv \frac{1}{1 + e^{-x/\sigma}},$$




$$\lim_{\sigma \rightarrow 0} \Theta_\sigma(x) = \theta(x)$$




$$R_{ud}^{(\tau, \text{I})}(\sigma) = 12\pi S_{\text{EW}} \frac{|V_{ud}|^2}{m_\tau^3} \int_0^\infty dE K_{\text{I}}^\sigma \left(\frac{E}{m_\tau} \right) E^2 \rho_{\text{I}}(E^2)$$

Infinite volume/continuum extrapolations

It is convenient to carry out the extrapolations to the continuum at fixed σ



The infinite volume limit involves no power-law correction: $R_{ud}^{(\tau, \text{I})}(\sigma, \infty) - R_{ud}^{(\tau, \text{I})}(\sigma, L) \sim \mathcal{O}(L^{-\infty})$



The extrapolation to zero smoothing is done at the end, exploiting the asymptotic relation

$$R_{ud}^{(\tau, \text{I})}(\sigma) = R_{ud}^{(\tau, \text{I})} + C\sigma^4 + \mathcal{O}(\sigma^6)$$


Numerical setup

Four physical-point $N_f=2+1+1$ Wilson-clover-twisted mass ensembles

Lattice spacing: $a \in [0.057 \text{ fm} - 0.080 \text{ fm}]$ Iwasaki action

$L \sim 5.1 \text{ fm}$ and $L \sim 7.6 \text{ fm}$ to control Finite Size Effects


Automatic improved $\mathcal{O}(a)$ improvement of observables




ID	V/a^4	a fm	L fm	m_π GeV
B64	$64^3 \cdot 128$	0.07957(13)	5.09	0.1352(2)
B96	$96^3 \cdot 192$	0.07957(13)	7.64	0.1352(2)
C80	$80^3 \cdot 160$	0.06821(13)	5.46	0.1349(3)
D96	$96^3 \cdot 192$	0.05692(12)	5.46	0.1351(3)

Numerical aspects


- Multi GPU simulations on Leonardo
- ISCRA & EuroHPC competitive grants
- INFN contingent of computing time



Multigrid algorithms to accelerate solution of the Dirac Equation



Optimally implemented in QUDA library by community + hardware vendor effort




...in a fully GPU-ported suite of
~100k lines of C++17 code



Scaling up to hundreds of GPUs

HLT method stabilization

Naive minimization







HLT method ($\alpha = 4, r_{\max} = 4, \text{B64}$)



Stability analysis

Tuning of λ and tweaking the parameters $\{\alpha, E_{\min}, r_{\max}\}$ probes the stability of the reconstruction


$$d_I[g_I^\lambda] \equiv \sqrt{A_I^0[g_I^\lambda]/A_I^0[0]}$$



Comparison of different volumes


Systematic of FSE estimated by comparing two volumes

Observed difference added as an error







Continuum limit extrapolation

- Combination of two regularizations (OS and TM)
- Several fit ansatz variations




Extrapolating to vanishing smearing



Extrapolation to zero smearing well described by σ^4 asymptotic prediction

Comparison with the experiments



Using the average value $R_{ud}^{(\tau)}(\text{HFLAV}) = 3.471(7)$ we obtain $|V_{ud}| = 0.9752(39)$

Good agreement with $|V_{ud}| = 0.97373(31)$ from superallowed β -decay.

The us channel



PHYSICAL REVIEW LETTERS **132**, 261901 (2024)

Inclusive Hadronic Decay Rate of the τ Lepton from Lattice QCD: The $\bar{u}s$ Flavor Channel and the Cabibbo Angle



Constantia Alexandrou,^{1,2} Simone Bacchio,² Alessandro De Santis,³ Antonio Evangelista,³ Jacob Finkenrath,⁴
Roberto Frezzotti,³ Giuseppe Gagliardi,⁵ Marco Garofalo,⁶ Bartosz Kostrzewa,⁷ Vittorio Lubicz,⁸ Simone Romiti,⁶
Francesco Sanfilippo,⁵ Silvano Simula,⁵ Nazario Tantalo¹⁰,³ Carsten Urbach,⁶ and Urs Wenger⁹

(Extended Twisted Mass Collaboration)

$$R_{us}(\sigma) = 12\pi S_{EW} |V_{ud}|^2 \int_0^\infty \frac{dE E^2}{m_\tau^3} \left\{ K_T^\sigma \left(\frac{E}{m_\tau} \right) \rho_T(E^2) + K_L^\sigma \left(\frac{E}{m_\tau} \right) \rho_L(E^2) \right\}$$




The us channel



Alternative determination, not (yet) competitive with $K_{\ell 3}$ but independent

Some tension is visible indeed...

Perspective: Isospin Breaking



Needs to include the **photon exchange** between hadrons and tau

A few percent effect, but very important to perform effective prediction

Work in progress...

Thank
you!