

The Cabibbo Angle from Inclusive τ Decays

Thursday, 22 May 2025 15:40 (15 minutes)

The inclusive hadronic decays of the τ lepton offer an alternative method for extracting the CKM matrix elements V_{ud} and V_{us} . In this talk, I will discuss recent results from the ETM Collaboration on the inclusive hadronic decay rate of the τ , obtained in $N_f = 2 + 1 + 1$ QCD using the novel HLT method. This approach circumvents the well-known inverse Laplace transform problem that hinders this calculation, allowing us to obtain first-principles results without relying on the operator-product expansion or perturbative QCD. Apart from isospin-breaking (IB) effects, all systematic uncertainties are under control. In the $\bar{u}s$ channel we obtain $|V_{us}|_{\tau\text{-latt-incl}} = 0.2189(7)_{\text{th}}(18)_{\text{exp}}$, which reveals a 3σ tension with purely hadronic determinations of $|V_{us}|$. Since this tension can no longer be attributed to the OPE approximation, it prompts a closer examination of experimental uncertainties and highlights the importance of determining IB corrections from first principles. I will briefly present an update on the ongoing status of our calculation of these corrections.

Primary authors: SANFILIPPO, Francesco (Istituto Nazionale di Fisica Nucleare); GAGLIARDI, Giuseppe (Istituto Nazionale di Fisica Nucleare)

Presenter: SANFILIPPO, Francesco (Istituto Nazionale di Fisica Nucleare)

Session Classification: Hadron/Flavor/Lattice