

## Precision study of $K^{+-}$ into $\pi^{+-} \pi^0 \pi^0$ and $K^{+-}$ into $\pi^{+-} \pi^+ \pi^-$ Dalitz plot distributions by NA48/2.

*Tuesday, 22 May 2007 17:50 (20 minutes)*

The NA48/2 experiment at the CERN SPS has collected an unprecedented sample of 3-pion decays of charged kaons. The high statistics and the good resolution of the detectors allow a unique investigation of the detailed phase space distributions of these decays. The effects of final state pion rescattering observed in the Dalitz plot distribution of the  $K^{+-}$  into  $\pi^{+-} \pi^0 \pi^0$  decays turned out to be a powerful tool for extraction of the S-wave pion-pion scattering lengths. The recent results obtained using a number of different theoretical approaches will be discussed, together with future prospects. The large statistics also allowed a precise measurement of the Dalitz plot slope parameters for the  $K^{+-}$  into  $\pi^{+-} \pi^+ \pi^-$  decays, which will be discussed.

**Primary author:** Dr GOUDZOVSKI, Evgueni (Scuola Normale Superiore and INFN, Pisa)

**Presenter:** Dr GOUDZOVSKI, Evgueni (Scuola Normale Superiore and INFN, Pisa)

**Session Classification:** Session III

**Track Classification:** Low energy QCD