

# KLOE measurements of the BR(Ks into gamma gamma) and direct search for Ks into e+e-

Tuesday, 22 May 2007 16:10 (20 minutes)

A precise measurement of the  $K_S \rightarrow \gamma\gamma$  rate is an important test of Chiral Perturbation Theory predictions. The decay amplitude can be evaluated at the leading  $p^4$  order providing an estimate of the BR for this decay of  $(2.1 \pm 0.1) \times 10^{-6}$ . The latest experimental determination of  $BR(K_S \rightarrow \gamma\gamma)$  is a precise measurement from NA48,  $(2.78 \pm 0.07) \times 10^{-6}$ , which differs from  $\chi$ PT  $p^4$  prediction of about 30%. This seems to indicate the presence of important contributions from higher order corrections. KLOE analysis on  $1.6 \text{ fb}^{-1}$  of data acquired during years 2001-2002 and 2004-2005 benefits from the tagging technique, which allows for the first time this decay to be identified with a pure  $K_S$  "beam", without the background from  $K_L \rightarrow \gamma\gamma$  decay, and with completely different systematics respect to fixed target experiments. Event counting is performed from a fit to the bidimensional distribution of the two-photon invariant mass versus the angle between photon momenta in the  $K_S$  rest frame. In this plane the best separation is achieved between the signal and the main source of background, which is represented by  $K_S \rightarrow \pi^0\pi^0$  events with two missing photons. The result of this analysis is presented, which is competitive with present measurements.

$K_S \rightarrow e^+e^-$  decay is a  $\Delta S = 1$  weak neutral current process. The Standard Model expectation for its BR is  $1.6 \times 10^{-15}$ , which has been evaluated by Chiral Perturbation Theory with 10% error. The best experimental limit on this decay, achieved by CPLEAR experiment, is  $BR < 1.4 \times 10^{-7}$  at 90% CL. We performed a direct search of  $K_S \rightarrow e^+e^-$  decay by analysing  $1.3 \text{ fb}^{-1}$  of data. The analysis exploits the excellent KLOE drift chamber momentum resolution to identify the signal through  $e^+e^-$  invariant mass reconstruction. Further background rejection comes from calorimeter particle identification, which is based on time of flight, shower longitudinal profile and  $E/p$ . The result of this search is presented, which improves on the previous experimental limit by a factor of 10.

**Primary author:** KLOE, Collaboration (INFN/LNF)

**Presenter:** Dr MARTINI, Matteo (LNF -INFN)

**Session Classification:** Session II

**Track Classification:** Non leptonic/ radiative decays